

Composite health behaviour classifier as the basis for targeted interventions and global comparisons in men's health

Ryan Kendrick Flannigan^{1,2}; John L. Oliffe³; Donald R. McCreary⁴; Nahid Punjani⁵; Khushabu Kasabwala¹; Nick Black⁶; Joe Rachert⁷; Larry S. Goldenberg²

¹Department of Urology, Weill Cornell Medicine, New York City, NY, United States; ²Department of Urologic Sciences, University of British Columbia, Vancouver, BC, Canada; ³School of Nursing, University of British Columbia, Vancouver, BC, Canada; ⁴Department of Psychology, Brock University, St. Catharines, ON, Canada; ⁵Division of Urology, Western University, London, ON, Canada; ⁶Intensions Consulting, Vancouver, British Columbia; ⁷Canadian Men's Health Foundation

Acknowledgements: Study supported by Canadian Men's Health Foundation; Public Health Agency of Canada.

Cite as: *Can Urol Assoc J* 2018 September 27; Epub ahead of print.

<http://dx.doi.org/10.5489/cuaj.5454>

Published online September 27, 2018

Abstract

Introduction: Lifestyle-related diseases are the leading cause of death among North American men. We evaluated health behaviours and their predictors that contribute to morbidity and mortality among Canadian men as a means to making recommendations for targeted interventions.

Methods: A cross-sectional analysis of Canadian men drawn from 5362 visitors to our online survey page was conducted. The current study sample of 2000 men (inclusion: male and >18 years; exclusion: incomplete surveys) were stratified to the 2016 Canadian census. The primary outcome was the number of unhealthy men classified using our Canadian Composite Classification of Health Behaviour (CCCHB) score. Secondary outcomes included the number of men with unhealthy exercise, diet, smoking, sleep, and alcohol intake, as well as socioeconomic and demographic factors associated with unhealthy behaviours to be used for targeting future interventions.

Results: Only 118/2000 (5.9%) men demonstrated 5/5 healthy behaviours, and 829 (41.5%) had 3/5 unhealthy behaviours; 391 (19.6%) men currently smoked, 773 (38.7%) demonstrated alcohol overuse, 1077 (53.9%) did not get optimal sleep (< 7 or ≥9 hours per night), 977 (48.9%) failed to exercise ≥150 minutes/week, and 1235 (61.8%) had an unhealthy diet. Multivariate analysis indicated that men with high school education were at increased risk of unhealthy

behaviours (odds ratio [OR] 1.58; 95% confidence interval [CI] 1.15–2.18; $p=0.005$), as were men living with relatives (OR 2.10; 95% CI 1.04–4.26; $p=0.039$), or with their partner and children (OR 1.34; 95% CI 1.02–1.76; $p=0.034$).

Conclusions: An overwhelming 41.5% of Canadian men had 3/5 unhealthy behaviours, affirming the need for targeted lifestyle interventions. Significant health inequities within vulnerable sub-groups of Canadian men were identified and may guide the content and delivery of future interventions.

Introduction

Health disparities and inequities within Canadian men's health are rarely articulated and poorly understood. Men die prematurely from preventable causes at an unprecedented rate, best reflected by a large potential years of life lost (PYLL) of 3,836 per 100,000.^{1,2} The etiology of increased mortality among men is unclear, though the evidence suggests that biological, environmental, psychological and behavioural factors are at play.

Men identify individually with masculine ideals and norms, which guide lifestyle and behaviour choices. Strong alignments to competitiveness and stoicism may heighten men's likelihood of participating in risky behaviours.^{3,4} Key health behaviours associated with mortality and morbidity have repeatedly identified lack of exercise, poor diets, smoking, alcohol overuse and poor sleep to be primary contributors⁵⁻¹⁶. Men demonstrate high rates of alcohol and tobacco use.¹⁷ Canadian men are less likely to consume a healthy diet¹⁸⁻²² resulting in high rates of obesity and predisposing them to lifestyle-related chronic diseases.²³ No accepted ideal diet has been identified; however, dietary components associated with negative health outcomes include those high in salt,²⁴ saturated fats,^{25,26} and refined sugar.²⁷⁻²⁹ Dietary choices associated with positive health outcomes include mono or polyunsaturated fats intake,³⁰ and 5 or more servings of fruits and/or vegetables per day.³¹⁻³³ Collectively, the impact of these health behaviours upon chronic disease is significant. Studies have identified compounding risks of mortality and morbidity with each added unhealthy behaviour.^{13,16} Studies have predicted that elimination of unhealthy behaviours would prevent 80% of heart disease, stroke, type 2 diabetes and 40% of cancers in the general population.^{34,35}

Demographic and socioeconomic status (SES) disparities have been identified to be strongly associated with increased morbidity and mortality.³⁶⁻⁴⁰ This association may, in part, be explained by poor health behaviours, lack of social support for minorities and a relative lack of resources among those with low SES. Identifying subgroups of men with dispirit demographics and SES contributing to poor health behaviours in Canada is important to identify those at greatest risk for morbidity and mortality.

An early step to preventing chronic illness and improving quality of life in men involves identifying behaviours as they relate to overall health, including men's health promotion practices (or lack thereof). The objective of the current study was to construct a composite behavioural classification (CCCHB) to quantitate morbidity and mortality associated health

behaviours among a representative sample of the Canadian male population, as a means to guiding recommendations for targeted interventions and global comparisons.

Methods

Institutional Research Board (IRB) approval was obtained from the University of British Columbia. Between April 20, 2017 and April 28, 2017, 5,362 participants were sourced from an online sample provider and screened to ensure they met inclusion criteria: men aged 19 or older, who were able to read French or English and residing in Canada. The exclusion criteria included an incomplete survey, providing non-differential responses, completing the survey significantly faster than average speed, and respondents not identifying as male. Weighted randomization was used to select 40,000 potential respondents from a large panel, who then received the initial survey invitation. Stratification was performed to ensure that the sample's composition reflected the relative distribution of the Canadian population by age and geography as determined by the 2016 Canadian Census data.⁴¹ The survey topic was not disclosed in the initial survey invitation, and only potential respondents who went to the survey introduction page were advised that the focus was on health behaviours (Figure 1). Informed consent was obtained prior to completing the survey. In total, 2000 men met study criteria and were included in the final analysis.

Demographic variables were collected including: age, minority status, employment status, household income, sexual orientation, province of residence, living arrangement, education level, and medical comorbidities.

The survey consisted of 5 distinct sections including: geographic sampling, demographic profiling, Men's Health literacy, Men's Health stigma, and Men's Health behaviours. The survey consisted of 94 questions dispersed among the 5 sections and was estimated to take respondents 15 minutes in total duration. Question types included Likert scales and multiple-choice designs.

Health behaviours

We assessed the established key health behaviours that are known to impact male morbidity and mortality. To do this, we used a priori evidence based thresholds to classify behaviours as being 'healthy' versus 'unhealthy'.

Smoking

Participants were asked about their cigarette smoking habits based upon questions adapted from the Canadian Tobacco Use Monitoring Survey (CTUMS)⁴². Men were classified as healthy if they were non-smokers or ex-smokers.

Alcohol

Participants were administered the validated Audit-C screening questionnaire⁴³ which functions with a sensitivity of 0.86 and specificity of 0.89 when used as a screening tool for alcohol overuse among men.⁴⁴ A score of less than 4 was considered healthy.

Sleeping

Participants were asked the duration of sleep obtained each night based upon the Canadian Health Measures Survey (CHMS).⁴⁵ Men sleeping 7 or 8 hours per night were classified as healthy in accordance with the National Sleep Foundation's updated sleep duration recommendations⁴⁶ and literature consensus that demonstrates optimal health outcomes associated with 7 hours sleep per night.⁴⁷⁻⁴⁹

Exercise

The validated Godin Leisure-time exercise questionnaire,⁵⁰ designed to assess exercise behaviour in the community,⁵¹ was administered to participants. Men achieving at least 150 minutes of moderate to strenuous exercise per week were classified as healthy, in accordance with the Canadian Physical Activity Guidelines for 18-64 year olds,⁵² 65+ year olds⁵³ and available literature.⁵⁴

Diet

Participants were asked about their weekly frequency of food consumption. Unhealthy foods were defined as those identified in the literature to be associated with negative health outcomes including foods high in salt,²⁴ saturated fats,^{25,26} and refined sugar.²⁷⁻²⁹ Healthy foods were defined as those associated with positive health outcomes and included high in mono or polyunsaturated fats,³⁰ and 5 or more servings of fruits and/or vegetables per day.³¹⁻³³ Lower frequencies of unhealthy food choices and higher frequencies of healthy food choices resulted in greater scores. Men with a composite score greater than 11 of a maximum score of 15 were classified as healthy. This questionnaire was adapted from the 'youcheck' tool, with the addition of a refined sugar component.⁵⁵

Total health behaviour classification

Men were classified as 'very healthy' if they demonstrated no unhealthy behaviours; 'healthy' if they had 1 unhealthy behaviour; 'borderline' if they had 2 unhealthy behaviours; and 'unhealthy' if they had 3-5 unhealthy behaviours. These categories were then dichotomized into a combined group of "very healthy, healthy, borderline" group and "unhealthy" group. This division was based upon previous work where 0-2 unhealthy behaviours has been used as the referent healthy group, validated by increased mortality among men and women with additional unhealthy behaviours.¹³

Statistical analysis

Descriptive analysis was performed to identify the number of men with healthy and unhealthy classification of behaviours. Multivariate logistic regression was performed to analyze predictive factors for our dichotomized categories of healthy behaviour. The most populous categories were chosen as the referent within each sub group. All analyses were unadjusted. A

two-tailed p-value of <0.05 was considered significant. Statistical analyses were performed using Stata 14.1.

DRAFT

Results

Sample demographic characteristics are reported in Table 1. The frequency of men's engagement with these unhealthy behaviours are described in Table 2.

Our findings reveal that only 5.9% (118/2000) of Canadian men engaged in all 5 healthy behaviours and 21.9% (437/2000) engaged in 4/5 so as to be classified as healthy. More than a quarter of the men had 3/5, or borderline healthy behaviours (30.8%; 616/2000), and 41.5% (829/2000) of men were classified as unhealthy with 0-2/5 healthy behaviours. Table 2 shows which unhealthy behaviours contributed to each of the CCCHB classifications. The most common unhealthy behaviours included poor sleep, lack of exercise and poor diet.

With respect to smoking, 1,609 (80.5%) of men reported healthy smoking behaviours, with 1,052 (52.6%) non-smokers, and 557 (27.9%) ex-smokers. The remaining 391 (19.6%) men were classified as unhealthy with 145 (7.3%) occasional smokers, and 246 (12.3%) regular smokers.

Audit-C questionnaire results identified 1,227 (61.4%) men reporting healthy drinking habits, while 773 (38.7%) reported behaviours that screen positively for alcohol overuse.

Less than half, 923 (46.2%) men slept a healthy 7 to <9 hours/night. Among the 1077 (53.9%) of men that reported unhealthy sleeping behaviours, 994 (92.3%) under-slept, and 83 (7.7%) overslept.

Only 1,023 (51.2%) men met the criteria of at least 150 minutes of moderate to strenuous activity/week, while 977 (48.9%) did not exercise enough. Most concerning, 20% of men were sedentary, reporting no moderate exercise.

One third, (765; 38.3%) of men were classified as consuming a healthy diet receiving a score of 12/15 or greater, while 1235 (61.8%) were classified as having unhealthy eating. Most of these men consumed a lack of vegetables/fruits and mono-polyunsaturated fats rather than excessive refined sugars, and sodium laden processed foods.

Multivariate logistic regression revealed that men with high school education were at increased risk of unhealthy behaviours, OR 1.58 (95% CI: 1.15-2.18, $p = 0.005$), as were men living with relatives, OR 2.10 (1.04-4.26, $p=0.039$), or with their partner and children, OR 1.34 (1.02-1.76, $p = 0.034$) (Table 3).

Discussion

Creation of the CCCHB included behaviours directly associated with morbidity and mortality; when available, validated questionnaires and evidence based a priori thresholds were used.

Classification of Canadian men using our CCCHB, identified that 94.1% of men have at least 1 unhealthy behaviour amenable to modification. Most concerning is that nearly half of men demonstrate 3 to 5/5 unhealthy behaviours. A cumulative effect of each additional unhealthy behaviour has been associated with increased mortality risk; in one series, hazard ratios sequentially increased from 1.37 to 6.15 as the number of unhealthy behaviours increased.^{56,57}

To understand the behaviours contributing to CCCHB, we also assessed each behaviour independently.

A sizable portion of men (53.9%) did not achieve healthy sleep (7 to <9 hours). Thus, more than half of Canadian men with poor sleep behaviours are at increased risk for mortality and significant comorbidities such as cardiovascular disease, stroke, diabetes, cancer and inflammatory states.

In our study population, only 51% of men met the minimal 150 minutes of weekly exercise as recommended by the Canadian Society of Exercise Physiology⁵⁸ and the U.S. Department of Health physical activity guidelines for Americans.⁵⁹ An 11-year longitudinal study of 12,201 older men demonstrated that those meeting this threshold had decreased risk of mortality (HR 0.74).⁵⁴ An inverse relationship between exercise duration and mortality has been described in a multitude of studies.⁶⁰ Specifically, exercise is associated with a decreased risk of conditions that heavily contribute to mortality among western nations, including: cancers,⁶¹ cardiovascular disease,⁶² hypertension,⁶² hyperlipidemia,⁶² metabolic syndrome, obesity, diabetes mellitus⁶² and suicidal ideation.⁶³ Therefore, nearly half of the current study respondents reported inadequate exercise which by extension exposed them to modifiable risks for a multitude of preventable disease, morbidity and early mortality.

A healthy diet is difficult to define from the literature,⁶⁴ in part because there have been no long-term comparative studies to place the merits of all available diets against each other.⁶⁵ However, there are a common set of dietary principles associated with health-related outcomes. We defined unhealthy dietary items as those associated with negative health outcomes in the literature. These included dietary items high in salt,²⁴ saturated fat^{25,26} and refined sugar²⁷⁻²⁹ which are independently associated with morbidity and mortality including increased risk of cardiovascular disease, and cancers.⁶⁶⁻⁶⁹ Diets rich in vegetables³¹ and mono- and poly-unsaturated fats³⁰ have been associated with reduced risk of cardiovascular disease, neurodegenerative disease, diabetes and numerous cancers.^{67,70-72} Our results identify that nearly two thirds of the respondents had unhealthy diets, placing themselves at unnecessarily high risk for morbidity and mortality. These findings are supported by previous literature where many men exhibit diets higher in fat, meat and salt with less vegetables, fruit and fiber.²²

Only a minority (19.6%) of the respondents were current smokers, in keeping with national rates,⁷³ and 27.9% of men were ex-smokers, supporting previous findings that smoking rates are declining in Canada and globally.⁷⁴ The literature characterizing the negative impact of smoking on health outcomes is strong and longstanding. Smoking is associated with cardiovascular disease, numerous cancers, pulmonary disease and significantly increased mortality. Smoking is thought to contribute 30-39% of behaviour-mediated loss of life expectancy.⁷⁵ Given the impact of smoking on health outcomes it is imperative for efforts to continue to decrease the rate of smokers and prevent premature death and morbidity.

We identified alcohol overuse in 38.7% of Canadian men. This is higher than previous studies among Canadian men, which identified 29.0% of adult men to be heavy drinkers.⁷⁶ This is concerning since alcohol overuse has been associated with self-inflicted injuries, homicide, liver disease, cancers, neuropsychiatric disease and cardiovascular disease.⁷⁷

Our study has uniquely assessed the contribution of demographic and socioeconomic factors contributing to health behaviours among Canadian men. Men with a high school graduate level of education were at greater risk of unhealthy behaviours compared to men with undergraduate degrees. Increased education has consistently demonstrated healthier behaviours and lifestyle,⁷⁸ and is integral to future health promotion. A trend also existed for reduced household income and poorer CCCHB classification, suggesting behaviours among men of lower SES likely contribute to the disparities of increased morbidity and mortality.⁷⁹

Men living with relatives or with their partner and children engaged in fewer healthy behaviours compared to men living only with their partners. Previous studies have also shown that men living with their partner have better health behaviours and overall health.^{80,81} We postulate that men living with extended families may face additional time and financial constraints, precluding opportunities to optimize health behaviours.

In our model, retired men had healthier behaviours compared to men working full time. Men working full time, likely face increased time constraints and fatigue associated with work, resulting in decreased time to focus on self-health resulting in more unhealthy behaviours.⁸² Interestingly, both part-time and full-time students demonstrated healthier behaviours which may be due to increasing trends of dietary and exercise health awareness and education.⁸³ However, these results contrast with older publications, suggesting that today's students may be making greater strides to live healthier lives.

Strengths of our study include the creation and use of a composite CCCHB classification system which includes evidence based health behaviours and respective thresholds associated with morbidity and mortality. Limitations include potential recall biases which are difficult to eliminate in self-reporting surveys such as ours. Given our aim was to capture current behaviours, this bias may be less of an issue. Our survey was conducted via an online platform, which may not be accessible to all Canadians who were not technologically savvy enough to access the survey, potentially introducing a selection bias.

Conclusion

It is fair to say that the 'problems' of men's health behaviours have attracted significant attention. The current study confirms the rhetoric in this regard, with the majority of men reporting multiple unhealthy behaviours. Specifically, ever present is the need to attend to disparities and health inequities among men, wherein greater resources might be directed toward men of lower SES to enhance their likelihood of making positive health behaviour adjustments. In addition, given that unhealthy lifestyles comprise behaviour there might be significant benefit in advocating and empowering men for small changes to an array of health practices – as distinct

from abstracting and making wholesale changes to one behaviour, as advocated by the Canadian Men's Health Foundation (dontchangemuch.ca). Although findings and recommendations from our study are representative of Canadian men, adoption of this composite classification may also provide a much-needed platform for comparisons amongst global populations and in longitudinal outcomes assessment following programmatic interventions.

DRAFT

References

1. Elterman DS, Kaplan SA, Pelman RS, Goldenberg SL. How 'male health' fits into the field of urology. *Nat Rev Urol*. 2013;10(10):606-612.
2. Information CIH. Canada's International Health System Performance Over 50 Years: Examining Potential Years of Life Lost. 2016.
3. Katon WJ, Russo JE, Heckbert SR, et al. The relationship between changes in depression symptoms and changes in health risk behaviours in patients with diabetes. *Int J Geriatr Psychiatry*. 2010;25(5):466-475.
4. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med*. 2000;160(14):2101-2107.
5. Belloc NB, Breslow L. Relationship of physical health status and health practices. *Prev Med*. 1972;1(3):409-421.
6. Haveman-Nies A, de Groot L, Burema J, et al. Dietary quality and lifestyle factors in relation to 10-year mortality in older Europeans: the SENECA study. *Am J Epidemiol*. 2002;156(10):962-968.
7. Khaw KT, Wareham N, Bingham S, Welch A, Luben R, Day N. Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS Med*. 2008;5(1):e12.
8. King DE, Mainous AG, 3rd, Geesey ME. Turning back the clock: adopting a healthy lifestyle in middle age. *Am J Med*. 2007;120(7):598-603.
9. Knuops KT, de Groot LC, Kromhout D, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA*. 2004;292(12):1433-1439.
10. Kvaavik E, Batty GD, Ursin G, Huxley R, Gale CR. Influence of individual and combined health behaviours on total and cause-specific mortality in men and women: the United Kingdom health and lifestyle survey. *Arch Intern Med*. 2010;170(8):711-718.
11. Nechuta SJ, Shu XO, Li HL, et al. Combined impact of lifestyle-related factors on total and cause-specific mortality among Chinese women: prospective cohort study. *PLoS Med*. 2010;7(9).
12. Spencer CA, Jamrozik K, Norman PE, Lawrence-Brown M. A simple lifestyle score predicts survival in healthy elderly men. *Prev Med*. 2005;40(6):712-717.
13. Tamakoshi A, Tamakoshi K, Lin Y, Yagyu K, Kikuchi S, Group JS. Healthy lifestyle and preventable death: findings from the Japan Collaborative Cohort (JACC) Study. *Prev Med*. 2009;48(5):486-492.
14. van Dam RM, Li T, Spiegelman D, Franco OH, Hu FB. Combined impact of lifestyle factors on mortality: prospective cohort study in US women. *BMJ*. 2008;337:a1440.
15. Wingard DL, Berkman LF, Brand RJ. A multivariate analysis of health-related practices: a nine-year mortality follow-up of the Alameda County Study. *Am J Epidemiol*. 1982;116(5):765-775.
16. Ford ES, Bergmann MM, Boeing H, Li C, Capewell S. Healthy lifestyle behaviours and all-cause mortality among adults in the United States. *Prev Med*. 2012;55(1):23-27.

17. Bilsker D, Goldenberg L, Davison J. A Roadmap to Men's Health: Current Status, Research Policy & Practice. *Mens Health Initiative of British Columbia* 2010.
18. Gough B, Conner MT. Barriers to healthy eating amongst men: a qualitative analysis. *Soc Sci Med.* 2006;62(2):387-395.
19. De Visser RO, Smith JA, McDonnell EJ. 'That's not masculine': masculine capital and health-related behaviour. *J Health Psychol.* 2009;14(7):1047-1058.
20. Gough B. Try to be healthy, but don't forgo your masculinity: deconstructing men's health discourse in the media. *Soc Sci Med.* 2006;63(9):2476-2488.
21. Courtenay WH. Constructions of masculinity and their influence on men's well-being: a theory of gender and health. *Soc Sci Med.* 2000;50(10):1385-1401.
22. Courtenay WH. Key determinants of the health and the well-being of men and boys. *International journal of Men's Health.* 2003;2:1-27.
23. Canada S. *Body mass index, overweight or obese, self-reported, adult, by sex, provinces and territories.* 2012.
24. Rust P, Ekmekcioglu C. Impact of Salt Intake on the Pathogenesis and Treatment of Hypertension. *Adv Exp Med Biol.* 2017(956):61-84.
25. Clifton PM, Keogh JB. A systematic review of the effect of dietary saturated and polyunsaturated fat on heart disease. *Nutr Metab Cardiovasc Dis.* 2017;27(12):1060-1080.
26. Imamura F, Micha R, Wu JH, et al. Effects of Saturated Fat, Polyunsaturated Fat, Monounsaturated Fat, and Carbohydrate on Glucose-Insulin Homeostasis: A Systematic Review and Meta-analysis of Randomised Controlled Feeding Trials. *PLoS Med.* 2016;13(7):e1002087.
27. Skop-Lewandowska A, Zajac J, Kolarzyk E. Overweight and obesity vs. simple carbohydrates consumption by elderly people suffering from diseases of the cardiovascular system. *Ann Agric Environ Med.* 2017;24(4):575-580.
28. Li N, Petrick JL, Steck SE, et al. A pooled analysis of dietary sugar/carbohydrate intake and esophageal and gastric cardia adenocarcinoma incidence and survival in the USA. *Int J Epidemiol.* 2017;46(6):1836-1846.
29. Ebrahimof S, Hosseini-Esfahani F, Mirmiran P, Fallah-Kezabi M, Azizi F. Food Patterns and Framingham Risk Score in Iranian Adults: Tehran Lipid and Glucose Study: 2005-2011. *Metab Syndr Relat Disord.* 2018.
30. Nicoll R, Howard JM, Henein MY. A review of the effect of diet on cardiovascular calcification. *Int J Mol Sci.* 2015;16(4):8861-8883.
31. Sabate J, Wien M. A perspective on vegetarian dietary patterns and risk of metabolic syndrome. *Br J Nutr.* 2015;113 Suppl 2:S136-143.
32. Van Duyn MA, Pivonka E. Overview of the health benefits of fruit and vegetable consumption for the dietetics professional: selected literature. *J Am Diet Assoc.* 2000;100(12):1511-1521.
33. Riboli E, Norat T. Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *Am J Clin Nutr.* 2003;78(3 Suppl):559S-569S.
34. Medicine Io. Crossing the Quality Chasm: A New Health System for the 21st Century. . *National Academy of Science Press.* Washington, DC2001.

35. M. E, AD. L, A R. Comparative Quantification of Health Risks: Global and Regional Burden of Disease Attributable to Selected Major Risk Factors. *World Health Organization*. Geneva 2004.
36. Lund Jensen N, Pedersen HS, Vestergaard M, Mercer SW, Glumer C, Prior A. The impact of socioeconomic status and multimorbidity on mortality: a population-based cohort study. *Clin Epidemiol*. 2017;9:279-289.
37. Turrell G, Mathers C. Socioeconomic inequalities in all-cause and specific-cause mortality in Australia: 1985-1987 and 1995-1997. *Int J Epidemiol*. 2001;30(2):231-239.
38. Singh GK, Siahpush M. Increasing inequalities in all-cause and cardiovascular mortality among US adults aged 25-64 years by area socioeconomic status, 1969-1998. *Int J Epidemiol*. 2002;31(3):600-613.
39. James PD, Wilkins R, Detsky AS, Tugwell P, Manuel DG. Avoidable mortality by neighbourhood income in Canada: 25 years after the establishment of universal health insurance. *J Epidemiol Community Health*. 2007;61(4):287-296.
40. Rogers R, Hummer R, Nam C. Demographic, Socioeconomic, and behavioural Factors Affecting Ethnic Mortality by Cause. *Social Forces*. 1996;74(4):1419-1438.
41. Canada S. 2016 Census of Population, Statistics Canada Catalogue no. 98-400-X2016001. 2016; <http://www12.statcan.gc.ca/census-recensement/2016/dp-pd/dt-td/Rp-eng.cfm?LANG=E&APATH=3&DETAIL=0&DIM=0&FL=A&FREE=0&GC=0&GID=0&GK=0&GRP=1&PID=109523&PRID=10&PTYPE=109445&S=0&SHOWALL=0&SUB=0&Temporal=2016&THEME=115&VID=0&VNAMEE=&VNAMEF=>. Accessed September 25 2017, 2017.
42. Statscan. Canadian Tobacco Use Monitoring Survey (CTUMS) 2005 Cycle 1 Questionnaire. 2005; http://www23.statcan.gc.ca/imdb-bmdi/instrument/4440_Q1_V11-eng.pdf, 2017.
43. Bush K, Kivlahan DR, McDonnell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Arch Intern Med*. 1998;158(16):1789-1795.
44. Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res*. 2007;31(7):1208-1217.
45. Chaput J, Wong S, Michaud I. Duration and quality of sleep among Canadians aged 18-79. 2017. Accessed January 7 2018, 2018.
46. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *Sleep Health*. 2015;1(4):233-243.
47. Hall MH, Smagula SF, Boudreau RM, et al. Association between sleep duration and mortality is mediated by markers of inflammation and health in older adults: the Health, Aging and Body Composition Study. *Sleep*. 2015;38(2):189-195.
48. Cai H, Shu XO, Xiang YB, et al. Sleep duration and mortality: a prospective study of 113 138 middle-aged and elderly Chinese men and women. *Sleep*. 2015;38(4):529-536.
49. Yin J, Jin X, Shan Z, et al. Relationship of Sleep Duration With All-Cause Mortality and Cardiovascular Events: A Systematic Review and Dose-Response Meta-Analysis of Prospective Cohort Studies. *J Am Heart Assoc*. 2017;6(9).

50. Godin G. The Godin-Shephard Leisure-Time Physical Activity Questionnaire. *Health & Fitness Journal of Canada*. 2011;4(1):18-22.
51. Godin G, Shephard RJ. A simple method to assess exercise behaviour in the community. *Can J Appl Sport Sci*. 1985;10(3):141-146.
52. Physiology CSFE. Canadian Physical Activity Guidelines for Adults 18-64 years 2011; http://www.csep.ca/CMFiles/Guidelines/CSEP_PAGuidelines_adults_en.pdf, 2017.
53. Physiology CSFE. Canadian Physical Activity Guidelines For Adults 65 years & Older. 2011; http://www.csep.ca/CMFiles/Guidelines/CSEP_PAGuidelines_older-adults_en.pdf, 2017.
54. Almeida OP, Khan KM, Hankey GJ, Yeap BB, Golledge J, Flicker L. 150 minutes of vigorous physical activity per week predicts survival and successful ageing: a population-based 11-year longitudinal study of 12 201 older Australian men. *Br J Sports Med*. 2014;48(3):220-225.
55. Pourmalek F, Goldenberg SL, Ho K, Skeldon SC, Patrick DM. Development and case-control validation of the Canadian Men's Health Foundation's self risk-assessment tool: "You Check". *Journal of Men's Health*. 2017;13(1):e9-e18.
56. Hamer M, Bates CJ, Mishra GD. Multiple health behaviours and mortality risk in older adults. *J Am Geriatr Soc*. 2011;59(2):370-372.
57. Krokstad S, Ding D, Grunseit AC, et al. Multiple lifestyle behaviours and mortality, findings from a large population-based Norwegian cohort study - The HUNT Study. *BMC Public Health*. 2017;17(1):58.
58. Canada Go. Physical Activity. 2011; <https://www.canada.ca/en/health-canada/services/food-nutrition/canada-food-guide/food-guide-basics/physical-activity-canada.html>. Accessed September 27 2017, 2017.
59. Services USDoHaH. Appendix 1. Physical Activity guidelines for Americans. 2015; <https://health.gov/dietaryguidelines/2015/guidelines/appendix-1/>. Accessed January 22 2018, 2018.
60. Gebel K, Ding D, Chey T, Stamatakis E, Brown WJ, Bauman AE. Effect of Moderate to Vigorous Physical Activity on All-Cause Mortality in Middle-aged and Older Australians. *JAMA Intern Med*. 2015;175(6):970-977.
61. Brown JC, Winters-Stone K, Lee A, Schmitz KH. Cancer, physical activity, and exercise. *Compr Physiol*. 2012;2(4):2775-2809.
62. Zhang X, Devlin HM, Smith B, et al. Effect of lifestyle interventions on cardiovascular risk factors among adults without impaired glucose tolerance or diabetes: A systematic review and meta-analysis. *PLoS One*. 2017;12(5):e0176436.
63. Vancampfort D, Hallgren M, Firth J, et al. Physical activity and suicidal ideation: A systematic review and meta-analysis. *J Affect Disord*. 2017;225:438-448.
64. Shim JS, Oh K, Kim HC. Dietary assessment methods in epidemiologic studies. *Epidemiol Health*. 2014;36:e2014009.
65. Katz DL, Meller S. Can we say what diet is best for health? *Annu Rev Public Health*. 2014;35:83-103.
66. Htun NC, Suga H, Imai S, Shimizu W, Takimoto H. Food intake patterns and cardiovascular risk factors in Japanese adults: analyses from the 2012 National Health and nutrition survey, Japan. *Nutr J*. 2017;16(1):61.

67. Schwingshackl L, Schwedhelm C, Hoffmann G, et al. Food groups and risk of all-cause mortality: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr*. 2017;105(6):1462-1473.
68. DiNicolantonio JJ, Lucan SC, O'Keefe JH. The Evidence for Saturated Fat and for Sugar Related to Coronary Heart Disease. *Prog Cardiovasc Dis*. 2016;58(5):464-472.
69. Baldo MP, Rodrigues SL, Mill JG. High salt intake as a multifaceted cardiovascular disease: new support from cellular and molecular evidence. *Heart Fail Rev*. 2015;20(4):461-474.
70. Capurso C, Vendemiale G. The Mediterranean Diet Reduces the Risk and Mortality of the Prostate Cancer: A Narrative Review. *Front Nutr*. 2017;4:38.
71. Cernea S, Hancu N, Raz I. Diet and coronary heart disease in diabetes. *Acta Diabetol*. 2003;40 Suppl 2:S389-400.
72. Dinu M, Pagliai G, Casini A, Sofi F. Mediterranean diet and multiple health outcomes: an umbrella review of meta-analyses of observational studies and randomised trials. *Eur J Clin Nutr*. 2017.
73. Canada S. Smokers, by sex, provinces and territories (Percent). <https://www.statcan.gc.ca/tables-tableaux/sum-som/101/cst01/health74b-eng.htm>. Accessed January 22 2018, 2018.
74. Ng M, Freeman MK, Fleming TD, et al. Smoking prevalence and cigarette consumption in 187 countries, 1980-2012. *JAMA*. 2014;311(2):183-192.
75. Manuel DG, Perez R, Sanmartin C, et al. Measuring Burden of Unhealthy Behaviours Using a Multivariable Predictive Approach: Life Expectancy Lost in Canada Attributable to Smoking, Alcohol, Physical Inactivity, and Diet. *PLoS Med*. 2016;13(8):e1002082.
76. Lukassen J, Beaudet MP. Alcohol dependence and depression among heavy drinkers in Canada. *Soc Sci Med*. 2005;61(8):1658-1667.
77. Room R, Babor T, Rehm J. Alcohol and public health. *Lancet*. 2005;365(9458):519-530.
78. Cowell AJ. The relationship between education and health behaviour: some empirical evidence. *Health Econ*. 2006;15(2):125-146.
79. Rehkopf DH, Berkman LF, Coull B, Krieger N. The non-linear risk of mortality by income level in a healthy population: US National Health and Nutrition Examination Survey mortality follow-up cohort, 1988-2001. *BMC Public Health*. 2008;8:383.
80. Hughes ME, Waite LJ. Health in household context: living arrangements and health in late middle age. *J Health Soc Behav*. 2002;43(1):1-21.
81. Rossow I, Rise J. Living arrangements and health behaviours in adolescence and young adulthood. *Health Education Research*. 1993;8(4):495-503.
82. M. I. The health consequences of retirement. *Journal of Human Resources*. 2014;49(1):195-233.
83. Kyrkou C, Tsakoumaki F, Fotiou M, et al. Changing Trends in Nutritional behaviour among University Students in Greece, between 2006 and 2016. *Nutrients*. 2018;10(1).

Figures and Tables

Fig. 1. Among 5362 Canadians that visited the survey page, only 2000 met inclusion and exclusion criteria. Of note, 144 patients were excluded for visiting “DontChangeMuch.ca,” which is a website created by the Canadian Men’s Health Foundation aimed to address many of these health behaviours; these men were excluded to prevent bias in selecting men actively pursuing healthy behaviour and lifestyle changes.




Table 1. Baseline respondent characteristics and demographics	
Demographics and baseline characteristics	Number of participants (%)
Sex	
Male	2000 (100)
Age	
19–29	379 (19.0)
30–54	934 (46.7)
55+	687 (34.4)
Province	
British Columbia	265 (13.3)
Alberta	217 (10.9)
Saskatchewan	131 (6.6)
Ontario	768 (38.4)
Quebec	476 (23.8)
Martime provinces	142 (7.1)
Territories	1 (0.1)
Minority	
Yes	218 (10.9)
No	1782 (89.1)
Household income	
<\$20 000	129 (6.5)
\$20 000–39999	282 (14.1)
\$40 000–59 999	336 (16.8)
\$60 000–79 999	355 (17.8)
\$80 000–99 999	278 (13.9)
\$100 000–119 999	222 (11.1)
\$120 000–139 999	134 (6.7)
\$140 000 +	264 (13.2)
Highest level of education	
Primary school or less	6 (0.3)
Some high school	48 (2.4)
High school graduate	301 (15.1)
Some college/trade school	225 (11.3)
Graduated college/trade school	422 (21.1)
Some university	175 (8.8)
University undergraduate degree	495 (24.8)
University graduate degree	328 (16.4)
Sexual preference	
Heterosexual	1805 (90.3)
Homosexual	118 (5.9)
Bisexual	48 (2.4)
Not sure or questioning	25 (1.3)
Other	4 (0.2)

Number of children age <19 living with participant	
None	1623 (81.2)
1	177 (8.9)
2	140 (7.0)
3	42 (2.1)
4+	18 (0.9)
Living arrangement	
Partner	818 (40.9)
Alone	449 (22.5)
Partner & children	392 (19.6)
Parent	178 (8.9)
Non-relatives	54 (2.7)
Children	49 (2.5)
Relatives	41 (2.1)
University or college campus	14 (0.7)
Other	5 (0.3)
Employment	
Employed full-time	1003 (50.2)
Employed part-time	155 (7.8)
Self-employed	170 (8.5)
Looking for employment	93 (4.7)
Unable to work	63 (3.2)
Retired	453 (22.7)
Studying full-time	108 (5.4)
Studying part-time	47 (2.4)
Home caregiver	14 (0.7)
Total number of participants	2000 (100)

Total health behaviour classification			Men classified as “unhealthy” (%) per category				
	Health classification	Total men (%)	Smoking	Alcohol	Sleep	Exercise	Diet
Very Healthy	0 unhealthy behaviours	118/2000 (5.9)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Healthy	1 unhealthy behaviour	437/2000 (21.9)	12/437 (2.7)	84/437 (19.2)	137/437 (31.4)	74/437 (16.9)	130/437 (29.7)
Borderline	2 unhealthy behaviours	616/2000 (30.8)	55/616 (8.9)	205/616 (33.3)	297/616 (48.2)	293/616 (47.6)	382/616 (62.0)
Unhealthy	3 unhealthy behaviours	571/2000 (28.5)	140/571 (24.5)	281/571 (49.2)	418/571 (73.2)	394/571 (69.0)	480/571 (84.0)
	4 unhealthy behaviours	219/2000 (11.0)	145/219 (66.2)	164/219 (74.9)	186/219 (84.9)	177/219 (80.8)	204/219 (93.2)
	5 unhealthy behaviours	39/2000 (2.0)	39/39 (100)	39/39 (100)	39/39 (100)	39/39 (100)	39/39 (100)

For each group of men with a number of unhealthy behaviours (0–5), a breakdown is provided demonstrating the number (%) of men with each type of unhealthy behaviour.

Table 3. Multivariate analysis of demographic, socioeconomic, and comorbid conditions impacting total health behaviour classification		
Predictors of unhealthy total health behaviour	Odds ratio (95% CI)	p
Demographics		
Minority	1.01 (0.74–1.38)	0.949
Young age	1.00 (0.99–1.01)	0.699
Employment status (employed full-time reference)		
Employed part-time	0.76 (0.49–1.16)	0.202
Looking for employment	0.62 (0.38–1.02)	0.061
Unable to work	1.38 (0.75–2.51)	0.290
Retired	0.63 (0.45–0.88)	0.007*
Student full-time	0.59 (0.36–0.96)	0.034*
Student part-time	0.51 (0.26–1.00)	0.049*
Home caregiver	3.39 (0.67–17.24)	0.142
Orientation (heterosexual reference)		
Gay or lesbian	1.13 (0.82–1.56)	0.464
Living location (Ontario reference)		
British Columbia	0.74 (0.54–1.01)	0.054*
Alberta	1.00 (0.73–1.37)	0.986
Saskatchewan & Manitoba	1.08 (0.73–1.60)	0.698
Quebec	1.01 (0.79–1.29)	0.954
Atlantic provinces & Territories	0.92 (0.63–1.35)	0.665
Living arrangement (living with partner reference)		
Alone	1.03 (0.79–1.36)	0.809
Parent	1.07 (0.71–1.60)	0.752
Partner & children	1.34 (1.02–1.76)	0.034*
Non-relatives	1.35 (0.74–2.48)	0.328
Children	1.36 (0.74–2.50)	0.325
Relatives	2.10 (1.04–4.26)	0.039*
University or college campus	0.60 (0.17–2.12)	0.432
Other	1.31 (0.19–8.88)	0.782
Highest level of education (university undergraduate reference)		
Primary school	0.77 (0.13–4.59)	0.777

Some high school	1.48 (0.13–4.59)	0.228
High school graduate	1.58 (1.15–2.18)	0.005*
Some college/trade school	1.23 (0.87–1.73)	0.235
Graduated college/trade school	1.09 (0.82–1.44)	0.563
Some university	0.98 (0.67–1.43)	0.908
University graduate degree	0.86 (0.64–1.17)	0.341
Household income (\$60 000–79 999 reference)		
\$19 999 or less	1.13 (0.71–1.82)	0.595
\$20 000–39 999	0.95 (0.68–1.34)	0.785
\$40 000–59 999	1.33 (0.97–1.84)	0.076
\$80 000–99 999	1.12 (0.81–1.56)	0.494
\$100 000–119 999	0.71 (0.49–1.03)	0.068
\$120 000–139 999	0.71 (0.72–1.69)	0.650
\$140 000 or more	0.83 (0.58–1.19)	0.313

*Statistically significant predictor variable $p < 0.05$. CCCHB total health behaviour classification is dichotomized into: very healthy, healthy, borderline vs. unhealthy. CI: confidence interval.