

Canadian consensus algorithm for erectile rehabilitation following prostate cancer treatment

Dean S. Elterman¹; Anika R. Petrella¹; Lauren M. Walker²; Brandon Van Asseldonk¹; Leah Jamnicky¹; Gerald B. Brock³; Stacy Elliott²; Antonio Finelli¹; Jerzy B. Gajewski⁴; Keith A. Jarvi⁵; John Robinson⁶; Janet Ellis⁷; Shaun Shepherd¹; Hossein Saadat¹; Andrew Matthew¹

¹Division of Urology, University Health Network, Toronto, ON, Canada; ²Department of Psychiatry, University of British Columbia, Vancouver, BC, Canada; ³Department of Surgery, St. Joseph's Hospital, London, ON, Canada;

⁴Department of Urology, Dalhousie University, Halifax, NS, Canada; ⁵Murray Koffler Urologic Wellness Centre, Mount Sinai Hospital, Toronto, ON, Canada; ⁶Clinical Psychology Department, University of Calgary, Calgary, AB, Canada; ⁷Department of Psychology, Sunnybrook Health Sciences Centre, Toronto, ON, Canada

Funding: The consensus meeting was funded by a Prostate Cancer Canada Movember grant.

Cite as: *Can Urol Assoc J* 2018 December 3; Epub ahead of print.

<http://dx.doi.org/10.5489/cuaj.5653>

Published online December 3, 2018

Abstract

Introduction: The present descriptive analysis carried out by a pan-Canadian panel of expert healthcare practitioners (HCPs) summarizes best practices for erectile rehabilitation following prostate cancer (PCa) treatment. This algorithm was designed to support an online sexual health and rehabilitation e-clinic (SHARe-Clinic), which provides biomedical guidance and supportive care to Canadian men recovering from PCa treatment. The implications of the algorithm may be used inform clinical practice in community settings.

Methods: Men's sexual health experts convened for the TrueNTH Sexual Health and Rehabilitation Initiative Consensus Meeting to address concerns regarding erectile dysfunction (ED) therapy and management following treatment for PCa. The meeting brought together experts from across Canada for a discussion of current practices, latest evidence-based literature review, and patient interviews.

Results: An algorithm for ED treatment following PCa treatment is presented that accounts for treatment received (surgery or radiation), degree of nerve-sparing, and level of pro-erectile treatment invasiveness based on patient and partner values. This algorithm provides an approach from both a biomedical and psychosocial focus that is tailored to the patient/partner presentation.

Algorithm for erectile rehabilitation after PCa treatment

Regular sexual activity is recommended, and the importance of partner involvement in the treatment decision-making process is highlighted, including the management of partner sexual concerns.

Conclusions: The algorithm proposed by expert consensus considers important factors like the type of PCa treatment, the timeline of erectile recovery, and patient values, with the goal of becoming a nationwide standard for erectile rehabilitation following PCa treatment.

Introduction

Prostate cancer (PCa) is the second most frequent cancer found in men, accounting for 21% of the estimated new annual cases.^{1,2} A number of treatment options are available to patients with PCa, including active surveillance (AS), radiotherapy (RT), and radical prostatectomy (RP). Treatment options are chosen based on a number of factors, such as clinical stage, patient's age, and the presence of co-morbid diseases in the patient.^{3,4}

PCa treatments, regardless of modality, increase the likelihood of erectile dysfunction (ED).⁵ Rates for ED lasting 2 years or more following RP range from 66-75%.^{5,6} Similar results are reported in men following RT at 3 years (37-81%).⁷ ED rates have been found to be 10 to 15-fold higher in men with PCa than their similarly aged peers.⁸ ED is a significant threat to the quality of life of men diagnosed with PCa, as 60% of affected men experience severe distress from ED.⁹⁻¹¹ The loss of sexual activity and resultant challenge to masculinity have been shown to negatively affect quality of life.¹²

Currently, there remains a gap in the systematic and comprehensive care of sexual dysfunction after PCa treatment. This presents a significant barrier to continuity of care for PCa survivors across Canada. The TrueNTH SHARe Clinic was developed to provide sexual health support to men located across Canada. The present manuscript incorporates research evidence, patient perspectives and clinical expertise from experts in the field. To address the lack of consistency in care of sexual dysfunction after PCa treatment a meeting was held. The TrueNTH Sexual Health and Rehabilitation Initiative Consensus Meeting was held with the purpose of developing an ED therapy algorithm following PCa treatment. The algorithm would be disseminated and used in the TrueNTH SHARe Clinic. The present manuscript describes the development of an ED therapy algorithm that uniquely accounts for patient and partner values and goals for erectile recovery, type of PCa treatment, time since treatment, nerve-sparing status, and also includes thematic recommendations for psychosocial support

Methods

A pan-Canadian panel of men's sexual health experts convened for the TrueNTH Sexual Health and Rehabilitation Initiative Meeting to develop a consensus on treating ED in patients treated for localized PCa. The consensus panel meeting was held on October 31, 2016 in Toronto. A limited peer esteem snowballing technique (PEST) was used to identify expert opinion panellists

Algorithm for erectile rehabilitation after PCa treatment

based on research or clinical expertise in sexual dysfunction post-PCa treatment.¹³ A 17-member expert opinion panel provided commentary on variation in ED treatment approaches through various programs and services across Canada. The group represented a wide range of backgrounds including nursing, urology, urologic oncology, radiation oncology, psychology, psychiatry, as well as patient advocates.

The meeting involved several key decision-making components. Before the consensus meeting, all panelists reviewed the evidence-based medical literature on ED, particularly concerning physiology, pathophysiology, diagnosis, and treatment of ED following PCa treatment. The meeting began with a discussion of various cancer-related sexual health rehabilitation programs that exist across the country. Special attention was paid to patients' concerns about the current practice of ED therapy and management following PCa treatment, specifically the advantages and disadvantages of existing practices. Subsequently, the latest clinical guidelines on sexual rehabilitation after PCa treatment were reviewed and summarized.¹⁴⁻¹⁶

Patients' perspectives and feedback regarding gaps in current practice and desired practice were incorporated into the development of the TrueNTH SHARe Clinic algorithm. Possibilities for content and structure were outlined and discussed. Proposed strategies for the uptake of the algorithm to the medical community were also outlined.

Results

The TrueNTH Sexual Health and Rehabilitation Initiative Consensus Meeting established that a tailored, comprehensive ED therapy algorithm for patients (and their partners) after localized PCa treatment should recognize the following:

- 1) real-life results are often more modest than reported in the literature;
- 2) patients are overly optimistic about the likelihood that they will be in the minority of patients who do not experience ED and about the ease with which they will adapt to use of pro-erectile therapies;^{17,18}
- 3) poorly managed patient expectations can be demotivating in sustaining use of pro-erectile therapies;
- 4) while the best time to introduce erectile rehabilitation remains unclear, pre-PCa treatment psychoeducation on sexual dysfunction and available ED therapies is a necessity;¹⁹⁻²²
- 5) post-PCa treatment ED recovery typically occurs over a minimum of two (or more) years;²³
- 6) desired pace of return of ED function and willingness to engage in invasive treatment varies across patients and partners;^{24,25}
- 7) exclusive focus on achieving erections via any means, may overlook the values or goals of patients and their partners;
- 8) the uptake and adherence to pro-erectile therapies is generally poor;²⁶⁻²⁸
- 9) detailed education on the systematic use of pro-erectile therapies is often lacking in post-PCa sexual health care;¹⁹

Algorithm for erectile rehabilitation after PCa treatment

- 10) inclusion of the partner in the recovery process is optimal;^{24,29-31}
- 11) the process of ED therapy re-challenging over the recovery period is necessary to achieve optimal erectile functioning and to manage perceived treatment failure;³²
- 12) maintaining regular sexual activity (penetrative or non-penetrative) during the course of erectile recovery is advantageous for individuals and couple's well-being.^{33,34}

Algorithm

The algorithm for managing ED is illustrated in Table 1. The algorithm focuses on therapeutic strategies to erectile recovery, with psychosocial considerations at each stage of treatment. The focus of the algorithm is split based on patients' values or goals for erectile recovery. Further considerations are offered based on the cancer treatment (radiation or surgery), nerve-sparing status, preference for less invasive vs. more aggressive treatment approach, and inclination for a pharmaceutical vs. mechanical approach to pro-erectile therapy.

Section 1: Algorithm process

Patients will choose a management pathway (Table 1) based on the type of PCa treatment received (surgery or radiation), followed by the patients desired level of invasiveness (low or high), and the nerve sparing status (bilateral nerve sparing; BNS, unilateral nerve sparing; UNS, and non-nerve sparing; NNS), and lastly if they prefer a pharmaceutical vs. mechanical approach to pro-erectile therapy. Thus, vacuum erection devices (VED) are available in each quadrant for a non-biomedical approach. The inflatable penile prosthesis is available for patients who are refractory to both biomedical and non-biomedical approaches, however as this strategy is often reserved later in the treatment trajectory (beyond 2 years) to allow sufficient time for natural recovery and sufficient retrial with first-line treatments, it is not presented in the algorithm.³⁵ Patient's preferences are also taken into account in terms of tolerance of degree of treatment invasiveness, and the patients' goals for erectile recovery (articulated further in section 2 below). In addition to the specific recommendations for pro-erectile therapy, patients are recommended to maintain regular sexual activity (at least weekly) whether penetrative or non-penetrative, and/or masturbation. Clinicians may also consider combination therapies if necessary, depending on the patient's desire to challenge ED over time.^{36,37}

Section 2: Patient goals for erectile recovery and psychosocial considerations

As a supplemental document to the algorithm, Table 2 outlines patients' values and goals for erectile rehabilitation and should be used as a resource to guide treatment planning. The panel agreed that there were essentially two pathways that the patient might choose. The primary goals of "long-term penile health" and (i.e. optimising changes for natural recovery of erections with a relatively non-invasive approach) vs. "short-term erectile function" were identified in the algorithm. Clinicians are encouraged to assess the patient's goals for outcomes as this may change with time. Furthermore, partners may also wish to provide input into these goals.

Section 3: Benchmarking and normalization

In addition to the algorithm based on patient treatment and preference for ED rehabilitation (pharmaceutical vs. mechanical), the panel noted the importance of providing patients with a point of reference for their progress in terms of expected timeline of erectile function recovery. This attempt at typical response benchmarking is intended to help patients manage their expectations and normalize their recovery process. The panel determined that sufficient empirical evidence for typical response benchmarking was only available for the post-radical patient population (see Table 3). More research is needed to determine the timeline for patients treated with radiation therapy.³⁸

Discussion

Developing a treatment plan for ED after PCa involves balancing a number of factors including different trajectories for radiation vs. surgical treatment, nerve-sparing status, and the degree of invasiveness of various pro-erectile therapies, along with determining patient and partner's goals for erectile recovery.²⁰⁻²³ Other patient related factors must all be considered in ensuring that pro-erectile therapies are optimally successful including: patient expectations of pro-erectile therapy and timeline of recovery, the role of the partner in erectile recovery, and the patients' sexual beliefs, masculine values, possible grief in response to sexual losses, and potential performance anxiety.³⁹⁻⁴¹

Sub-algorithms were proposed for each group of patients (e.g. those treated with radiation vs. surgery), as the panel recognized advantages to more accurately capture between-group differences such as different trajectory of impact of PCa treatment on erectile function. Consensus on structures of the sub-algorithms was achieved; the structure would include consideration for timeline benchmarks, long-term penile health vs. short-term erectile function, as well as additional sexual concerns, all while integrating patient preferences. Suggested timeline benchmarks for the sub-algorithms were 1-3, 6, 12, 18, and 24 months from baseline, for surgical outcomes only. The suggested benchmarks were based on historical practice adopted in clinical trials and the expected success rates of relevant treatment approaches. However, a significant limitation is the limited empirical evidence due to heterogeneity in treatment timelines and a low risk of external validity in clinical studies. Further discussion is required to establish outcome benchmarks for patients with a delayed response to treatment and also for patients receiving radiation therapy.

The sub-algorithm structure for penile rehabilitation (i.e. long-term care of penile health) was designed to focus on treatment flow with emphasis placed on the patient's desire to engage in strategies that would promote penile health and improve capacity for natural return of erections with time. This sub-algorithm also considers the patients' tolerance of treatment invasiveness. Consistent with treatment guidelines, phosphodiesterase type 5 inhibitors (PDE5i) were identified as first-line treatment for ED post RP with either daily or as-needed dosing.³⁵ Daily PDE5i dosing has demonstrated to be an effective alternative to as-needed dosing in patients in patients who had an incomplete response to therapy with maximum-dose as-required

Algorithm for erectile rehabilitation after PCa treatment

PDE5i.⁴² If patient response to PDE5i is poor at 3 months, the introduction of intracavernous injection (ICI) treatment can be introduced as an option, with the possible retreat of PDE5i. Alternatively, treatment with VED can be used in place of ICI or PDE5i at any point in the recovery timeline thus patient treatment preference should be prioritized to guide long-term clinical management.

In response to direct feedback from patients and clinicians on the panel, patient preference for expedient recovery of erections was identified as an alternative goal to long-term penile health. In this scenario, the short-term erectile function component of the algorithm should be followed (right column). The first increment of the treatment timeline should be 1-3 months. Patients motivated to achieve an erection immediately, should initiate ICI therapy as first-line treatment. Otherwise, if treating with PDE5i, a higher dose is recommended with the possibility of combined treatment with ICI and/or the VED³⁶

The following considerations were identified for inclusion in the content of the sub-algorithms: climacturia, dysorgasmia, alterations to penile anatomy, and reduced sexual desire. Other psychosexual factors that interact with patient preferences were considered for inclusion such as the importance of sexual activity and intimacy, expectations for recovery with pro-erectile therapy, performance anxiety, and the recognition of the impact of loss and grief. Important highlights included interventions to communicate the importance of persistence to therapy, are required to ensure patients can realize the full benefits of clinical therapy. Psychosocial interventions are required to redefine patient sex life, integrating the sexual partner(s) when possible, and focusing on building on therapeutic gains.

Patients should be counselled about the likelihood of natural recovery of erections and timelines for recovery, including the significant role of the patients' baseline erectile function prior to PCa treatment. In fact, baseline erectile function may have even more influence on erectile recovery than use of pro-erectile therapies.⁴³ Age is also a major predictor, with younger men showing better natural erectile recovery.^{23,44} The percent probability of erectile recovery by 24 months, for men who have full erections at baseline is 63% ≤ 60 years vs. 37% in men ≥ 65 years. In contrast, in men with recently diminished erectile function at baseline, the percent probability is lower at 48% ≤ 60 years vs. 26% in men ≥ 65 years, and even worse for men who had partial erectile function at baseline, 35% ≤ 60 years vs. 18% in men ≥ 65 years. Rates in all categories increase slightly out to 36 months.²³

Additional insights from the literature should be considered when counselling patients about erectile recovery, in order to promote adherence to, and acceptance of the use of pro-erectile therapies. For example, in the general population, sustained use of PDE5i is better when prescribed with the knowledge and involvement of the patients' partner.⁴⁵ Given the overly optimistic mindset of patients about the probability of experiencing ED and the ease of treatments for ED, patients need adequate preparation before starting pro-erectile therapy. The algorithm was designed with the explicit purpose of being embedded in sexual health programming for PCa patients.¹⁸

Algorithm for erectile rehabilitation after PCa treatment

The most common reason for discontinuation of PDE5i is lack of treatment efficacy (e.g. hardness of erection),²⁷ therefore, in a context when the need to re-challenge is the norm, patients need to be forewarned that PDE5i may not work for all patients immediately, and that likelihood of effectiveness increases the further out the patient is from surgery.^{16,46} In the context of radiation therapy, the opposite effect is observed. Patients often do not understand that tactile stimulation is necessary to prompt an erectile response even with the use of PDE5i and that spontaneous erections are unlikely.⁴⁷ Furthermore, patients should be informed that none of the pro-erectile therapies promote sexual desire or interest.

In addition, a variety of psychosocial factors are listed throughout the Table 2. Ideally good erectile rehabilitation should be provided in a bio-psychosocial context. Where this is not possible, clinicians may wish to read the evidence-based literature for suggestions on enhancing the likelihood of successful sexual recovery.

The long-term goal is to utilize the treatment algorithms in the development of the web-based TrueNTH SHARE-Clinic. The opinion of the panel was that the application of a personalized clinical treatment tool with the TrueNTH SHARE-Clinic will improve treatment for ED in PCa patients by tailoring individualized therapies in a clinical environment that promotes patients' participation in decision making. The web-based clinic features tailored content including: personalized sexual health coaching, a multimodal virtual library, and symptom monitoring with feedback mechanisms. Thus, the TrueNTH SHARE Clinic uniquely combines web-based sexual health counselling, with an ED therapy algorithm designed to provide patients and their partners with accessible, personalized, post-PCa long-term sexual health care.

Conclusion

Because of the significantly high rate of ED after PCa treatment, it is critical to establish practices guidelines for the management of ED in this patient population. There exist a number of clinical guidelines to inform first-line through fourth or fifth line therapies, however, to date, none of these guidelines attempt to directly incorporate patient values into the treatment decision making process. Instead they seem to focus on a linear progression of pro-erectile aids from least invasive to most invasive. However, ideal clinical management must be tailored to patient (and partner) preferences including specific goals for erectile recovery. In addition, considerations from the diverse representation on the multi-disciplinary panel highlighted the inclusion of patient experiences directly identified by patient advocates, as well as psychosocial recommendations to enhance patient and partner efficacy, acceptance, and compliance with pro-erectile aids. Furthermore, sensitivity to the discrepancies in trajectory of ED development after radiation treatment vs. surgery need to be accommodated when making recommendations in order to maximize applicability. Further work is required to define subsequent clinical management benchmarks, particularly in the context of radiation treatment. The development of an algorithm that incorporates these many different facets, should prove a novel and useful tool for clinicians wishing to support patient's erectile recovery after PCa treatment.

References

1. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: Sources, methods and major patterns in globocan 2012. *Int J Cancer* 2015;136:E359-86. 2014/09/16.
2. Canadian Cancer Society's Steering Committee for Canadian Cancer Statistics. *Canadian cancer statistics 2017*. Toronto, ON: Canadian Cancer Society;2017.
3. Zumsteg ZS, Spratt DE, Pei I, et al. A new risk classification system for therapeutic decision making with intermediate-risk prostate cancer patients undergoing dose-escalated external-beam radiation therapy. *Eur Urol* 2013;64:895-902. 2013/04/02.
4. Epstein JI, Zelefsky MJ, Sjoberg DD, et al. A contemporary prostate cancer grading system: A validated alternative to the gleason score. *Eur Urol* 2016;69:428-35. 2015/07/15.
5. Robinson JW, Moritz S, Fung T. Meta-analysis of rates of erectile function after treatment of localized prostate carcinoma. *Int J Radiat Oncol Biol Phys* 2002;54:1063-8. 2002/11/07.
6. Haglind E, Carlsson S, Stranne J, et al. Urinary incontinence and erectile dysfunction after robotic versus open radical prostatectomy: A prospective, controlled, nonrandomised trial. *Eur Urol* 2015;68:216-25. 2015/03/17.
7. Akbal C, Tinay I, Simsek F, et al. Erectile dysfunction following radiotherapy and brachytherapy for prostate cancer: Pathophysiology, prevention and treatment. *Int Urol Nephrol* 2008;40:355-63. 2007/10/26.
8. Bacon CG, Mittleman MA, Kawachi I, et al. Sexual function in men older than 50 years of age: Results from the health professionals follow-up study. *Ann Intern Med* 2003;139:161-8. 2003/08/06.
9. Nelson CJ, Mulhall JP, Roth AJ. The association between erectile dysfunction and depressive symptoms in men treated for prostate cancer. *J Sex Med* 2011;8:560-6. 2010/12/16.
10. Cooperberg MR, Koppie TM, Lubeck DP, et al. How potent is potent? Evaluation of sexual function and bother in men who report potency after treatment for prostate cancer: Data from capsure. *Urology* 2003;61:190-6. 2003/02/01.
11. Penson DF, Latini DM, Lubeck DP, et al. Is quality of life different for men with erectile dysfunction and prostate cancer compared to men with erectile dysfunction due to other causes? Results from the exceed data base. *J Urol* 2003;169:1458-61. 2003/03/12.
12. Bourke L, Boorjian SA, Briganti A, et al. Survivorship and improving quality of life in men with prostate cancer. *Eur Urol* 2015;68:374-83. 2015/05/06.
13. Christopoulos D. Peer esteem snowballing: A methodology for expert surveys. Paper presented at: Eurostat conference for new techniques and technologies for statistics2009.
14. Chung E, Brock GB. Emerging and novel therapeutic approaches in the treatment of male erectile dysfunction. *Curr Urol Rep* 2011;12:432-43. 2011/09/17.
15. Salonia A, Burnett AL, Graefen M, et al. Prevention and management of postprostatectomy sexual dysfunctions part 2: Recovery and preservation of erectile function, sexual desire, and orgasmic function. *Eur Urol* 2012;62:273-86. 2012/05/12.
16. Kirby MG, White ID, Butcher J, et al. Development of uk recommendations on treatment for post-surgical erectile dysfunction. *Int J Clin Pract* 2014;68:590-608. 2013/11/06.

Algorithm for erectile rehabilitation after PCa treatment

17. Wittmann D, He C, Coelho M, et al. Patient preoperative expectations of urinary, bowel, hormonal and sexual functioning do not match actual outcomes 1 year after radical prostatectomy. *J Urol* 2011;186:494-9. 2011/06/18.
18. Bronner G, Shefi S, Raviv G. Sexual dysfunction after radical prostatectomy: Treatment failure or treatment delay? *J Sex Marital Ther* 2010;36:421-9. 2010/10/07.
19. Boberg EW, Gustafson DH, Hawkins RP, et al. Assessing the unmet information, support and care delivery needs of men with prostate cancer. *Patient Educ Couns* 2003;49:233-42. 2003/03/19.
20. Vickers AJ, Kent M, Mulhall J, et al. Counseling the post-radical prostatectomy patients about functional recovery: High predictiveness of current status. *Urology* 2014;84:158-63. 2014/05/16.
21. Wittmann D, Montie JE, Hamstra DA, et al. Counseling patients about sexual health when considering post-prostatectomy radiation treatment. *Int J Impot Res* 2009;21:275-84. 2009/07/18.
22. Forbat L, White I, Marshall-Lucette S, et al. Discussing the sexual consequences of treatment in radiotherapy and urology consultations with couples affected by prostate cancer. *BJU Int* 2012;109:98-103. 2011/06/03.
23. Rabbani F, Stapleton AM, Kattan MW, et al. Factors predicting recovery of erections after radical prostatectomy. *J Urol* 2000;164:1929-34. 2000/11/04.
24. Wittmann D, Carolan M, Given B, et al. Exploring the role of the partner in couples' sexual recovery after surgery for prostate cancer. *Support Care Cancer* 2014;22:2509-15. 2014/04/15.
25. Davison BJ, Keyes M, Elliott S, et al. Preferences for sexual information resources in patients treated for early-stage prostate cancer with either radical prostatectomy or brachytherapy. *BJU Int* 2004;93:965-9. 2004/05/15.
26. Lee DJ, Cheetham P, Badani KK. Penile rehabilitation protocol after robot-assisted radical prostatectomy: Assessment of compliance with phosphodiesterase type 5 inhibitor therapy and effect on early potency. *BJU Int* 2010;105:382-8. 2009/09/15.
27. Buvat J, Buttner H, Hatzimouratidis K, et al. Adherence to initial pde-5 inhibitor treatment: Randomized open-label study comparing tadalafil once a day, tadalafil on demand, and sildenafil on demand in patients with erectile dysfunction. *J Sex Med* 2013;10:1592-602. 2013/04/05.
28. Matthew AG, Goldman A, Trachtenberg J, et al. Sexual dysfunction after radical prostatectomy: Prevalence, treatments, restricted use of treatments and distress. *J Urol* 2005;174:2105-10. 2005/11/11.
29. Davison BJ, Matthew A, Elliott S, et al. Assessing couples' preferences for postoperative sexual rehabilitation before radical prostatectomy. *BJU international* 2012;110:1529-35.
30. Fisher WA, Eardley I, McCabe M, et al. Erectile dysfunction (ed) is a shared sexual concern of couples ii: Association of female partner characteristics with male partner ed treatment seeking and phosphodiesterase type 5 inhibitor utilization. *J Sex Med* 2009;6:3111-24. 2009/08/14.
31. Dunn ME. Restoration of couple's intimacy and relationship vital to reestablishing erectile function. *J Am Osteopath Assoc* 2004;104:S6-10. 2004/04/16.
32. Jiann BP, Yu CC, Su CC, et al. Rechallenge prior sildenafil nonresponders. *Int J Impot Res* 2004;16:64-8. 2004/02/14.

Algorithm for erectile rehabilitation after PCa treatment

33. Wittmann D, Carolan M, Given B, et al. What couples say about their recovery of sexual intimacy after prostatectomy: Toward the development of a conceptual model of couples' sexual recovery after surgery for prostate cancer. *The journal of sexual medicine* 2015;12:494-504.
34. Walker LM, Wassersug RJ, Robinson JW. Psychosocial perspectives on sexual recovery after prostate cancer treatment. *Nature Reviews Urology* 2015;12:167.
35. Bella AJ, Lee JC, Carrier S, et al. 2015 CUA practice guidelines for erectile dysfunction. *Canadian Urological Association Journal* 2015;9:23.
36. Kimura M, Caso JR, Banez LL, et al. Predicting participation in and successful outcome of a penile rehabilitation programme using a phosphodiesterase type 5 inhibitor with a vacuum erection device after radical prostatectomy. *BJU Int* 2012;110:E931-8. 2012/04/24.
37. Defade BP, Carson CC, 3rd, Kennelly MJ. Postprostatectomy erectile dysfunction: The role of penile rehabilitation. *Rev Urol* 2011;13:6-13. 2011/08/10.
38. Wiegner EA, King CR. Sexual function after stereotactic body radiotherapy for prostate cancer: Results of a prospective clinical trial. *Int J Radiat Oncol Biol Phys* 2010;78:442-8. 2010/02/09.
39. Zaider T, Manne S, Nelson C, et al. Loss of masculine identity, marital affection, and sexual bother in men with localized prostate cancer. *J Sex Med* 2012;9:2724-32. 2012/09/20.
40. Gannon K, Guerra-Blanco M, Patel A, et al. Re-constructing masculinity following radical prostatectomy for prostate cancer. *Aging Male* 2010;13:258-64. 2010/11/12.
41. Wittmann D, Foley S, Balon R. A biopsychosocial approach to sexual recovery after prostate cancer surgery: The role of grief and mourning. *J Sex Marital Ther* 2011;37:130-44. 2011/03/15.
42. Kim E, Seftel A, Goldfischer E, et al. Comparative efficacy of tadalafil once daily in men with erectile dysfunction who demonstrated previous partial responses to as-needed sildenafil, tadalafil, or vardenafil. *Curr Med Res Opin* 2015;31:379-89. 2014/12/03.
43. Castiglione F, Nini A, Briganti A. Penile rehabilitation with phosphodiesterase type 5 inhibitors after nerve-sparing radical prostatectomy: Are we targeting the right patients? *Eur Urol* 2014;65:673-4. 2014/01/07.
44. Gallina A, Ferrari M, Suardi N, et al. Erectile function outcome after bilateral nerve sparing radical prostatectomy: Which patients may be left untreated? *J Sex Med* 2012;9:903-8. 2012/01/14.
45. Carvalheira AA, Pereira NM, Maroco J, et al. Dropout in the treatment of erectile dysfunction with pde5: A study on predictors and a qualitative analysis of reasons for discontinuation. *J Sex Med* 2012;9:2361-9. 2012/05/24.
46. Montorsi F, Brock G, Stolzenburg JU, et al. Effects of tadalafil treatment on erectile function recovery following bilateral nerve-sparing radical prostatectomy: A randomised placebo-controlled study (reactt). *Eur Urol* 2014;65:587-96. 2013/10/31.
47. Sadovsky R, Brock GB, Gutkin SW, et al. Toward a new 'epoch': Optimising treatment outcomes with phosphodiesterase type 5 inhibitors for erectile dysfunction. *Int J Clin Pract* 2009;63:1214-30. 2009/07/25.

Figures and Tables

Table 1. Pro-erectile therapy algorithm. BNS: bilateral nerve sparing; ICI: intracavernous injection; NNS: non-nerve sparing; PC: prostate cancer; PDE5i: phosphodiesterase type 5 inhibitors; UNS: unilateral nerve sparing; VED: vacuum erection devices.

Pro-erectile Therapy Algorithm

Baseline recommendation for all patients: Regular sexual activity (at least once a week)
When applicable – Patient's partners should be included in the treatment plan

Patient Preference:		LOW Invasiveness			HIGH Invasiveness		
Patient Goals:		Long-term penile health			Short-term erectile function		
PC Treatment Status:		BNS	UNS	NNS	BNS	UNS	NNS
Surgery	Pharmaceutical Approach	Daily & PRN full dose PDE5i	PRN use of full dose PDE5i	MUSE	ICI	ICI	ICI
	Mechanical Approach	VED	VED	VED	VED	VED	VED
Radiation	Pharmaceutical Approach	Daily & PRN full dose PDE5i			ICI		
	Mechanical Approach	VED			VED		

Algorithm for erectile rehabilitation after PCa treatment

Table 2. Timelines and patient goals for erectile recovery. ICI: intracavernous injection; PDE5i: phosphodiesterase type 5 inhibitors; VED: vacuum erection devices.

• **Consider time since treatment and expectation management**

Timeline	Patient Goal for Erectile Recovery	
	Long-term penile health	Short-term erectile function
Prior to treatment	Treatment flow post treatment focused on the patient's desire to maintain penile health over the long term. Psychosocial focus: Normalize use of pro-erectile aids, education on timeline of recovery, manage expectation for erectile recovery and success with use of pro-erectile therapies, exploring patient goals for erectile recovery. *Note: patient goals should be continuously evaluated	Treatment flow post treatment focused on the patient's desire to achieve functional erections in the short term.
6-weeks post treatment	<ul style="list-style-type: none"> PDE5i first-line; dosage, daily vs. PRN Include penile stimulation in treatment flow 	<ul style="list-style-type: none"> Include penile stimulation in treatment flow If immediate erection is the goal, consider ICI therapy first-line; or possibly PDE5i full dose if responsive
	Psychosocial focus: Importance of regular sexual activity (penetrative or non-penetrative), education on timeline of recovery, manage expectation for erectile recovery and success with use of pro-erectile therapies, encourage couples to maintaining sexual intimacy (penetrative or non-penetrative)	
10-weeks post treatment	<ul style="list-style-type: none"> PDE5i first-line; dosage, daily vs. PRN Include penile stimulation in treatment flow 	<ul style="list-style-type: none"> Include penile stimulation in treatment flow If immediate erection is the goal, consider ICI therapy first-line; or possibly PDE5i full dose if responsive
	Psychosocial focus: Assessment and treatment of Partner Concerns, Encouraging adherence to pro-erectile treatment plan, encourage couples to maintaining sexual intimacy (penetrative or non-penetrative)	
4 months post treatment	<ul style="list-style-type: none"> For patients who do not respond initially to PDE5i's, introduce ICI or MUSE or VED 	<ul style="list-style-type: none"> Depending on response to PDE5i, consider continued use of effective PDE5i or ICI or MUSE or VED
	Psychosocial focus: Recognition of grief response to sexual losses or performance anxiety, acknowledge impact on masculinity, encourage couples to maintaining sexual intimacy (penetrative or non-penetrative)	
6 months post treatment	<ul style="list-style-type: none"> For patients who do not respond initially to PDE5i's, introduce ICI or MUSE or VED 	<ul style="list-style-type: none"> Depending on response to PDE5i, consider continued use of effective PDE5i or ICI or MUSE or VED
	Psychosocial focus: Re-visit expectations and goals & explore level of patient bother due to ED; Sexual desire and fantasy, encourage couples to maintaining sexual intimacy (penetrative or non-penetrative)	
12 months post treatment	<ul style="list-style-type: none"> Rechallenge with PDE5i as needed For patients who do not respond to PDE5i's, introduce ICI or MUSE or VED 	<ul style="list-style-type: none"> Depending on response to PDE5i, consider continued use of effective PDE5i or ICI or MUSE or VED
	Psychosocial focus: Redefining sex life & building on success	
18 months post treatment	<ul style="list-style-type: none"> Rechallenge with PDE5i as needed For patients who do not respond to PDE5i's, introduce ICI or MUSE or VED 	<ul style="list-style-type: none"> Depending on response to PDE5i, consider continued use of effective PDE5i or ICI or MUSE or VED
	Psychosocial focus: confirming goals and expectations	
24 months post treatment	<ul style="list-style-type: none"> Rechallenge with PDE5i as needed For patients who do not respond to PDE5i's, introduce ICI or MUSE or VED 	<ul style="list-style-type: none"> Depending on response to PDE5i, consider continued use of effective PDE5i or ICI or MUSE or VED
	Psychosocial focus: long term goal setting and management, Adaptation and Acceptance, Satisfaction with pro-erectile therapy	
	*ICI = Intracavernosal injections, PDE5i = phosphodiesterase-5 inhibitors, MUSE = intraurethral alprostadil, VED = vacuum erection device	

Table 3. Timelines and benchmarks post-surgery. ED: erectile dysfunction.**Timelines & Benchmarks Post-Surgery**

- Use of pro-erectile aids or devises encouraged at all timepoints
- Consider time since treatment and expectation management

Timeline	Benchmarks using the Erectile Firmness Scale (1-10; 1 flaccid, 6 just hard enough to achieve penetration, 10 full erection)
Prior to treatment	Evaluate baseline erectile function (poorer function at baseline may require more aggressive approach) * some men experience ED post diagnosis/pre-treatment due to stress and anxiety
6-weeks post treatment	1-3: Lack of natural sexual function to be expected.
10-weeks post treatment	1-3: Lack of natural sexual function to be expected.
4 months post treatment	1-4: Some early recovery of mild to moderate tumescence in <10% of patients.
6 months post treatment	2-6: Some early recovery of mild to moderate tumescence in <10% of patients.
12 months post treatment	3-7: 20-40% recover natural erectile function hard enough for penetration.
18 months post treatment	4-7: 20-40% recover natural erectile function hard enough for penetration.
24 months post treatment	5-8: 30-50% recover natural erectile function hard enough for penetration.