

Case — Recurrent incarcerated inguinal bladder hernia with small bowel obstruction

Jeff McLellan, BComm¹; Zuhaib M. Mir, MD²; Michael Di Lena, MD³; Sulaiman Nanji, MD, PhD FRCSC²; Darren Beiko, MD, MBA, FRCSC³

¹Northern Ontario School of Medicine, Sudbury, ON, Canada; ²Department of Surgery, Division of General Surgery, Queen's University, Kingston, ON, Canada; ³Department of Urology, Queen's University, Kingston, ON, Canada

Cite as: *Can Urol Assoc J* 2019;13(11):E382-4. <http://dx.doi.org/10.5489/cuaj.5740>

Published online February 26, 2019

Introduction

Inguinal hernias are a common surgical complaint, with a lifetime incidence requiring repair of 27% in men and 3% in women.¹ Recurrence of an inguinal hernia requiring repair depends on a number of factors, including repair type, surgeon experience, and patient factors, but has been estimated at 1–4%.^{2,3} The bladder is involved in 1–4% of inguinal hernias in the general population, and involvement is as high as 10% in men over age 50.⁴ Risk factors for urinary bladder involvement in an inguinal hernia include male gender, age over 50, and obesity.⁵ The mainstay of treatment for an inguinal bladder hernia is herniorrhaphy.

Most small bladder hernias are discovered intraoperatively due to the lack of urinary symptoms prior to herniorrhaphy. In fact, less than 7% are diagnosed preoperatively.⁴ Small urinary bladder hernias are often asymptomatic, but clinical manifestations can include urinary frequency, urgency, nocturia, hematuria, and double-phase micturition.⁶ Extensive herniation of the bladder, commonly referred to as a scrotal cystocele, is much less frequent. These are more likely to present with urinary symptoms, and urgent intervention is usually required, as bladder infarction and/or obstruction may result.⁶ Inguinal bladder hernias with associated small bowel obstruction are rare and rates have not been characterized in the literature.

We present a case of a recurrent inguinal hernia with bladder and small bowel incarceration, as well as associated small bowel obstruction. In addition, aside from a case involving complete herniation of the bladder,⁶ we present the first such images of a Foley catheter balloon being found within the herniated portion of the bladder.

Case report

Our patient was an 80-year old male, who presented to the emergency department with a two-day history of vomiting, diarrhea, expressive aphasia, and confusion. Comorbidities prior to presentation included Alzheimer's disease, prostate adenocarcinoma (T2a, Gleason 6/10, treated with curative radiotherapy), hypertension, gout, previous atrial tachycardia, previously resected vestibular schwannoma, and a previous left inguinal hernia repaired with Shouldice technique. He was seen urgently by the neurology service, however, acute cerebrovascular pathology was subsequently ruled out. The focus then shifted to the patient's gastrointestinal symptoms, and initial bloodwork revealed a mild leukocytosis ($11.2 \times 10^9/L$), elevated creatinine ($165 \mu\text{mol/L}$), but an otherwise unremarkable hematological and biochemical profile. Blood cultures and urine culture were drawn and ultimately negative, ruling out underlying sepsis. Initial computed tomography (CT) of the abdomen and pelvis demonstrated a recurrent left-sided inguinal hernia containing a loop of dilated small bowel with resultant proximal small bowel dilatation. The hernia also contained a portion of non-opacified bladder with the Foley catheter balloon/tip within the herniated segment (Fig. 1), causing the bladder to remain distended.

The Foley catheter balloon was deflated and repositioned. In addition, manual reduction of the hernia was attempted in the emergency department under procedural sedation. Repeat CT images showed successful reduction of the small bowel from the hernia, but persistent incarceration of the urinary bladder within the left inguinal hernia (Fig. 2). There was no contrast extravasation seen from the bladder, suggesting that its wall was intact.

Nonetheless, given the persistent incarceration of the bladder, as well as the need to assess viability and ischemia via cystoscopy, the patient was consented and booked for urgent operative repair of the hernia. An open plug and patch repair was performed. Intraoperatively, the hernia was noted to be indirect, with no evidence of residual small bowel




Fig. 1. Initial computed tomography scan demonstrating a left inguinal hernia containing a portion of the urinary bladder with the Foley catheter balloon (white arrow) within the herniated segment. Proximal small bowel dilatation is also seen.

herniation. Externally, the bladder tissue did not appear to be necrotic or perforated. An intraoperative cystoscopy was also performed, which demonstrated mucosal edema at the dome of the bladder without any necrosis or perforation. A Foley catheter was left in situ and the patient tolerated the procedure well.

Followup cystogram on postoperative day 8 showed no perforation or filling defects and the Foley catheter was removed. Repeat cystourethroscopy at eight weeks postoperatively demonstrated mild meatal stenosis and bulbar urethral stricture requiring dilation, with minimal inflammation at the site of bladder incarceration but an otherwise well-healed bladder. On followup, the patient remained asymptomatic from a bowel and bladder perspective and was, therefore, discharged from care.

Discussion

Inguinal hernias are a common surgical complaint, with bladder involvement in approximately 1–4% of cases. While there are many reported cases of inguinal bladder hernias, we believe this to be the first case where the Foley catheter balloon was inflated in the herniated portion of the bladder, thereby helping identify the underlying pathology. There is one prior case where a suprapubic catheter balloon was seen in the hernia, however, that was a case involving complete bladder herniation.⁶

Small inguinal bladder hernias tend to be asymptomatic from a urinary perspective and, therefore, most often discovered intraoperatively, which can lead to a more challenging repair and the possibility of bladder injury during herniorrhaphy. Imaging modalities such as CT and cystograms




Fig. 2. Repeat computed tomography scan showing persistent incarceration of the urinary bladder and successful reduction of small bowel after bedside reduction attempted under procedural sedation.

should be considered preoperatively to clearly delineate the anatomy if there is suspicion for bladder involvement, as well as to guide surgical repair.

When recognized preoperatively, the mainstay of treatment is surgical repair through an open inguinal approach, reducing the bladder to its original location in the pelvis. Partial cystectomy is rarely indicated, except in the case of bladder wall necrosis, herniated bladder diverticulum, tight hernia neck, or bladder tumor within the herniated segment.^{7,8} As such, we recommend routine intraoperative cystoscopy as an easy strategy to avoid missing such pathologies.

In this case, there was also an associated small bowel obstruction. While the small bowel was able to be reduced without operative intervention, incarceration of the bladder segment was an indication for urgent surgery. A review of the literature returns 18 cases of incarcerated inguinal bladder hernias, only two of which had associated small bowel obstructions.

Conclusions

To the best of our knowledge, this is the first reported case of an incarcerated inguinal bladder hernia where the Foley catheter balloon was inflated in the herniated portion of the bladder. The incarcerated small bowel was reduced preoperatively, however, persistence of bladder incarceration was the indication for operative repair. Given that the herniated bladder tissue was edematous but not necrotic, partial cystectomy was not required and the patient recovered well after herniorrhaphy.

Competing interests: Dr. Beiko has received a grant from Cook Medical. The remaining authors report no competing personal or financial interests related to this work.

This paper has been peer-reviewed.

References

1. Primates P, Goldacre MJ. Inguinal hernia repair: Incidence of elective and emergency surgery, readmission, and mortality. *Int J Epidemiol* 1996;25:835-39. <https://doi.org/10.1093/ije/25.4.835>
2. Magnusson J, Gustafsson U, Nygren J, et al. Rates of and methods used at reoperation for recurrence after primary inguinal hernia repair with Prolene Hernia System and Lichtenstein. *Hernia* 2018;22:439-44. <https://doi.org/10.1007/s10029-017-1705-9>
3. Ramjist J, Dossa F, Stukel T, et al. Reoperation for inguinal hernia recurrence in Ontario: A population-based study. *Hernia* 2018;23:647-54. <https://doi.org/10.1007/s10029-018-1822-0>
4. Moufid K, Touiti D, Mohamed L. Inguinal bladder hernia: Four case analyses. *Rev Urol* 2013;15:32-6. <https://doi.org/10.4103/2156-7514.97758>
5. Oruç MT, Akbulut Z, Özozan Ö, et al. Urological findings in inguinal hernias: A case report and review of the literature. *Hernia* 2004;8:76-9. <https://doi.org/10.1007/s10029-003-0157-6>
6. Bisharat M, O'Donnell ME, Thompson T, et al. Complications of inguinoscrotal bladder hernias: A case series. *Hernia* 2009;13:81-4. <https://doi.org/10.1007/s10029-008-0389-6>
7. Wagner AA, Arcand P, Bamberger MH. Acute renal failure resulting from huge inguinal bladder hernia. *Urology* 2004;64:156-7. <https://doi.org/10.1016/j.urol.2004.03.040>
8. Vindlachervu RR, Zayyan K, Burgess NA, et al. Extensive bladder infarction in a strangulated inguinal hernia. *Br J Urol* 1996;77:926-7. <https://doi.org/10.1046/j.1464-410X.1996.07035.x>

Correspondence: Dr. Darren Beiko, Department of Urology, Queen's University, ON, Canada; Darren.Beiko@kingstonhsc.ca