

Pediatric bladder augmentation – Panacea or Pandora’s box?

James P.J. Ross; Melise Keays; Christopher Neville; Michael Leonard; Luis Guerra
Children’s Hospital of Eastern Ontario (CHEO), Department of Surgery, Division of Urology, University of
Ottawa, Ontario, Canada

Cite as: *Can Urol Assoc J* 2020 January 20; Epub ahead of print.
<http://dx.doi.org/10.5489/cuaj.6024>

Published online January 20, 2020

Abstract

Introduction: Bladder augmentation is a surgery that can increase bladder capacity and compliance. The objective of this study was to provide a longitudinal review of pediatric bladder augmentation at a tertiary Canadian center.

Methods: A retrospective review was performed on patients who underwent bladder augmentation at a tertiary pediatric hospital between 1986 and 2014. The primary objective was short- and long-term complications of augmentation. Secondary objectives were to review number of augmentation procedures performed over time and the utility of routine postoperative cystograms.

Results: A total of 56 procedures were performed on 54 patients (28 males, 26 females) of mean age 10 years (standard deviation [SD] 5) and mean followup eight years (SD 5). The most common bowel segment used was ileum (87.5%). Twenty-eight patients (50%) received catheterizable channels. Overall complication rate was 15%, and the most common complications were urinary tract infections (68.5%), worsening hydronephrosis (14.8%), bladder stone formation (14%), and hematuria (13%). In total, 19 of 54 (35.2%) patients returned to the operating room. The incidence of bladder perforation was 3.6%. Complications with the catheterizable channel occurred in 13 of 28 (46.4%), of which 10 were related to stomal stenosis. Forty patients had postoperative cystograms and extravasation was seen in three (7.5%). There was no malignancy during the followup. Only four augmentations were performed from 2008–2014.

Conclusions: Bladder augmentation likely represents a safe surgical treatment option. Extravasation on postoperative cystogram was uncommon and thus it may not be indicated routinely. The number of augmentation procedures performed has declined in recent years.

Introduction

Patients with neurogenic bladders and high intra-vesicle pressures are at risk for a number of complications including recurrent infection, urinary incontinence, and renal deterioration.¹ A number of conservative treatments have been used to help optimize bladder function in these patients, including clean intermittent catheterization, oral and intravesical anticholinergics, as well as intra-detrusor Botox injections.² Bladder augmentation, or augmentation cystoplasty, is a surgical procedure that is used when more conservative treatment methods have failed.³ The surgery involves anastomosis of a harvested bowel segment to the native bladder to increase capacity and compliance.

In spite of the potential benefits of bladder augmentation, there have been a number of studies which have reported high incidence of both short and long-term morbidity associated with this procedure. Common post-operative complications include bladder stones, bladder rupture, renal deterioration, metabolic abnormalities, and malignancy.³⁻¹⁰ Due to a decreased incidence of associated congenital disorders in the western world as well as improvements in early detection and treatment of patients with neurogenic bladders, the rates of augmentation cystoplasty performed appear to be declining in both the United Kingdom and the United States.^{3,4}

To our knowledge there have been no Canadian series reporting on long-term outcomes of pediatric bladder augmentation. The objective of this study is to review our experience with bladder augmentation in the Canadian pediatric population and to evaluate the incidence of post-operative complications at our institution. As part of this analysis, we will also look to evaluate the rate of augmentation procedures performed over time. A secondary objective is to evaluate the utility of routine post-operative cystograms in augmented patients. We hypothesize that for augmentation cystoplasty in our tertiary Canadian centre: 1) the rate of postoperative complication is low, 2) the rate of augmentation cystoplasty has decreased over time, and 3) routine post-operative cystogram is not necessary.

Methods

Following IRB approval, a retrospective chart review was conducted for all patients who were followed post-bladder augmentation at a tertiary pediatric hospital between 1986 and 2014. Patients were followed from the time of their surgery up to 18 years of age, at which point they were transitioned to the care of an adult Urologist. Patient records were reviewed for demographic information including date of birth, gender, age at time of procedure, and indication for augmentation. Operative details were collected including date of procedure, tissue segment used in augmentation, as well as any secondary procedures performed. Length of follow-up as well as the number of patients who were lost to follow-up prior to

the age of 18 years were also collected. Details of post-operative investigations, including ultrasound and cystoscopy were reviewed. The number of patients who underwent post-operative cystogram as well as the rate of extravasation was also collected.

Specific attention was directed towards the short and long-term complications of augmentation. Post-operative complications related to the augmentation procedure included urinary tract infection (UTI), renal deterioration, worsening hydronephrosis, bladder perforation, gastric reservoir perforation, bladder stone formation, metabolic abnormalities, wound infection, and hematuria. UTI was defined as any positive urine culture in addition to at least one of the following associated symptoms: dysuria, abdominal pain, hematuria, and/or fever. Renal deterioration was defined as the development of chronic kidney disease following augmentation in the context of documented normal pre-operative renal function. Metabolic abnormalities included patients who developed persistent electrolyte disturbances necessitating ongoing medical management. Wound/channel infections were identified based on the presence of localized inflammatory symptoms, including erythema, edema, and/or purulent discharge. Hematuria included any instance of gross hematuria, which was not secondary to traumatic catheterization. The incidence as well as the indication for reoperation was collected for both augmentation and catheterizable channel related complications. All eligible patients underwent routine screening for malignancy with annual cystoscopy starting at 10 years following their augmentation.

Results

A total of 56 augmentation procedures were performed on 54 patients, of which there were 28 males and 26 females. Five different Urologists performed augmentation procedures during this time period. The mean age at the time of augmentation was 9.6 (SD 5) years. The mean follow-up was 7.6 (SD 5) years. Catheterizable channels were created in 28 (50%) patients. In total three patients were lost to follow-up prior to the age of 18 (5.6%). Twenty-two (40.7%) patients underwent an additional procedure at the time of augmentation. This included unilateral ureteric reimplantation in 10, bilateral ureteric reimplantation in eight, bladder neck closure in four, bladder neck reconstruction in two, open appendectomy in two, cystolithotomy in one, and rectus fascial sling in one. Gastric reservoirs were performed in two patients in 1995 and 1996 respectively. A summary of patient demographics as well as a breakdown of the indication for augmentation and type of augmentation performed can be seen in Table 1.

Thirty-seven (68.5%) patients experienced at least one symptomatic UTI. Among the patients with normal baseline renal functioning, worsening of renal function occurred in one patient with ileocystoplasty and two patients with gastric reservoirs. Metabolic abnormalities were noted in one patient with a gastric reservoir and two patients with ileocystoplasty. The patient with the gastric reservoir developed hypochloremic hypokalemic metabolic alkalosis, whereas one patient with the ileocystoplasty developed

hyperchloremic metabolic acidosis and the other developed vitamin B12 deficiency. Complications with the catheterizable channel were noted in 13 out of 28 (46.4%) patients, which included stomal stenosis in 10 patients, bleeding in two, and infection in one (Table 2).

In total 19 out of 54 (35.2%) patients returned to the OR due to post-operative complications. The mean time to reoperation was 31.4 months (SD 28.8 months). Of the patients who required a second operation, 10 (52.6%) were due to complications with the catheterizable channel. Repeat bladder augmentation was required in one ureterocystoplasty and one ileocystoplasty. Bladder perforation occurred in one ileo-colocystoplasty and one ileocystoplasty. Bladder calculi extraction was performed using an open approach in two patients and an endoscopic approach in two patients (Table 3).

Table 4 shows the timing of post-operative complications in patients who underwent augmentation cystoplasty. Bladder perforation occurred within 8 days post-operatively in all three patients. Other complications, including stoma stenosis, electrolyte abnormalities, bladder stone formation, and worsening hydronephrosis were seen in more long-term follow-up.

There was an increase in the rate of augmentation procedures performed between 2002 and 2008. The maximum number of augmentations performed in a single year was six in 2003. Between 2008 and 2014 only four augmentations were performed (Figure 1).

Forty patients received a post-operative cystogram (74%). The mean time to post-operative cystogram was 21.4 days. Extravasation on post-operative cystogram was reported in three (7.5%) patients, however, these were minor and all resolved with longer suprapubic drainage. There were no malignancies identified on surveillance cystoscopy for those 18 patients with a follow-up period greater than or equal to 10 years. Mean duration of annual cystoscopy surveillance was 4.7 years for these patients. There were no deaths attributable to augmentation.

Discussion

The overall post-operative complication rate of augmentation cystoplasty was 15%. This incidence of complication post-augmentation is not unexpected as other studies have also documented a high rate of morbidity in these patients. The majority of complications encountered, however, were minor in severity and required only minimal intervention. The incidence of more serious complications of bladder augmentation, including bladder perforation, malignancy, and renal deterioration were low in comparison. Of note, the incidence of complications secondary to the catheterizable channel was relatively high in comparison, with an overall rate of 46.4%.

While symptomatic UTIs were common in the post-operative period (68.5%), they were not considered a direct surgical complication. It is well known that there is a high incidence of bacteriuria in these patients, particularly in those patients employing clean

intermittent catheterization.^{4,11–15} Furthermore, there is a lack of consensus in terms of how to properly define and treat UTIs amongst this population.¹⁵ Given the high incidence of UTIs and the difficulty of diagnosis within this population, physicians should be vigilant for symptoms of infection.

Metabolic abnormalities were noted in 5.6% of patients. Of these, two patients received ileocystoplasty, of which one developed hyperchloremic metabolic acidosis while the other developed vitamin B12 deficiency. This is consistent with existing literature, which suggests that up to 50% of patients with ileocystoplasty will develop metabolic acidosis while the incidence of vitamin B12 deficiency was between 3% and 20%.^{10,16,17} Furthermore, one patient with a gastric reservoir developed hypochloremic hypokalemic metabolic alkalosis. This is a known risk factor for the use of stomach tissue in augmentation due to gastric secretion of hydrochloric acid and has been found to occur in up to 7% of patients post-gastrocystoplasty.^{18,19} The recognized risk of electrolyte disturbance necessitates the need for routine laboratory monitoring of these patients. At our institution, patients underwent annual screening for electrolyte and other metabolic disturbances post-operatively.

Bladder stone formation was a common post-operative complication (14.8%). Our rate of bladder stones is consistent with other studies which have described an incidence of between 11% and 52%.^{7,10,20–23} The high rate of bladder stones amongst augmented patients suggests that regular screening may be indicated. At our institution, patients underwent annual screening with ultrasound to assess for development of possible complications including stones. Bladder stones result from calcification of accumulated bowel mucous as well as chronic bacterial colonization of the urinary tract. Therefore, the use of preventative strategies, including frequent bladder irrigation, should be promoted.²⁴ Furthermore, there was a high rate of reoperation amongst patients who developed bladder stones (50%). Therefore, the potential for repeat intervention secondary to stone formation should be discussed with the patient and their family prior to augmentation.

The incidence of decrease in renal function post-augmentation was 5.6%. This is in keeping with existing literature which suggest an incidence of between 3% and 20.3%.^{20,25} Unlike previous studies, however, patients with a diagnosis of chronic kidney disease due to a pre-existing condition, such as lower urinary tract obstruction, were not attributed as having worsening renal function as a complication of augmentation. It is of note that both patients with gastric reservoir had normal pre-operative renal function, and experienced a subsequent renal deterioration. The exact cause is unclear but is thought to be secondary to chronic reflux of acidic urine from the stomach.

Worsening hydronephrosis was noted in 15% of patients. Despite this relatively high incidence, a large majority were managed expectantly. Only two patients required repeat surgical intervention for worsening hydronephrosis, with one patient requiring

nephrostomy tube drainage and eventual repeat augmentation, and another requiring bilateral ureteric reimplantation.

There were no cases of malignancy identified. This is reflective of literature to date, which have noted a low overall incidence of malignancy post-augmentation, ranging from 0 to 5.5%.^{10,21,26–29} Patients underwent routine annual cystoscopy surveillance starting at 10 years post-operatively. Our screening protocol is based on the recommendations from previous studies based on the long latency period between operation and development of malignancy.²⁹ The mean follow-up period was 8 years, however, and the majority of patients did not reach the point where regular screening for malignancy was indicated. Furthermore, for the 18 patients in whom regular surveillance was initiated, the mean surveillance period was short at 4.7 years. Therefore, it is possible that additional cases of malignancy could develop as the follow-up period for these patients is extended.

The incidence of bladder perforation was 3.6%. This is lower than other studies, which have reported rates of between 5% and 12.8%.^{20,30,31} The low rate of perforation could be attributed to the regular access to medical care and follow-up in our pediatric population. Furthermore, at our institution patients received regular education and training from a multi-disciplinary care team, which promoted a strict protocol of frequent bladder irrigation and catheterization to prevent bladder over-distention. In spite of the relatively low incidence, bladder perforation represents a potential source of significant morbidity and mortality, which necessitates vigilant follow-up and patient awareness.

The rate of return to OR post-augmentation was high (35.2%) but not unexpected as other studies have also reported a high incidence of surgical complications.^{4,11–13,23,24} Most patients requiring repeat operation, however, were due to complications with the catheterizable channel (52.6%). Of these patients, most required only simple dilatation or revision of a stenotic stoma. A study by Husmann et al. (2017) found a similar high incidence of complications associated with continent catheterizable channels, with a reported complication rate of 40% for Mitrofanoff and 71% for Monti channels.¹⁰ The high rate of complication and reoperation for those patients with a catheterizable channel necessitates regular follow-up and screening. We suggest that inquiries regarding the functioning of the channel along with thorough inspection of the stomal opening should be incorporated into every follow-up visit.

The timing of post-operative complications was variable depending on the type of complication. Bladder perforation was seen almost immediately in all three cases, with timing ranging from 3 to 8 days post-operatively. Other complications including stoma stenosis, bladder calculi, electrolyte abnormalities, and worsening hydronephrosis were seen well over one year post-operatively on average. The wide range of timing of complications further necessitates the need for long term follow-up in these patients.

Within our institution, there was an increase in the number of procedures performed between 2000 and 2008, with a total of 36 augmentations performed during this time. During this period there was a new staff surgeon hired with special expertise in this field which may have led to the relative higher number of augmentation surgeries. From 2009 to 2014 there was a relative decrease in the number of operations with only 4 augmentations performed despite no change in staff surgeons during this period. Such a decrease in rate of augmentation has also been documented in other centres and may be attributed to improvements in early management of neurogenic bladder through UTI prevention and medical optimization, as well as a declining incidence of spina bifida and other congenital conditions in the Western world.^{3,4}

There is lack of existing literature on the utility of routine post-operative cystogram in patients who have undergone augmentation cystoplasty. In our series, the incidence of extravasation on post-operative cystogram was low at 7.5%. Furthermore, all patients with extravasation were managed expectantly with a longer period of catheterization and no patient required active surgical intervention. We would therefore suggest that routine post-operative cystogram might not be indicated in regular post-operative screening. Instead, it should be reserved for select cases where a leak or perforation is suspected based on clinical suspicion.

As these patients reach the age of majority, transition to appropriate adult care is of utmost importance. At our institution, all patients 18 years of age were transitioned to an adult Urologist with a specific expertise and interest in management of these complex patients. Given the complexity of these patients and need for routine ongoing follow-up, a thorough patient history as well as specific follow-up instructions, including the need for annual cystoscopic surveillance starting at 10 years post-operatively was provided to the accepting Urologist. For the three patients that were lost to follow-up prior to the age of 18 years, two had moved to another city and one's parents wished to be followed closer to home. Appropriate referrals to Urologists in the area of their new residence had been made prior to departure from our care.

There are some limitations to our study. First of all, the retrospective nature of our study represents a potential source of selection bias. In addition, we have a relatively small sample size. There was also a relatively short mean follow-up period of 8 years; therefore, we may have under-reported certain long-term complications. Finally, our study examined the outcomes of bladder augmentation at only a single tertiary center which may limit the applicability of our results to a broader pediatric population.

Conclusions

Overall, bladder augmentation represents a safe surgical alternative to increasing bladder capacity and compliance when other more conservative methods have failed. Despite this, there is an evident decrease in the number of augmentation procedures performed in recent years, which may be due to better medical management options coupled with a decreased incidence of associated congenital disorders in the western world. Catheterizable channels represent a significant source of morbidity and reoperation thereby necessitating prudent follow-up and screening for associated complications. Although no malignancies were identified, this may be due to our relatively short follow-up period. The concern for developing malignancy warrants careful surveillance. Post-operative cystograms may not be necessary, as the rate of extravasation is low. The lower rate of bladder perforation may be due to improved access to care along with a strict bladder catheterization and irrigation protocol in our institution. Further study would be required to confirm this.

DRAFT


References

1. Roth JD and Cain MP. Neuropathic Bladder and Augmentation Cystoplasty. *Urol Clin North Am* 2018;45: 571–85.
2. Leu R and Stearns GL. Complications of Botox and their Management. *Curr Urol Rep* 2018;19:90.
3. Biers SM, Venn SN and Greenwell TJ. The past, present and future of augmentation cystoplasty. *BJU Int* 2012;109:1280–93.
4. Schlomer BJ and Copp HL. Cumulative incidence of outcomes and urologic procedures after augmentation cystoplasty. *J Pediatr Urol* 2014;10:1043–50.
5. Shekarriz B, Upadhyay J, Demirbilek S, et al. Surgical complications of bladder augmentation: comparison between various enterocystoplasties in 133 patients. *Urology* 2000;55:123–28.
6. Metcalfe PD, Casale AJ, Kaefer MA, et al. Spontaneous bladder perforations: a report of 500 augmentations in children and analysis of risk. *J Urol* 2006;175:1466–70.
7. Balachandra B, Swanson PE, Upton MP, et al. Adenocarcinoma arising in a gastrocystoplasty. *J Clin Pathol* 2007;60:85–87.
8. Khoury JM, Timmons SL, Corbel L, et al. Complications of enterocystoplasty. *Urology* 1992;40:9–14.
9. Palmer LS, Franco I, Kogan SJ, et al. Urolithiasis in children following augmentation cystoplasty. *J Urol* 1993;150:726–29.
10. Husmann DA. Lessons learned from the management of adults who have undergone augmentation for spina bifida and bladder exstrophy: Incidence and management of the non-lethal complications of bladder augmentation. *Int J Urol* 25:94–101.
11. Greenwell TJ., Venn SN, and Mundy AR. Augmentation cystoplasty. *BJU Int* 2001;88: 511–25.
12. Veeratterapillay R, Thorpe AC and Harding C. Augmentation cystoplasty: Contemporary indications, techniques and complications. *Indian J Urol* 2013;29: 322–27.
13. Hendren WH and Hendren RB. Bladder augmentation: experience with 129 children and young adults. *J Urol* 1990;144:445–53.
14. Khoury JM, Timmons SL, Corbel L, et al. Complications of enterocystoplasty. *Urology* 1992;40:9–14.
15. Madden-Fuentes RJ, McNamara ER, Lloyd JC, et al. Variation in Definitions of Urinary Tract Infections in Spina Bifida Patients: A Systematic Review. *Pediatrics* 2013;132: 132–39.
16. Poulsen AL and Steven K. Acid-base metabolism after bladder substitution with the ileal urethral Kock reservoir. *Br J Urol* 1996;78:47–53.
17. Matsui U, Topoll B, Miller K, et al. Metabolic long-term follow-up of the ileal neobladder. *Eur Urol* 1993;24:197–200.

18. Gilbert SM and Hensle TW. Metabolic consequences and long-term complications of enterocystoplasty in children: a review. *J Urol* 2005;173:1080-86.
19. Çetinel B, Kocjancic E and Demirdağ Ç. Augmentation cystoplasty in neurogenic bladder. *Investig Clin Urol* 2016;57:316–23.
20. Shekarriz B, Upadhyay J, Demirbilek S, et al. Surgical complications of bladder augmentation: comparison between various enterocystoplasties in 133 patients. *Urology* 2000;55:123–28.
21. Metcalfe PD, Cain MP, Kaefer M, et al. What is the need for additional bladder surgery after bladder augmentation in childhood? *J Urol* 2006;176:1801–05.
22. Kaefer M, Hendren WH, Bauer SB, et al. Reservoir calculi: a comparison of reservoirs constructed from stomach and other enteric segments. *J Urol* 1998;160: 2187–90.
23. DeFoor W, Minevich E, Reddy P, et al. Bladder calculi after augmentation cystoplasty: risk factors and prevention strategies. *J Urol* 2004;172:1964–66.
24. Hensle TW, Bingham J, Lam J, et al. Preventing reservoir calculi after augmentation cystoplasty and continent urinary diversion: the influence of an irrigation protocol. *BJU Int* 2004;93:585–87.
25. Obermayr F, Szavay P, Schaefer J, et al. Outcome of augmentation cystoplasty and bladder substitution in a pediatric age group. *Eur J Pediatr Surg* 2011;21:116–19.
26. North AC and Lakshmanan Y. Malignancy associated with the use of intestinal segments in the urinary tract. *Urol Oncol* 2007;25:165–67.
27. Soergel TM, Cain MP, Misseri R, et al. Transitional cell carcinoma of the bladder following augmentation cystoplasty for the neuropathic bladder. *J Urol* 2004;172: 1649–51.
28. Lane T and Shah J. Carcinoma following augmentation ileocystoplasty. *Urol Int* 2000; 64:31–32.
29. Biardeau X, Chartier-Kastler E, Rouprêt M, et al. Risk of malignancy after augmentation cystoplasty: A systematic review. *Neurourol Urodyn* 2016;35:675–82.
30. DeFoor W, Tackett L, Minevich E, et al. Risk factors for spontaneous bladder perforation after augmentation cystoplasty. *Urology* 2003;62:737–41.
31. Bauer SB, Hendren WH, Kozakewich H, et al. Perforation of the augmented bladder. *J Urol* 1992;148:699–703.
32. Mehmood S, Alhazmi H, Al-Shayie M, et al. Long-term outcomes of augmentation cystoplasty in a pediatric population with refractory bladder dysfunction: a 12 year follow-up experience at single center. *Int Neurourol J* 2018;22:287–94.
33. Telli O, Ozcan C, Hamidi N, et al. Preoperative risk factors predicting complication rates of augmentation cystoplasty using the modified Clavien classification system in pediatric population. *Urology* 2016; 97: 166-71.

Figures and Tables

Fig. 1. Number of augmentation cystoplasties performed per year.



DR

Table 1. Patient gender, indication for augmentation, and type of augmentation procedure performed	
	Number (%)
Number of patients	54
Males	28 (51.9)
Females	26 (48.1)
Indication for augmentation:	
Spina bifida	32 (59.3)
Bladder/cloacal exstrophy	9 (16.7)
Posterior urethral valves	4 (7.4)
Non-neurogenic neurogenic bladder	3 (5.6)
Sacral agenesis	3 (5.6)
Traumatic spinal cord injury	2 (3.7)
Bilateral ectopic ureters	1 (1.9)
Number of procedures	56
Ileocystoplasty	49 (87.5)
Ileo-colocystoplasty	2 (3.6)
Colocystoplasty	1 (1.8)
Gastrocystoplasty	1 (1.8)
Ureterocystoplasty	1 (1.8)
Gastric reservoir	2 (3.6)

Table 2. Incidence of postoperative complications following bladder augmentation	
Complications (Total of patients 54)	Incidence n (%)
Decreased renal function	3 (5.6%)
Bladder perforation	2 (3.6%)
Gastric reservoir perforation	1 (1.9%)
Bladder stone formation	8 (14.8%)
Metabolic abnormality	3 (5.6%)
Wound infection	3 (5.6%)
Hematuria	7 (13.0%)
Worsening hydronephrosis	8 (14.8%)

Table 3. The incidence of re-operation secondary to channel and augmentation complications	
	Number (%)
Channel complications	
Total	10/28 (35.7)
Debridement of infection	1/28 (3.6)
Stenotic channel revision	2/28 (7.1)
Dilation/revision of stenotic stoma	7/28 (25.0)
Augmentation complications	
Total	9/54 (16.7)
Repeat augmentation	2/54 (3.7)
Bladder calculi extraction	4/54 (7.4)
Bladder perforation	2/54 (3.7)
Repair of gastric reservoir leak	1/54 (1.9)

Table 4. Timing of postoperative complications for augmentation cystoplasty	
Complication	Mean time to complication (range)
Stomal stenosis	21 months (6–64)
Bladder calculi	58 months (36–72)
Metabolic abnormalities	22 months (8–37)
Worsening hydronephrosis	34 months (5–83)
Bladder perforation	5 days (3–8)