

Can negative ureteroscopy be predicted in ureteral stone treatment?

Mehmet Oguz Sahin, MD¹; Volkan Sen, MD, FEBU¹; Bora Irer, MD²; Guner Yildiz, MD³

¹Manisa State Hospital, Department of Urology, Manisa, Turkey; ²Izmir Metropolitan Municipality Esrefpasa Hospital, Department of Urology, Izmir, Turkey; ³Dr Suat Seren Chest Diseases and Surgery Training and Research Hospital, Department of Urology, Izmir, Turkey

Cite as: *Can Urol Assoc J* 2020;14(5):E209-13. <http://dx.doi.org/10.5489/cuaj.6026>

Published online November 29, 2019

Abstract

Introduction: We aimed to evaluate factors predictive of negative ureteroscopy (URS) in ureteral stones.

Methods: Patients who underwent URS between January 2007 and June 2018 were included in the study. Patients were divided into two groups: group 1 – positive URS (841 patients); and group 2 – negative URS (75 patients). These two groups were compared in terms of demographic data, stone characteristics, and postoperative outcomes.

Results: The mean age of the study patients was 44.5±15.1 years. The absence of collecting system dilatation due to the present stone was found to be a significant predictive factor for negative URS in univariate analysis, but there was no significant difference in multivariate analysis. In the multivariate analysis, low body mass index (BMI), no history of stone surgery, stone located in the distal ureter, small stone area, longer time between the last imaging procedure and URS, and medical expulsive therapy (MET) application were statistically significant in predicting negative URS.

Conclusions: In this study, the parameters that significantly predicted negative URS were found to be low BMI, no history of stone surgery, distal localization of the stone, small stone area, longer time between the last imaging procedure and URS, and MET applied for the current stone. These parameters should be considered to avoid negative URS and patients should be informed of the possibility of negative URS prior to operation.

Introduction

Urolithiasis is seen in approximately 15% of the population and is known to be caused by ureteral stones in 20% of cases.¹ In recent years, non-contrast computed tomography (NCCT) has become the preferred method for the diagnosis of ureteral stones, with a sensitivity of almost 100%.² However, due to its mutagenic and carcinogenic side effects depending on the amount of radiation exposure, it is recommended that patients undergo repeated computed tomography (CT) only at certain intervals.³ The main treatment modalities of

ureteral stones are medical expulsive therapy (MET), ureteroscopy (URS), and extracorporeal shock wave lithotripsy (ESWL), as well as open or laparoscopic ureterolithotomy (only rarely employed in recent years).⁴ URS is the most commonly used method of operation and despite the increasing experience of surgeons, complications related to URS or anesthesia are still encountered. In the early period, ureteral injuries, sepsis, and even death may occur, while ureteral stenosis may develop in the late period.⁵ Considering such negative conditions, MET presents a significant method for the treatment of stones ≤10 mm in particular. In MET, stone expulsion time varies between four and six weeks, and if it is decided that the stone is not likely to be passed at the end of the waiting period, URS may be required.⁶ However, there is no consensus on the imaging techniques to be used to determine the final status of the stone during or within the waiting period. Although these cases can be evaluated by ultrasonography (US), intravenous urography (IVU), kidney-ureter-bladder (KUB), or CT, each of these modalities have disadvantages. In addition, it is not possible to use KUB in the followup of non-radiopaque ureteral stones. Despite all imaging modalities and the feedback received concerning whether the patient was able to pass the stone, the stone present in the urinary system is not visualized in 3.8–9.8% of all URS procedures, known as negative URS.⁷⁻⁹

In this study, we investigated the predictive factors in cases that were pre-diagnosed with proximal or distal ureteral stones and were planned to receive an endoscopic ureteral stone treatment with URS, but were found to have no stones in the urinary system during the procedure and, thus, were reported as negative URS.

Methods

Following the approval of the local ethics committee, we retrospectively evaluated 841 patients that underwent endoscopic ureteral stone therapy after being diagnosed with a ureteral stone and found to have one on URS (positive URS), and 75 patients diagnosed with a ureteral stone that could not be visualized on URS and were reported as diagnostic URS (negative URS) in our clinic between January 2007 and

June 2018. We assessed demographic data, history of urinary system stone disease, localization and other imaging data of the ureteral stone, whether MET was applied, time between the last imaging procedure and URS, and operative results. Alpha adrenergic receptor blockers (tamsulosin and silodosin) were used for MET. Tamsulosin 0.4 mg once a day or silodosin 8 mg once a day were used for three weeks and the patients were asked to present to the emergency service when needed for the management of pain, drink at least 2.5–3 L/day water, be mobile, and strain their micturition into a glass to catch the stones spontaneously passed. The patients were divided into two groups as positive URS (group 1) and negative URS (group 2) for comparison. For the diagnosis of ureteral stones, NCCT (300 mA, 130 kV, 16 slice, Alexion-Toshiba®, Japan) or US together with KUB were used. For the calculation of the stone area, the maximum width and length of the stone in mm were multiplied and the result was obtained as mm². One of the alpha receptor blockers was prescribed to the patients who underwent MET. In addition, 50 mg/day diclofenac tablets were given orally. The followup assessments were undertaken using KUB, US, or CT. A semi-rigid ureteroscope was used in all operations. For all negative URS cases, the absence of a stone was confirmed by a postoperative NCCT.

Patients with multiple or bilateral ureteral stones, those requiring staged procedures, those suspected to have perioperative migration of the stone to the kidney, and those that had previously received a double J stent were excluded from the study.

For the analysis of the data, the Statistical Package for the Social Sciences (SPSS, Inc., Chicago, IL) v.22 was used. The data were presented as mean \pm standard deviation (SD), number (n) and percentages (%). P values of <0.05 were considered statistically significant. Student's t-test was conducted to compare the continuous variables between the groups and the X² test for the comparison of categorical variables. The parameters that were found to statistically significantly difference between the groups according to the univariate analysis were further examined using a multivariate analysis to determine their effect on predicting negative URS.

Results

A total of 916 patients, 630 (68.8%) male and 286 (31.2%) female, were included in the study. The mean age of these patients was 44.5 \pm 15.1 years, the mean body mass index (BMI) was 26.3 \pm 3.3 kg/m², and the mean stone area was 68.0 \pm 51.8 mm². Of the stones, 299 (32.6%) were proximal and 617 (67.4%) were in the distal ureter. URS revealed a ureteral stone in 841 patients (91.8%) but no stone was visualized in 75 patients (8.2%) during this procedure. The patients' general demographic data, history of urinary system stone disease, and other data related to the current stone and applied procedure are shown in Table 1.

Table 1. General demographic data and stone characteristics

Age (years)	44.5 \pm 15.1
Gender	
Male	630 (68.8%)
Female	286 (31.2%)
BMI (kg/m ²)	26.3 \pm 3.3
Systemic disease	210 (22.9%)
Stone disease history	380 (41.5%)
ESWL history	88 (9.6%)
Stone passage history	376 (41.0%)
Stone area (mm ²)	68.0 \pm 51.8
Stone localization	
Proximal	299 (32.6%)
Distal	617 (67.4%)
Collecting system dilatation	673 (73.5%)
Operative time (min)	34.8 \pm 19.1
Hospitalization time (days)	1.4 \pm 1.3
URS	
Negative	75 (8.2%)
Positive	841 (91.8%)

BMI: body mass index; ESWL: extracorporeal shock wave lithotripsy; URS: ureteroscopy.

There were 582 (69.2%) male and 259 (30.8%) female patients with a mean age of 44.5 \pm 15.0 years in group 1, and 48 (64.0%) male and 27 (36.0%) female patients with a mean age of 44.8 \pm 15.6 years in group 2, with no statistically significant difference between the groups ($p=0.857$ and $p=0.364$, respectively). When the two groups were compared in terms of systemic disease, stone disease history, stone expulsion history, laterality of the current stone, and preoperative NCCT evaluation, no significant difference was observed ($p=0.886$, $p=0.808$, $p=0.903$, $p=0.547$, $p=0.800$ respectively). The mean BMI was 26.4 \pm 3.3 kg/m² in group 1, which was significantly higher compared to group 2 (25.2 \pm 2.8 kg/m²) ($p=0.003$). The presence of a history of ESWL was significantly higher in group 1 ($n=86$; 10.2%) than in group 2 ($n=2$; 2.7%) ($p=0.038$). Similarly, the history of stone surgery was significantly higher in group 1 ($n=77$; 9.2%) compared to group 2 ($n=1$; 1.3%) ($p=0.016$). When the groups were compared in relation to ureteral localization of the current stone, distal stones were more common in group 2 ($n=67$; 89.3%) compared to group 1 ($n=550$; 65.4%) ($p<0.001$). Collecting system dilatation was higher in group 1 ($n=631$; 75.0%) than in group 2 ($n=42$; 56.0%) ($p=0.001$). The stone area was also significantly larger in group 1 (70.9 \pm 52.6 mm² vs. 35.5 \pm 22.7 mm²; $p<0.001$). The time between the last imaging procedure and URS was 4.6 \pm 4.9 days in group 1 and 12.1 \pm 7.2 days in group 2, with a statistically significant difference ($p<0.001$). A higher percentage of patients in group 2 ($n=20$; 26.7%) were found to have received MET for the current stone compared to group 1 ($n=23$; 2.7%) ($p<0.001$). The results of intergroup comparisons are shown in Tables 2, 3.

Table 2. Comparison of the groups in terms of demographic data and history of stone disease

	Group 1 Positive URS (n=841)	Group 2 Negative URS (n=75)	p
Gender			0.364
Male	582 (69.2%)	48 (64.0%)	
Female	259 (30.8%)	27 (36.0%)	
Age (years)	44.5±15.0	44.8±15.6	0.857
BMI (kg/m ²)	26.4±3.3	25.2±2.8	0.003
System disease			0.886
Absent	647 (76.9%)	59 (78.7%)	
Present	194 (23.1%)	16 (21.3%)	
Stone disease history			0.808
Absent	491 (58.4%)	45 (60.0%)	
Present	350 (41.6%)	30 (40.0%)	
Stone passage history			0.903
Absent	495 (58.9%)	45 (60.0%)	
Present	346 (41.1%)	30 (40.0%)	
Stone surgery history			0.016
Absent	764 (90.8%)	74 (98.7%)	
Present	77 (9.2%)	1 (1.3%)	

BMI: body mass index; URS: ureteroscopy.

The absence of collecting system dilatation due to the present stone was found to be a significant predictive factor for negative URS in univariate analysis, but there was no significant difference in the multivariate analysis. In the multivariate analysis, the parameters of a low BMI, no history of stone surgery, stone being located in the distal ureter, small stone area, longer time between the last imaging procedure and URS, and MET application were statistically significant in predicting negative URS (Table 4).

Discussion

Although negative URS is relatively rare, it is a disturbing phenomenon for both the patient and the urologist. Performing a negative procedure may result in unnecessary complications and costs. The main reason for encountering negative URS is that the diagnosis and treatment of ureteral stones depend very much on preferences and are affected by several factors related to the patient, surgeon, and equipment. It is possibly for these reasons that the guidelines related to the diagnosis and followup of ureteral stones, the radiographic modality to be used, intervals of followup, and when to refer to surgery remain unclear. In addition, it is known that exposure to radiation presents with many risks, especially skin injury, cataract, malignancy, and chromosome damage.¹⁰ This has led to a shift in the first-choice imaging technique for ureteral stones from NCCT, a radiation imaging modality, to US in recent years.¹¹ However, as the stone size decreases, the sensitivity of US is also reduced. In addition, NCCT can cause more errors in calculating the size of the stone, and there may also be unnecessary aggressive interventions.¹² It

Table 3. Comparison of the groups in terms of the current stone characteristics

	Group 1 Positive URS (n=841)	Group 2 Negative URS (n=75)	p
Ureteral localization			<0.001
Proximal	291 (34.6%)	8 (10.7%)	
Distal	550 (65.4%)	67 (89.3%)	
Laterality			0.547
Right	402 (47.8%)	39 (52.0%)	
Left	439 (52.2%)	36 (48.0%)	
Stone area (mm ²)	70.9±52.6	35.5±22.7	<0.001
Preoperative NCCT			0.800
Absent	287 (34.1%)	27 (36.0%)	
Present	554 (65.9%)	48 (64.0%)	
Collecting system dilatation			0.001
Absent	210 (25.0%)	33 (44.0%)	
Present	631 (75.0%)	42 (56.0%)	
MET application			<0.001
Absent	818 (97.3%)	55 (73.3%)	
Present	23 (2.7%)	20 (26.7%)	
Time between the last imaging procedure and URS (days)	4.6±4.9	12.1±7.2	<0.001

MET: medical expulsive therapy; NCCT: non-contrast computed tomography; URS: ureteroscopy.

has previously been reported that the Doppler URS measurement of the urinary jet flow from the ureteral orifices within the bladder could be used to determine the presence of stones in the ureter and predict spontaneous passage.¹³ Most urologists have also adopted the strategy of combining USG with KUB to minimize cumulative radiation exposure due to radiation, which is 4.3–6.5 mSv in NCCT, but only 0.2–0.7 mSv in KUB.^{14,15} In addition, low-dose CT has been shown to provide successful results, similar to standard-dose CT in the diagnosis of ureteral stones.¹⁶

In their study conducted with pregnant women suspected to have urolithiasis, White et al reported a negative URS rate of 14%. The rate of negative URS was found to be 23%, 4.2%, and 20% in cases that underwent renal US alone, renal US with low-dose CT, and renal US with magnetic resonance urography, respectively.¹⁷ In another study, Youssef et al evaluated the patients scheduled for surgery due to a ureteral stone using KUB, NCCT, or both on the day of operation. The authors noted that 14% of their patients did not have any stones, and their approach avoided unnecessary URS.¹⁸ In addition to these studies, there are three studies in the literature reporting a negative URS rate in the range of 3.8–8%.⁷⁻⁹ In the current study, negative URS was detected in 8.2% of the patients, which is consistent with the literature.

Lambert et al determined the negative URS rate to be higher in women than in men. Although the authors could not clearly explain the reason for this, they noted that pelvic phleboliths and parenchymal calcifications associated with

Table 4. Outcomes of multivariate analysis for predictive factors associated with negative URS

	OR	95% CI	p
BMI	0.863	0.782–0.953	0.004
Stone surgery history	0.098	0.011–0.870	0.037
Ureteral localization (distal ureter)	2.810	1.192–6.622	0.018
Stone area	0.964	0.950–0.978	<0.001
Collecting system dilatation	0.783	0.421–1.454	0.438
MET application	4.251	1.829–9.877	0.001
Time between the last imaging procedure and URS	1.193	1.140–1.248	<0.001

BMI: body mass index; CI: confidence interval; MET: medical expulsive therapy; OR: odds ratio; URS: ureteroscopy.

medullary sponge kidney were more common in women and might have been confused with a stone. They also stated that women might not notice passage of stone because of urinating in a sitting position and having less dysuria complaints due to their shorter urethra.⁸ In the present study, there were a higher number of female patients in the negative URS group, but we found no statistically significant difference.

Katafigiotis et al reported that female gender, presence of a non-radiopaque stone that cannot be detected on KUB, and smaller stone surface area as significant predictive markers for negative URS.⁹ In our study, the groups did not significantly differ in terms of stone disease and stone passage history. This may be because the patients were able to pass the stone more easily or there were a greater number of non-radiopaque stones. The higher incidence of non-radiopaque stones might be another reason for the ill-informed decisions concerning stone presence on preoperative KUB. In addition, the lack of a difference between the groups in terms of ESWL application for the current stones may also indicate that there was no difference between the groups concerning the distribution of opaque and non-opaque stones, although such a differentiation was not made for all the current stones. In addition, the statistically balanced distribution of patients with a preoperative NCCT diagnosis, i.e., this parameter not resulting in a significant difference, can explain why there was no difference between the groups in relation to the accuracy of stone diagnosis. Furthermore, despite the absence of a significant difference between the groups regarding the history of stone disease or stone expulsion, we found a higher rate of negative URS among patients without a history of stone surgery, which can be attributed to the possibility that these patients had passed smaller stones easier.

Kreshover et al suggested that the presence of stones located in the distal ureter and a smaller stone size, which had also been reported by Katafigiotis et al, were significant data for predicting negative URS.^{7,9} Similarly, in our study, we found a higher rate of negative URS in the presence of distal ureteral stones or in ureteral stones with a smaller area. We believe that this finding is associated with the increased possibility of spontaneous stone passage.

Spontaneous passage through conservative treatment is 71–98% for distal ureteral stones of <5 mm and 25–53% for 5–10 mm stones.¹⁹ In addition, for proximal, mid, and distal localization of ureteral stones, the spontaneous passage rate is reported to be 22%, 46%, and 71%, respectively.²⁰ Therefore, the waiting period for stone expulsion usually ranges from four to six weeks. In our study, with the increase of the probability of spontaneous passage of current stones through MET, the rate of negative URS increased. Furthermore, the higher negative URS rates, in parallel with the increased time between the last imaging procedure and URS and absence of preoperative pain, can be explained by the linear relationship between the likelihood of spontaneous stone passage and duration.^{21,22}

We found that the possibility of negative URS was increased in cases with a lower BMI. This may be because these patients were more mobile or their stone diameter was smaller. In contrast to the increased risk of stone-formation due to inactivity and high BMI, physical activity alters the transportation of vitamins and minerals in the body that play an important role in the formation of stones. Exercise stimulates thirst, causes excess fluid intake during exercise, and contributes to the chronic expansion of total body water. In general, these effects result in an increase in circulating blood volume of 20–25%, thus increasing urinary excretion.^{23,24} There are studies indicating no relationship between BMI and urinary stone excretion.²⁵ On the other hand, depending on the exercise mechanism, similar to the reduction of stone-formation, ureteral stone expulsion can be increased.

The lower number of patients with collecting system dilatation in the negative URS group, which was significant in univariate analysis but ineffective according to multivariate analysis, can be attributed to the smaller area and distal localization of the stones in this group. Supporting this finding is the study by Sahin et al, who showed that the success of MET increased in stones with a smaller proximal ureter diameter and lower degree of hydronephrosis.²⁵

The main limitation of this study was its retrospective design. In addition, KUB was not performed in all patients in their initial diagnosis and followup; thus, it was not possible to differentiate between opaque and non-radiopaque stones in all cases. Another issue was that pain was evaluated according to the patients' subjective feedback and no pain scale was used. However, the strength of this study was the high number of patients. Using a data set of this size facilitated statistical analysis of the determinants for negative URS. Our results should be confirmed with further prospective and randomized studies.

Conclusions

In this study, the parameters that significantly predicted negative URS were found to be low BMI, no history of stone

surgery, distal localization of the stone, small stone area, longer time between the last imaging procedure and URS, and MET applied for the current stone. We consider that to avoid negative URS, these parameters should be considered, and the patients should be informed about the possibility of negative URS before operation.

Competing interests: The authors report no competing personal or financial interests related to this work.

This paper has been peer-reviewed.

References

- Dellabella M, Milanese G, Muzzonigro G. Randomized trial of efficacy of tamsulosin, nifedipine, and phloroglucinol in medical expulsive therapy for distal ureteral calculi. *J Urol* 2005;174:167-72. <https://doi.org/10.1097/01.ju.0000161600.54732.86>
- Coe FL, Evan A, Worcester E. Kidney stone disease. *J Clin Invest* 2005;115:2598-608. <https://doi.org/10.1172/JCI26662>
- Brenner DJ, Hall EJ. Computed tomography an increasing source of radiation exposure. *N Engl J Med* 2007;357:2277-84. <https://doi.org/10.1056/NEJMra072149>
- Heers H, Turney BW. Trends in urological stone disease: A 5-year update of hospital episode statistics. *BJU Int* 2016;118:785-9. <https://doi.org/10.1111/bju.13520>
- de la Rosette J, Denstedt J, Geavlete P, et al. The clinical research office of the endourological society ureteroscopy global study: Indications, complications, and outcomes in 11 885 patients. *J Endourol* 2014;28:131-9. <https://doi.org/10.1089/end.2013.0436>
- Skolarikos A. Medical treatment of urinary stones. *Curr Opin Urol* 2018;28:403-7. <https://doi.org/10.1097/MOU.0000000000000523>
- Kreshover JE, Dickstein RJ, Rowe C, et al. Predictors for negative ureteroscopy in the management of upper urinary tract stone disease. *Urology* 2011;78:748-52. <https://doi.org/10.1016/j.urol.2011.03.008>
- Lamberts Remy W, Conti Simon L, Leppert John T, et al. Defining the rate of negative ureteroscopy in the general population treated for upper tract urinary stone disease. *J Endourol* 2017;31:266-71. <https://doi.org/10.1089/end.2016.0751>
- Katafigiotis I, Sabler IM, Heifetz EM, et al. "Stoneless" or negative ureteroscopy: A reality in the endourologic routine or avoidable source of frustration? Estimating the risk factors for a negative ureteroscopy. *J Endourol* 2018;32:825-30. <https://doi.org/10.1089/end.2018.0291>
- Brenner DJ, Hall EJ. Computed tomography an increasing source of radiation exposure. *N Engl J Med* 2007;357:2277-84. <https://doi.org/10.1056/NEJMra072149>
- Ripolles T, Agramunt M, Errando J, et al. Suspected ureteral colic: Plain film and sonography vs. unenhanced helical CT. A prospective study in 66 patients. *Eur Radiol* 2004;14:129-36. <https://doi.org/10.1007/s00330-003-1924-6>
- Hangi L, Fuccai T, Caixia Z, et al. Limited sensitivity and size over measurements of ultrasound affect medical decisions for ureteral stone compared to non-contrast computed tomography. *World J Urol* 2019;37:907-11. <https://doi.org/10.1007/s00345-018-2444-y>
- Ongun S, Teken A, Yilmaz O, et al. Can ureteral jet flow measurement predict spontaneous passage of distal ureteral stones? *Urol Int* 2018;101:156-60. <https://doi.org/10.1159/000490498>
- Hyams ES, Shah O. Evaluation and followup of patients with urinary lithiasis: Minimizing radiation exposure. *Curr Urol Rep* 2010;11:80-6. <https://doi.org/10.1007/s11934-010-0092-x>
- Mitterberger M, Pinggera GM, Pallwein L, et al. Plain abdominal radiography with transabdominal native tissue harmonic imaging ultrasonography vs unenhanced computed tomography in renal colic. *BJU Int* 2007;100:887-90. <https://doi.org/10.1111/j.1464-410X.2007.07048.x>
- Poletti PA, Platon A, Rutschmann OT, et al. Low-dose vs. standard-dose CT protocol in patients with clinically suspected renal colic. *AJR Am J Roentgenol* 2007;188:927-33. <https://doi.org/10.2214/AJR.06.0793>
- White WM, Johnson EB, Zite NB, et al. Predictive value of current imaging modalities for the detection of urolithiasis during pregnancy: A multicenter, longitudinal study. *J Urol* 2013;189:931-4. <https://doi.org/10.1016/j.juro.2012.09.076>
- Youssef FR, Wilkinson BA, Hastie KJ, et al. Is preoperative imaging essential prior to ureteric stone surgery? *Ann R Coll Surg Engl* 2012;94:428-31. <https://doi.org/10.1308/003588412X13373405385458>
- Ibrahim AI, Shetty SD, Awad RM, et al. Prognostic factors in the conservative treatment of ureteric stones. *Br J Urol* 1991;67:358-61. <https://doi.org/10.1111/j.1464-410X.1991.tb15161.x>
- Miller OF, Kane CJ. Time to stone passage for observed ureteral calculi: A guide for patient education. *J Urol* 1999;162:688-91. <https://doi.org/10.1097/00005392-199909010-00014>
- Sáenz Medina J, Alarcón Parra RO, Redondo González E, et al. Prognostic factors of spontaneous expulsion in ureteral lithiasis. *Actas Urol Esp* 2010;34:882-7. <https://doi.org/10.1016/j.acuro.2010.05.006>
- Türk C, Knoll T, Seitz C, et al; European Association of Urology. Medical expulsive therapy for ureterolithiasis: The EAU recommendations in 2016. *Eur Urol* 2017;71:504-7. <https://doi.org/10.1016/j.eururo.2016.07.024>
- Convertino VA. Blood volume: Its adaptation to endurance training. *Med Sci Sports Exerc* 1991;23:1338-48. <https://doi.org/10.1249/00005768-199112000-00004>
- Aune D, Mahamat-Saleh Y, Norat T, et al. Body fatness, diabetes, physical activity, and risk of kidney stones: A systematic review and meta-analysis of cohort studies. *Eur J Epidemiol* 2018;33:1033-47. <https://doi.org/10.1007/s10654-018-0426-4>
- Sahin C, Eryildirim B, Kafkasli A, et al. Predictive parameters for medical expulsive therapy in ureteral stones: A critical evaluation. *Urolithiasis* 2015;43:271-5. <https://doi.org/10.1007/s00240-015-0762-8>

Correspondence: Dr. Mehmet Oguz Sahin, Manisa State Hospital, Department of Urology, Manisa, Turkey; urologoguz@yahoo.com