

**Incidence and mortality trends of metastatic prostate cancer: Surveillance, Epidemiology, and End Results database analysis**

Aaron C. Zhang, BA<sup>1</sup>; Rehana Rasul, MA<sup>2</sup>; Anne Golden, PhD<sup>2</sup>; Michael A. Feuerstein, MD<sup>3</sup>

<sup>1</sup>Donald and Barbara Zucker School of Medicine at Hofstra/Northwell,

Hempstead, NY, United States; <sup>2</sup>Feinstein Institutes for Medical Research, Department of Occupational Medicine, Epidemiology and Prevention, Zucker School of Medicine at Hofstra/Northwell, Hempstead, NY, United States;

<sup>3</sup>Lenox Hill Hospital, Department of Urology, Zucker School of Medicine at Hofstra/Northwell, Hempstead, NY, United States

**Cite as:** Zhang AC, Rasul R, Golden A, et al. Incidence and mortality trends of metastatic prostate cancer: Surveillance, Epidemiology, and End Results database analysis. *Can Urol Assoc J* 2021 June 22; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.7173>

Published online June 22, 2021

**Correspondence:** Dr. Michael A. Feuerstein, Assistant Professor, Zucker School of Medicine at Hofstra/Northwell, Department of Urology, Lenox Hill Hospital, New York, NY, United States; [mfeuerste1@northwell.edu](mailto:mfeuerste1@northwell.edu)

\*\*\*

**Abstract**

**Introduction:** In the past decade, prostate cancer screening decreased, raising the concern of delays in diagnosis and leading to increase in new cases of metastatic prostate cancer. This study evaluated whether these changes may have impacted trends in metastatic prostate cancer incidence and survival.

**Methods:** Metastatic prostate cancer diagnoses from 2008–2016 were identified from the Surveillance, Epidemiology, and End Results (SEER) 18 registries. Age-adjusted incidence rates per 100 000 were calculated by time periods and demographic variables. Two-year all-cause and prostate cancer-specific mortality were calculated for patients diagnosed from 2008–2014, and multivariable Cox proportional hazards models were used to evaluate the impact of demographic and clinical variables.

**Results:** Incidence rates of metastatic prostate cancer increased by 18% from 2008–2009 to 2014–2016 (Incidence rate ratio [IRR]=1.18, 95% confidence interval [CI] 1.14–1.21). This trend was observed across multiple subgroups but was greatest in non-Hispanic Whites and patients living in counties 0–10% below poverty level. There was an overall decreased risk of all-cause and prostate cancer-specific mortality, but unmarried men and men living in counties

>20% below poverty level showed statistically significant increased risk of prostate cancer-specific mortality.

**Conclusions:** Non-Hispanic Whites and the wealthiest subgroups had the largest increase in incidence of metastatic prostate cancer since 2008. Despite trends of decreased risk of prostate cancer-specific mortality, we found certain populations experienced increases in mortality risk. Studies exploring the role of socioeconomic factors on screening and access to newer treatments are needed.

## Introduction

Prostate cancer is the most common cancer in men in the United States with an estimated 191,000 new cases diagnosed in 2020.<sup>1</sup> Despite a 5-year relative-survival rate of greater than 99% for localized disease, metastatic prostate cancer (mPCa) remains the second leading cause of cancer-related death in men with a 5-year relative survival rate of 30.2%.<sup>2</sup> Prior studies have demonstrated there has been an increase in the diagnosis of *de novo* mPCa in the last several years, perhaps related to the United States Preventative Services Task Force (USPSTF) recommendation against routine prostate cancer screening for men of all ages in 2012.<sup>3-5</sup> Considering that race/ethnicity and socioeconomic status have been shown to influence incidence and mortality of cancer, we examined demographic differences in incidence of *de novo* mPCa and survival between 2008-2016.<sup>6,7</sup>

## Methods

### *Data sources and study sample*

Data were queried from the Surveillance, Epidemiology, and End Results (SEER) Program SEER 18 Registries Research Data + Hurricane Katrina Impacted Louisiana cases based on the November 2018 submission using SEER\*Stat 8.3.6 software.<sup>8</sup> SEER is sponsored by the National Cancer Institute to maintain and distribute cancer incidence and survival data from population-based cancer registries representing approximately 28 percent of the U.S. population based on the 2010 census.<sup>9</sup> The SEER registries collect information on site and extent of disease, first course of cancer-directed therapy and socio-demographic characteristics, with active follow-up for date and cause of death.<sup>10</sup> mPCA counts and population estimates generated from the incidence rate session, stratified by year of diagnosis, age group, race/ethnicity, region, poverty level, and stage, were extracted; health insurance status could not be considered for incidence rate analysis because it is not defined in the U.S. population data. Individual data from mPCa cases among males 45 years or older during 2008-2016 were also extracted from the case session. For 2016 cases, SEER classified metastatic stages based on the American Joint Committee on Cancer (AJCC) 7 staging criteria with clinical or pathological indication.<sup>11</sup> Cases with any clinical or pathological indication of metastatic cancer were included. However,

mPCa was defined using AJCC 6 staging criteria (M1a, M1b, M1c) for 2008-2015, which did not differentiate between clinical and pathologic staging.<sup>11</sup> Cases with unknown stage or poverty level, or where survival time was zero months after diagnosis, were excluded.

### *Study variables*

Demographic variables included age group (45-54, 55-64, 65-74, 75-84, 85+ years), race/ethnicity (Non-Hispanic White, Non-Hispanic Black, Non-Hispanic Asian or Pacific Islander, Hispanic (All races)), region (East, Pacific Coast/Alaska, Northern Plains, Southwest), county measure of percent of persons below the federal poverty level (0-10%, >10%-15%, >15%-20%, >20%), and health insurance (insured, uninsured, unknown). The impact of 2012 screening recommendations was examined using time periods of 2008-2009, 2010-2011, 2012-2013, and 2014-2016.

### *Statistical analysis*

Multivariable Poisson regression models for rate analyses were used to calculate mPCA incidence rates and evaluate whether they differed by period, while adjusting for demographic variables and tumor stage. They included an offset using the log (population). Interactions between period and demographic variables or stage were included in separate models to assess whether there were differential trends by these factors. Adjusted incidence rates, presented as the number of mPCa cases per 100,000 persons, as well as adjusted incidence rate ratios (IRR) and Bonferroni-corrected 95% confidence intervals (CI), were reported. The Bonferroni correction using  $p=0.0019$  (21 subgroup comparisons and 5 tests for interaction) was applied due to testing multiple comparisons.

Multivariable Cox proportional hazards models were used to determine whether there was a difference in 2-year all-cause mortality and prostate cancer mortality between time periods and whether it was modified by demographic variables or stage. Time to the event (all-cause mortality and prostate cancer mortality) was the interval in months between diagnosis and the date of death, loss of follow-up, or December 2016, whichever was earliest. This analysis was restricted to patients diagnosed between 2008 and 2014 so that all patients had at least 2 years of follow-up after diagnosis. The time periods 2008-2009, 2010-2011, and 2012-2014 were compared. Because hazards of treatment variables (chemotherapy, radiation, surgery) and stage were not constant over time, they were not included in the main analyses. As a sensitivity to the results, we performed models stratified by these treatment variables and the observed effects of time period and other adjustment variables were similar (Table 2).

mPCa counts and population estimates were extracted from SEER\*Stat software, version 8.3.6 (Surveillance Research Program, National Cancer Institute). All other statistical analyses were performed using SAS 9.4 (SAS Institute Inc., Cary, NC). Statistical significance was defined as  $p < 0.05$ .

## Results

The study cohort consisted of 24,407 men diagnosed with mPCa from 2008-2016 (Table 1). More than half of patients (55.9%) were aged 65-84 years old, non-Hispanic White (64.9%), and married (57.1%). The majority of the patients lived in the Pacific Coast region (50.8%) and the East region (35%). Stage M1b disease was most frequent (74.3%) followed by M1c (19.6%) and M1a (6.2%). Poverty level varied among the cohort, with the highest number of patients falling between 10-15% below the federal poverty level (33.3%).

Multivariable Poisson regression models of incidence trends are presented in Table 2. The increase in incidence trend was observed across several subgroups, including age, race/ethnicity, regions, and poverty levels ( $p < 0.05$  for all interactions). Comparing 2008-2009 to 2014-2016, patients 75-84 years of age showed a 24% increase in the mPCa incidence rate (IRR=1.24, [95% CI=1.15-1.35],  $p < 0.001$ ). Similarly, there was a 31% increase in incidence for patients living in counties 0-10% below federal poverty level (IRR=1.31, [95% CI=1.22-1.40],  $p < 0.001$ ). Patients living in counties >10-15% and >20% below poverty level also showed increases in mPCa incidence of 19% (IRR=1.19, [95% CI=1.18-1.454],  $p < 0.001$ ) and 18% (IRR=1.18, [95% CI=1.06-1.31],  $p < 0.001$ ), respectively.

The impact of interactions between time period and each variable on mortality rates was evaluated in separate survival models. Because none of these interactions were statistically significant ( $p < 0.05$ ), interactions were removed and models were rerun with main effects only, adjusted for all demographic variables (Table 3).

Despite increased incidence rates, men diagnosed with mPCa between 2012-2014 had a lower 2-year cancer-specific mortality (HR=0.93, [95% CI: 0.89-0.97],  $p=0.002$ ) and overall mortality (HR=0.95, [95% CI=0.90-1.00],  $p=0.033$ ) compared to those diagnosed in 2008-2009. Slight decreases in overall and cancer-specific mortality from 2008-2009 until 2010-2011 were not significantly.

## Discussion

Previous reports have shown that the incidence of mPCa increased after 2007, which may be partially attributed to the 2008 USPSTF recommendation against screening in those 75 and older.<sup>5</sup> Advances in imaging and screening techniques in the past decade is another plausible explanation for this trend. However, newer guideline changes in 2012 recommending against routine screening for men of all ages further supports the argument that this trend is most likely related to reduced screening and/or access to care. This study explores whether this trend is similar among different demographic subgroups and the interaction with disease and overall survival.

We found that there was a significant increase in incidence among certain groups, most notably those who were non-Hispanic White, those who were 0-10% below federal poverty level, and those 75-84 years old. We hypothesize that non-Hispanic Whites, men who lived in counties 0-10% below poverty level, i.e., the least impoverished group, and men over 65, the age

that Medicare begins, likely had better access to screening in the past and therefore were impacted the most during this time period.

Previous studies have demonstrated that lower socioeconomic status is associated with diagnoses of more aggressive cancers, such as prostate and breast, both of which have established screening methods.<sup>12,13</sup> After the changes in USPSTF guidelines on prostate cancer screening, patients in the lowest income brackets were likely impacted the least.<sup>14</sup>

Previous studies also described discrepancies in prostate cancer screening utilization in patients of different races. Caucasian men had shorter prostate cancer screening intervals as compared to African American men.<sup>15</sup> Likewise, there were more diagnoses of prostate cancer without prior history of screening in African American men as compared to that of Caucasian men.<sup>16</sup> Interestingly, after controlling for socioeconomic factors, the disparity in prostate cancer mortality from 1992-1999 between African Americans and Caucasians was reduced significantly.<sup>7</sup> The recent trends in incidence of mPCa illustrated in our study further reinforces many of the conclusions drawn from the literature by highlighting certain demographic subgroups that benefited most from screening prior to the USPSTF guideline change.

Our results show that despite increases in mPCa incidence over time, the risk of prostate cancer-specific mortality decreased in later time intervals. Specifically, patients diagnosed between 2012-2014 showed a 7% reduction in prostate cancer-specific mortality, while patients diagnosed between 2010-2011 only showed a 4% reduction in prostate cancer-specific mortality when compared to those diagnosed from 2008-2009. These findings may reflect the recent advances in treatment for prostate cancer and/or improved access to care through the Patient Protection and Affordable Care Act, enacted in 2010. Studies using more granular data including treatment information and insurance status should be considered to examine these trends.

Certain subgroups demonstrated a lower risk of cancer-specific mortality, including those who were insured compared to uninsured and non-Hispanic Asian or Pacific Islander compared to non-Hispanic White. Not only is there an increased likelihood of advanced prostate cancer in uninsured patients compared to insured patients, but also insurance status acts as an important predictor for favorable treatment outcomes in patients with low grade disease.<sup>14</sup> The decreased risk of cancer-specific mortality in patients of non-Hispanic Asian or Pacific Islander background for multiple types of cancer (lung, colorectal, prostate, breast) is less well-described in the literature and warrants further investigation.<sup>17</sup>

Conversely, we also found that certain subgroups demonstrated an increased risk of cancer-specific mortality, notably patients who were unmarried compared to married and who lived in counties >20% below federal poverty level compared to 0-10% below poverty level. Unmarried men may lack the social support systems that are critical for cancer treatment and survival as compared to married men. Prior studies demonstrate marital status as a predictor of cancer mortality and prostate cancer-specific mortality.<sup>18</sup> Using the SEER registry data, unmarried men had higher risk of prostate cancer-specific mortality compared to that of

unmarried men between 1988 and 2003.<sup>19</sup> Men living in areas below poverty level (>20%) likely suffer from lack of access to cancer-related treatments and care, which may contribute to the significantly higher risk of both prostate cancer-specific and all-cause mortality.<sup>20</sup> Although the results of our study show a direct association between percent below poverty level and increased risk of prostate cancer-specific mortality, only the >20% below poverty level group is found to be statistically significant.

Several limitations should be noted that are inherent in the SEER research data and were described previously.<sup>10</sup> First, this study is not able to assess whether screening guidelines were implemented for individual cases and can only make inferences based on time periods. Second, factors such as access to health insurance may influence these trends and were not accounted for in the incidence rate analysis because this is not collected for the general population in the US Census data. Likewise, SEER data does not include important risk factors and clinical factors such as smoking, alcohol intake, diet, family history, and comorbidities that may impact incidence and survival analyses by acting as confounding variables. Finally, this study did not incorporate a time-series design because there were too few years (9 years) for this type of analysis.

It is possible that the increased rates by periods are consistent from 2008-2016 and do not represent a change in the existing trend. It is also possible that there may be lagged effects. For example, in some subgroups, rates for 2012-2013 were not markedly higher than 2010-2011 rates. This could reflect the early adoption in the clinical community for restrictive screening guidelines before they were officially recommended in 2012. Nonetheless, the fact that only certain groups seem to be driving the increasing trend in mPCa incidence rates is worth noting. Furthermore, in order to ensure that all patients had at least 2 years of follow-up from diagnosis of mPCa, the survival analysis was restricted to patients diagnosed between 2008 and 2014. This poses a limitation because the patients had unequal follow-up time and 2-year follow-up time, rather than the more standard 5-year follow up period, may not be adequate for assessing cancer-specific or overall mortality.

## Conclusions

The overall incidence of mPCa increased between 2008 to 2016. We were able to show an increase in incidence of mPCa over time in patients living in the least impoverished areas and in non-Hispanic Whites, reflecting that the increase in diagnoses may be driven by demographic subgroups that benefited most from routine screening prior to the USPSTF recommendation against screening in 2012. We also demonstrated that despite decreased prostate cancer-specific mortality in recent years, perhaps due to the advances in treatment and access to care, certain demographic subgroups saw an increase in risk of prostate cancer-specific mortality. The differences in mortality risk were largely explained by these groups lacking access to resources and social support systems that are critical for favorable treatment outcomes. Future studies

should examine the impact of more recent USPTSF guidelines for reinstating screening and implementation of newer prostate cancer therapies in different subgroups.

DRAFT

## References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin.* 2020;70(1):7-30.
2. National Cancer Institute. Stat Facts: Prostate Cancer. Available at: <https://seer.cancer.gov/statfacts/html/prost.html>. Accessed August 1.
3. Moyer VA, Force USPST. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2012;157(2):120-134.
4. Hu JC, Nguyen P, Mao J, et al. Increase in Prostate Cancer Distant Metastases at Diagnosis in the United States. *JAMA Oncol.* 2017;3(5):705-707.
5. Weiner AB, Matulewicz RS, Eggener SE, Schaeffer EM. Increasing incidence of metastatic prostate cancer in the United States (2004-2013). *Prostate Cancer Prostatic Dis.* 2016;19(4):395-397.
6. Jemal A, Ward EM, Johnson CJ, et al. Annual Report to the Nation on the Status of Cancer, 1975-2014, Featuring Survival. *J Natl Cancer Inst.* 2017;109(9).
7. Du XL, Fang S, Coker AL, et al. Racial disparity and socioeconomic status in association with survival in older men with local/regional stage prostate carcinoma: findings from a large community-based cohort. *Cancer.* 2006;106(6):1276-1285.
8. Surveillance E, and End Results (SEER) Program ([www.seer.cancer.gov](http://www.seer.cancer.gov)) SEER\*Stat Database: Incidence - SEER 18 Regs Research Data + Hurricane Katrina Impacted Louisiana Cases, Nov 2018 Sub (2000-2016) <Katrina/Rita Population Adjustment> - Linked To County Attributes - Total U.S., 1969-2017 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2019, based on the November 2018 submission. .
9. SEER\*Stat Databases: November 2018 Submission. <https://seer.cancer.gov/data-software/documentation/seerstat/nov2018/>. Accessed August 1, 2020.
10. Scosyrev E, Messing J, Noyes K, Veazie P, Messing E. Surveillance Epidemiology and End Results (SEER) program and population-based research in urologic oncology: an overview. *Urol Oncol.* 2012;30(2):126-132.
11. National Cancer Institute. SEER Registrar Staging Assistant. Available at: [https://staging.seer.cancer.gov/tnm/input/1.9/prostate/clin\\_m/?breadcrumbs=\(~schema\\_list~\)](https://staging.seer.cancer.gov/tnm/input/1.9/prostate/clin_m/?breadcrumbs=(~schema_list~)). Accessed August 1, 2020.
12. Clegg LX, Reichman ME, Miller BA, et al. Impact of socioeconomic status on cancer incidence and stage at diagnosis: selected findings from the surveillance, epidemiology, and end results: National Longitudinal Mortality Study. *Cancer Causes Control.* 2009;20(4):417-435.
13. Rundle A, Neckerman KM, Sheehan D, et al. A prospective study of socioeconomic status, prostate cancer screening and incidence among men at high risk for prostate cancer. *Cancer Causes Control.* 2013;24(2):297-303.
14. Clegg LX, Li FP, Hankey BF, Chu K, Edwards BK. Cancer survival among US whites and minorities: a SEER (Surveillance, Epidemiology, and End Results) Program population-based study. *Arch Intern Med.* 2002;162(17):1985-1993.

15. Carpenter WR, Howard DL, Taylor YJ, Ross LE, Wobker SE, Godley PA. Racial differences in PSA screening interval and stage at diagnosis. *Cancer Causes Control*. 2010;21(7):1071-1080.
16. Carpenter WR, Godley PA, Clark JA, et al. Racial differences in trust and regular source of patient care and the implications for prostate cancer screening use. *Cancer*. 2009;115(21):5048-5059.
17. Awasthi S, Gerke T, Williams VL, et al. Interrelationship Between Health Insurance Status and Prostate Cancer Grade Can Have Critical Impact on Prostate Cancer Disease Control: A Retrospective Cohort Study. *Cancer Control*. 2019;26(1):1073274819837184.
18. Aizer AA, Chen MH, McCarthy EP, et al. Marital status and survival in patients with cancer. *J Clin Oncol*. 2013;31(31):3869-3876.
19. Tyson MD, Andrews PE, Etzioni DA, et al. Marital status and prostate cancer outcomes. *Can J Urol*. 2013;20(2):6702-6706.
20. Cheng I, Witte JS, McClure LA, et al. Socioeconomic status and prostate cancer incidence and mortality rates among the diverse population of California. *Cancer Causes Control*. 2009;20(8):1431-1440.

DRAFT

## Figures and Tables

<b>Table 1. Patient characteristics</b>	
<b>Variable</b>	<b>Total (n=24 407)</b>
	<b>n (%)</b>
<b>Age group</b>	
45–54 years	1513 (6.2)
55–64 years	5615 (23.0)
65–74 years	7236 (29.6)
75–84 years	6426 (26.3)
85+ years	3617 (14.8)
<b>Marital status</b>	
Married	13 930 (57.1)
Unmarried	8832 (36.2)
Unknown	1645 (6.7)
<b>Insurance status</b>	
Uninsured	840 (3.4)
Insured	22 709 (93.0)
Insurance status unknown	858 (3.5)
<b>Geographic region</b>	
Alaska, Pacific Coast	12 396 (50.8)
East	8535 (35.0)
Northern Plains	2385 (9.8)
Southwest	1091 (4.5)
<b>Race/ethnicity</b>	
Non-Hispanic White	15 845 (64.9)
Non-Hispanic Black	4271 (17.5)
Non-Hispanic American Indian/Alaska	165 (0.7)
Non-Hispanic Asian or Pacific Islander	1404 (5.8)
Hispanic (All Races)	2722 (11.2)
<b>Poverty level</b>	
0–10%	5060 (20.7)
>10–15%	8136 (33.3)
>15–20%	7583 (31.1)

>20%	3628 (14.9)
Stage	
M1a	1502 (6.2)
M1b	18124 (74.3)
M1c	4781 (19.6)

DRAFT

	Estimated incidence rates (CI) per 100 000				Estimated incidence rate ratios (CI) per 100 000 (ref=2008–2009)		
	2008–2009	2010–2011	2012–2013	2014–2016	2010–2011	2012–2013	2014–2016
Overall	4.22 (3.98–4.46)	4.29 (4.05–4.53)	4.63 (4.37–4.88)	4.97 (4.72–5.22)	1.02 (0.97–1.06)	1.10 (1.05–1.14)	1.18 (1.14–1.22)
Age group							
45–54 years	0.54 (0.47–0.62)	0.57 (0.50–0.65)	0.56 (0.49–0.64)	0.63 (0.56–0.70)	1.05 (0.88–1.27)	1.03 (0.86–1.24)	1.16 (0.98–1.37)
55–64 years	2.54 (2.33–2.75)	2.49 (2.28–2.69)	2.69 (2.48–2.90)	2.87 (2.68–3.06)	0.98 (0.89–1.08)	1.06 (0.96–1.17)	1.13 (1.04–1.23)
65–74 years	5.44 (5.01–5.88)	5.86 (5.41–6.30)	6.06 (5.62–6.49)	6.55 (6.15–6.96)	1.08 (0.98–1.18)	1.11 (1.02–1.21)	1.20 (1.11–1.3)
75–84 years	10.14 (9.34–10.95)	10.43 (9.61–11.24)	11.27 (10.42–12.11)	12.61 (11.81–13.41)	1.03 (0.94–1.13)	1.11 (1.02–1.21)	1.24 (1.15–1.35)
85+ years	18.10 (16.35–19.85)	16.83 (15.21–18.45)	20.22 (18.45–21.98)	19.87 (18.40–21.35)	0.93 (0.82–1.05)	1.12 (1.00–1.25)	1.10 (0.99–1.22)
Poverty level							
0–10%	4.26 (3.94–4.58)	4.44 (4.12–4.76)	4.72 (4.39–5.05)	5.08 (4.77–5.39)	1.04 (0.96–1.13)	1.11 (1.02–1.20)	1.19 (1.11–1.28)
>10–15%	4.2 (3.89–4.51)	4.07 (3.77–4.37)	4.4 (4.08–4.71)	4.63 (4.34–4.92)	0.97 (0.89–1.05)	1.05 (0.97–1.13)	1.10 (1.03–1.18)
>15–20%	4.31 (3.89–4.74)	4.36 (3.94–4.78)	4.83 (4.41–5.25)	5.10 (4.73–5.47)	1.01 (0.89–1.14)	1.12 (1–1.26)	1.18 (1.06–1.31)
>20%	3.97 (3.59–4.35)	4.24 (3.85–4.63)	4.58 (4.18–4.98)	5.19 (4.82–5.56)	1.07 (0.95–1.20)	1.15 (1.03–1.29)	1.31 (1.18–1.45)
Race/ethnicity							

## SEER: Incidence &amp; mortality trends of metastatic PCa

Hispanic (all races)	4.21 (3.76–4.66)	4.14 (3.71–4.56)	4.45 (4.03–4.87)	4.21 (3.89–4.53)	0.98 (0.85–1.13)	1.06 (0.92–1.21)	1.00 (0.88–1.13)
Non-Hispanic American Indian/Alaska	4.42 (2.56–6.29)	4.11 (2.38–5.84)	5.09 (3.24–6.94)	5.02 (3.60–6.45)	0.93 (0.51–1.69)	1.15 (0.66–2.01)	1.14 (0.68–1.89)
Non-Hispanic Asian or Pacific Islander	2.78 (2.38–3.17)	2.20 (1.86–2.53)	2.6 (2.25–2.95)	2.79 (2.51–3.07)	0.79 (0.65–0.97)	0.94 (0.77–1.13)	1.00 (0.85–1.19)
Non-Hispanic Black	9.45 (8.64–10.26)	9.54 (8.75–10.32)	9.52 (8.76–10.28)	9.88 (9.24–10.51)	1.01 (0.90–1.13)	1.01 (0.90–1.12)	1.05 (0.95–1.15)
Non-Hispanic White	3.58 (3.41–3.76)	3.74 (3.56–3.93)	4.09 (3.90–4.28)	4.54 (4.36–4.72)	1.04 (0.98–1.11)	1.14 (1.08–1.21)	1.27 (1.20–1.33)
Region							
East	3.78 (3.50–4.05)	4.01 (3.73–4.29)	4.15 (3.87–4.44)	4.59 (4.32–4.87)	1.06 (0.98–1.15)	1.1 (1.02–1.19)	1.22 (1.13–1.31)
Northern Plains	4.15 (3.66–4.65)	4.61 (4.10–5.13)	4.84 (4.32–5.36)	5.35 (4.88–5.82)	1.11 (0.95–1.29)	1.17 (1.00–1.35)	1.29 (1.13–1.47)
Alaska, Pacific Coast	4.91 (4.60–5.22)	4.73 (4.44–5.03)	5.23 (4.91–5.54)	5.48 (5.19–5.77)	0.96 (0.90–1.03)	1.06 (1.00–1.13)	1.12 (1.05–1.18)
Southwest	3.46 (2.85–4.08)	3.75 (3.13–4.37)	4.57 (3.90–5.24)	4.8 (4.25–5.35)	1.08 (0.86–1.37)	1.32 (1.06–1.65)	1.39 (1.13–1.70)
Stage							
M1a	1.04 (0.88–1.20)	1.13 (0.96–1.29)	1.24 (1.08–1.41)	1.68 (1.52–1.84)	1.09 (0.89–1.34)	1.2 (0.98–1.46)	1.62 (1.36–1.93)
M1b	14.24 (13.41–15.06)	15.38 (14.52–16.25)	16.65 (15.74–17.55)	17.79 (16.91–18.67)	1.08 (1.02–1.14)	1.17 (1.11–1.23)	1.25 (1.19–1.31)
M1c	5.02 (4.62–5.42)	4.10 (3.75–4.44)	4.35 (4.00–4.7)	4.42 (4.12–4.72)	0.82 (0.74–0.90)	0.87 (0.79–0.95)	0.88 (0.81–0.96)

SEER: Surveillance, Epidemiology, and End Results

<b>Table 3. Multivariable Cox proportional hazards models for all-cause and prostate cancer-specific mortality in men with metastatic prostate cancer, SEER 2008–2014</b>				
	<b>All-cause mortality</b>		<b>Prostate cancer mortality</b>	
	<b>HR<sup>†</sup></b>	<b>p</b>	<b>HR<sup>†</sup></b>	<b>p</b>
<b>Period (ref=2008–2009)</b>				
2010–2011	0.97 (0.92–1.02)	0.2	0.96 (0.92–1.01)	0.09
2012–2014	0.95 (0.90–1.00)	0.03	0.93 (0.89–0.97)	0.002
<b>Age group (ref=20–54 years)</b>				
55–64 years	0.93 (0.85–1.01)	0.08	0.99 (0.91–1.07)	0.7
65–74 years	0.96 (0.89–1.05)	0.4	1.10 (1.01–1.19)	0.02
75–84 years	1.27 (1.16–1.38)	<0.0001	1.52 (1.41–1.65)	<0.0001
85+ years	1.80 (1.64–1.97)	<0.0001	2.31 (2.13–2.51)	<0.0001
<b>Insurance status (ref=uninsured)</b>				
Insured	0.85 (0.77–0.95)	0.003	0.88 (0.80–0.97)	0.01
Insurance status unknown	0.88 (0.76–1.02)	0.1	0.91 (0.80–1.04)	0.2
<b>Marital status (ref=married)</b>				
Unmarried	1.20 (1.15–1.26)	<0.0001	1.23 (1.19–1.28)	<0.0001
Unknown	0.98 (0.90–1.06)	0.6	0.99 (0.92–1.06)	0.7
<b>Geographic region (ref=East)</b>				
Alaska, Pacific Coast	1.04 (0.99–1.09)	0.1	1.01 (0.97–1.05)	0.8
Northern Plains	0.99 (0.92–1.06)	0.7	1.00 (0.94–1.06)	0.9
Southwest	0.97 (0.87–1.07)	0.5	0.94 (0.86–1.03)	0.2
<b>Race/ethnicity (ref=White)</b>				
Non-Hispanic Black	1.02 (0.96–1.08)	0.5	1.04 (0.99–1.09)	0.1
Non-Hispanic American Indian/Alaska Native	1.04 (0.81–1.32)	0.8	1.04 (0.84–1.28)	0.7

Non-Hispanic Asian or Pacific Islander	0.73 (0.67–0.81)	<0.0001	0.77 (0.71–0.84)	<0.0001
Hispanic (All Races)	0.94 (0.88–1.01)	0.07	0.95 (0.89–1.00)	0.06
Poverty level (ref=0–10%)				
>10–15%	1.01 (0.96–1.07)	0.7	1.00 (0.95–1.05)	1.0
>15–20%	1.04 (0.98–1.11)	0.2	1.04 (0.99–1.09)	0.17
>20%	1.09 (1.02–1.17)	0.01	1.10 (1.04–1.17)	0.002

†Numbers represented as hazard ratio with 95% confidence intervals adjusted for age, race (White, Black, other), year of prostate cancer diagnosis, poverty level, marital status, insurance status, and geographic region of SEER registry. HR: hazard ratio; SEER: Surveillance, Epidemiology, and End Results.

DRAFT

<b>Supplementary Table 1. Sensitivity analysis of multivariable Cox proportional hazards models stratified by treatment variables for all-cause and prostate cancer-specific mortality in men with metastatic prostate cancer, SEER 2008–2014</b>				
	<b>All-cause mortality</b>		<b>Prostate cancer mortality</b>	
	<b>HR<sup>†</sup></b>	<b>p</b>	<b>HR<sup>†</sup></b>	<b>p</b>
<b>Period (ref=2008–2009)</b>				
2010–2011	0.96 (0.92–1.01)	0.1	0.97 (0.92–1.02)	0.2
2012–2014	0.94 (0.90–0.98)	0.006	0.95 (0.91–1.00)	0.06
<b>Age group (ref=20–54 years)</b>				
55–64 years	0.99 (0.92–1.08)	0.9	0.94 (0.86–1.02)	0.1
65–74 years	1.09 (1.01–1.19)	0.03	0.97 (0.89–1.06)	0.5
75–84 years	1.51 (1.39–1.64)	<0.0001	1.27 (1.16–1.39)	<0.0001
85+ years	2.29 (2.11–2.49)	<0.0001	1.81 (1.65–1.99)	<0.0001
<b>Insurance Status (ref=uninsured)</b>				
Insured	0.91 (0.83–1.00)	0.06	0.88 (0.79–0.98)	0.02
Insurance status unknown	0.93 (0.81–1.06)	0.3	0.90 (0.79–1.05)	0.2
<b>Marital Status (ref=married)</b>				
Unmarried	1.22 (1.18–1.27)	<0.0001	1.20 (1.15–1.25)	<0.0001
Unknown	0.98 (0.91–1.05)	0.5	0.97 (0.89–1.06)	0.5
<b>Geographic region (ref=East)</b>				
Alaska, Pacific Coast	1.00 (0.96–1.04)	0.9	1.03 (0.98–1.08)	0.2
Northern Plains	0.99 (0.93–1.05)	0.8	0.98 (0.91–1.05)	0.6
Southwest	0.93 (0.85–1.02)	0.1	0.96 (0.86–1.06)	0.4
<b>Race/ethnicity (ref=White)</b>				
Non-Hispanic Black	1.03 (0.98–1.08)	0.3	1.01 (0.96–1.07)	0.7
Non-Hispanic American Indian/Alaska Native	1.03 (0.83–1.27)	0.8	1.04 (0.81–1.32)	0.8

Non-Hispanic Asian or Pacific Islander	0.77 (0.71–0.83)	<0.0001	0.73 (0.66–0.80)	<0.0001
Hispanic (all races)	0.94 (0.89–1.00)	0.05	0.94 (0.88–1.00)	0.07
Poverty level (ref=0–10%)				
>10–15%	1.01 (0.96–1.06)	0.8	1.02 (0.96–1.08)	0.5
>15–20%	1.03 (0.98–1.09)	0.2	1.04 (0.98–1.10)	0.2
>20%	1.11 (1.04–1.18)	0.001	1.10 (1.02–1.18)	0.01

†Numbers represented as hazard ratio with 95% confidence intervals adjusted for age, race (White, Black, other), year of prostate cancer diagnosis, poverty level, marital status, insurance status, and geographic region of SEER registry. HR: hazard ratio; SEER: Surveillance, Epidemiology, and End Results.

DRAFT