

Fosfowash: Early proof of concept study investigating intravesical fosfomycin for recurrent urinary tract infections

Blayne Welk, MD, MSc¹; Christopher Munday, PhD²; Jeremy Burton, MSc, PhD³; Richard J. Baverstock, MD⁴

¹Department of Surgery and Epidemiology and Biostatistics, Western University, London, ON, Canada; ²Lawson Research Institute, London, ON, Canada; ³Departments of Surgery and Microbiology & Immunology, Western University, London, ON, Canada; ⁴Department of Surgery, Cumming School of Medicine, University of Calgary, AN, Canada

Cite as: Welk B, Munday C, Burton J, et al. Fosfowash: Early proof of concept study investigating intravesical fosfomycin for recurrent urinary tract infections. *Can Urol Assoc J* 2021;15(12):423-4. <http://dx.doi.org/10.5489/cuaj.7365>

Published online June 22, 2021

Introduction

Patients with neurogenic lower urinary tract dysfunction (NLUTD) rate urinary tract infections (UTIs) as one of the most bothersome problems.¹ This is a frustrating, multifactorial problem that often arises due to the use of catheters, abnormal bladder compliance, coexisting neurogenic bowel dysfunction, multidrug-resistant organisms, and specific deficiencies in host defense mechanisms.² The management of NLUTD patients who are plagued by an abnormally high frequency or severity of UTIs and are using clean intermittent catheters (CIC) may require tertiary treatment options to try and prevent UTIs. Clinicians often struggle with the potential harms of continuous oral antibiotic prophylaxis (such as antibiotic resistance) and this approach is often considered a last resort.^{3,4}

Intravesical antibiotic bladder instillation is an alternative option that is well-suited to patients already using CICs. Previous studies have generally focused on the use of gentamicin;⁵⁻⁷ however, it is an expensive medication, it is cumbersome to obtain for patients, and in recent years, it has been the victim of manufacturing delays and drug shortages. Our objective was to investigate if fosfomycin tromethamine, a readily available oral antibiotic that comes in powder form, may be a suitable agent for intravesical bladder instillation therapy.

Methods

Fosfomycin tromethamine is a bactericidal antibiotic with a broad range of activity against gram-positive and negative aerobic bacteria (such as *Escherichia coli*, *Enterococcus*, *Klebsiella*, *Proteus*, *Serratia*, *Pseudomonas*, and *Citrobacter*) that are commonly associated with UTIs. It has in vitro anti-

microbial activity against extended-spectrum beta-lactamase (ESBL)-producing bacteria and can penetrate biofilms. It inactivates the enolpyruvyl transferase enzyme (which inhibits bacterial cell wall synthesis) and reduces adherence of bacteria to the urothelium.^{8,9}

We conducted two in vitro experiments using fosfomycin. We tested three solutions: reconstituted fosfomycin (created by mixing a commercially available package of fosfomycin tromethamine [JAMP Pharma Corporation, QC, Canada] in 125 mL of distilled water), tobramycin 160 mg in 1 L normal saline (positive control), and buffered saline (negative control).

The objective of our first experiment was to determine if reconstituted fosfomycin tromethamine was bactericidal in the absence of metabolism by the human gut (where it is normally metabolized to its free acid, fosfomycin). We added 10 µL of the fosfomycin solution to a tryptic soy agar (TSA) plate of 10⁵ colony-forming units (CFU) of urinary isolate, *E. coli* 67, and the plate was incubated at 37°C aerobically overnight.

The objective of our second experiment was to determine the minimum inhibitory concentration of the fosfomycin and tobramycin solutions; 1 mL of each of the solutions was added to 1.5 mL Eppendorf tubes in triplicate using the following dilutions based on the original solutions (100%, 50%, 25%, and 12.5%). Then 1 × 10⁵ CFU of *E. coli* 67 was added to each tube. Tubes were incubated overnight at 37°C and agitated at 170 rpm. Bacterial survival was assessed the next day by dilution series and drop-plating on TSA plates.

Results

The fosfomycin solution had a similar pH (5.5) to the tobramycin solution (6.5). In the first experiment, fosfomycin and tobramycin were effective at inhibiting the growth of *E. coli* 67, with a zone of inhibition of 31 mm and 20 mm, respectively. This result was replicated after one week of storage at 4°C, showing that there is no loss of efficacy over this time period (data not shown). In our second experiment, all four dilutions of both the tobramycin and the fosfomycin inhibited all bacterial growth of *E. coli* 67; the negative control series had high growth of bacteria (Fig. 1).




Fig. 1. Fosfomycin at the concentration tested was effective at inhibiting the growth of *E. coli* 67 with a zone of inhibition that was similar to that of tobramycin. Results of fosfomycin are shown in panel A, and tobramycin in panel B.

Discussion

A recent systematic review identified 11 previous studies of intravesical antibiotic therapy and found that this approach was generally efficacious at both the treatment and prevention of UTIs, with minimal side effects.¹⁰ This conclusion supports the need to develop further feasible intravesical options. To our knowledge, this is the first investigation into the use of fosfomycin as an intravesical antibiotic. We have shown promising, preliminary data that support the further investigation of fosfomycin as an intravesical antibiotic agent for patients with NLUTD and frequent UTIs. As both the fosfomycin and tobramycin achieved complete bacterial kill in all the tested dilutions, we know the minimal inhibitory concentration for *E. coli* 67 is less than a 12.5% dilution (equivalent to one satchel of commercial fosfomycin in 1 L of water).

Ideally, intravesical antibiotics are instilled before bedtime, at the time of the last catheter of the day and allowed to dwell overnight. Potential advantages of fosfomycin as a bladder instillation include the ability of patients to obtain it from any outpatient pharmacy, the ease of storing a reconstituted satchel in the fridge for use over the course of a week (for example, reconstituting in a 500 mL or 1000 mL of commercial bottled water), and the wide antimicrobial spectrum of fosfomycin. Fosfomycin has often been used as a once-per-week oral therapy for recurrent UTIs, however, using it as an intravesical therapy may have benefits of avoiding systemic side effects, reducing bacterial resistance in the gut flora, and improved patient acceptance compared to continuous oral use. Three patients with NLUTD who were unable to access intravesical aminoglycosides due to the current pandemic switched to using fosfomycin instillations

due to lack of alternatives. Anecdotally, they found it to be tolerable and to have a similar efficacy to their previous tobramycin instillations.

Limitations of this work include the use of a single UTI-relevant bacterial strain, no direct assessment of potential cytotoxic effects (although it is typically well-tolerated orally), and the potential for resistance to develop with long-term use. Further basic science and clinical studies will be necessary to determine clinical efficacy and safety and to test different dilution options and instillations schedules.

Competing interests: The authors do not report any competing personal or financial interests related to this work.

This paper has been peer-reviewed.

References

1. Pannek J. Prevention of recurrent urinary tract infections in neurourology. *Eur Urol Focus* 2020;6:817-9. <https://doi.org/10.1016/j.euf.2020.01.015>
2. Kennelly M, Thiruchelvam N, Averbeck MA, et al. Adult neurogenic lower urinary tract dysfunction and intermittent catheterization in a community setting: Risk factors model for urinary tract infections. *Adv Urol* 2019;2019:2757862-13. <https://doi.org/10.1155/2019/2757862>
3. Fisher H, Oluboyede Y, Chadwick T, et al. Continuous low-dose antibiotic prophylaxis for adults with repeated urinary tract infections (AnTIC): A randomized, open-label trial. *Lancet Infect Dis* 2018;18:957-68. [https://doi.org/10.1016/S1473-3099\(18\)30279-2](https://doi.org/10.1016/S1473-3099(18)30279-2)
4. Blaser M. Antibiotic overuse: Stop the killing of beneficial bacteria. *Nature* 2011;24;476:393-4. <https://doi.org/10.1038/476393a>
5. Abrams P, Hashim H, Tomson C, et al. The use of intravesical gentamicin to treat recurrent urinary tract infections in lower urinary tract dysfunction. *Neurourol Urodyn* 2017;36:2109-16. <https://doi.org/10.1002/nau.23250>
6. Cox L, He C, Bevins J, et al. Gentamicin bladder instillations decrease symptomatic urinary tract infections in neurogenic bladder patients on intermittent catheterization. *Can Urological Assoc J* 2017;11:350-4. <https://doi.org/10.5489/cuaj.4434>
7. Stalenhoeft JE, Nieuwkoop C van, Menken PH, et al. Intravesical gentamicin treatment for recurrent urinary tract infections caused by multidrug resistant bacteria. *J Urol* 2019;201:549-55. <https://doi.org/10.1016/j.juro.2018.10.004>
8. Falagas ME, Vouloumanou EK, Samonis G, et al. Fosfomycin. *Clin Microbiol Rev* 2016;29:321-47. <https://doi.org/10.1128/CMR.00068-15>
9. Maraki S, Samonis G, Rafailidis PI, et al. Susceptibility of urinary tract bacteria to fosfomycin. *Antimicrob Agents Ch* 2009;53:4508-10. <https://doi.org/10.1128/AAC.00721-09>
10. Pietropolo A, Jones P, Moors M, et al. Use and effectiveness of antimicrobial intravesical treatment for prophylaxis and treatment of recurrent urinary tract infections (UTIs): A systematic review. *Curr Urol Rep* 2018;19:78. <https://doi.org/10.1007/s11934-018-0834-8>

Correspondence: Dr. Blayne Welk, Department of Surgery and Epidemiology and Biostatistics, Western University, London, ON, Canada; bkwelk@gmail.com