

## Which renal access technique for PCNL is more difficult to teach using simulation?

Anne Yin<sup>1</sup>, Christian Diab<sup>1</sup>, Sero Andonian<sup>1</sup>, Naeem Bhojani<sup>2</sup>, Ahmad Al Marzouq<sup>1</sup>, Caroline White<sup>3</sup>, Nader Fahmy<sup>1</sup>

<sup>1</sup>Division of Urology, Department of Surgery, McGill University, Montreal, QC, Canada; <sup>2</sup>Division of Urology, Department of Surgery, Université de Montréal, Montreal, QC, Canada; <sup>3</sup>Department of Education, McGill University, Montreal, QC, Canada

**Cite as:** Yin A, Diab C, Andonian S, et al. Which renal access technique for PCNL is more difficult to teach using simulation? *Can Urol Assoc J* 2023;17(5):E128-33. <http://dx.doi.org/10.5489/kuaj.8085>

Published online February 2, 2023

Appendix available at [cuaj.ca](http://cuaj.ca)

### ABSTRACT

**INTRODUCTION:** Percutaneous nephrolithotomy (PCNL) is a challenging procedure that urology trainees should be familiar with during residency. Simulators, such as the PERC Mentor, allow the development of this competency in a safer, stress-free environment. There are two primary fluoroscopic methods of gaining percutaneous renal access: the triangulation method and the bull's eye method. Our goal was to assess which method is easier to teach novices by using the PERC Mentor simulator. A secondary goal was to assess differences in subjective and objective outcomes.

**METHODS:** Fifteen simulator and procedure-naïve medical trainees were randomized into two groups using a crossover, randomized study design. Participants were provided with written, video, in-person demonstrations and hands-on practice for each technique. They then performed each method and were assessed objectively using the PERC Mentor performance data report and subjectively using the PCNL global rating scale (GRS) scoring system. Statistical analysis was performed using Student's T-test and non-parametric Wilcoxon signed rank test.

**RESULTS:** There was no statistical difference in the outcomes and complication rates between the two methods. The bull's eye method of obtaining percutaneous access was associated with a significant decrease in operative time (91 seconds vs. 128 seconds,  $p=0.03$ ) and fluoroscopy time (87 seconds vs. 123 seconds,  $p=0.03$ ) compared to the triangulation method.

**CONCLUSIONS:** Teaching of both techniques was equally well acquired by students. Both techniques had similar outcomes; however, the bull's eye method was associated with less operative and fluoroscopy time when compared to the triangulation method among novices.

### INTRODUCTION

Percutaneous nephrolithotomy (PCNL) is considered the gold standard for treatment of patients with larger renal stones. Gaining percutaneous access (PCA) is an essential, yet challenging step in the PCNL procedure.<sup>1</sup> There are two primary fluoroscopic methods used to obtain PCA: eye of the needle (bull's eye) or triangulation.<sup>2</sup> To date, no significant difference has been shown in terms of outcomes and complication rates when comparing the two fluoroscopic methods.<sup>2,3</sup> Various anatomical considerations exist that make gaining PCA difficult. The kidneys are well vascularized, with many vital structures nearby. Proper orientation and puncture needle placement are therefore necessary to avoid complications such as hemorrhage and visceral injuries.<sup>4,5</sup> The difference of couple of millimeters in needle placement can distinguish between a successful procedure and a complication.<sup>6</sup> Optimal placement also ameliorates the ease of PCNL, increases stone-free rate, and decreases bleeding due to less torque on the renal parenchyma.<sup>7-9</sup>

However, even though 70% of urologists perform percutaneous procedures, only 11–17% actually establish their own access.<sup>5,10</sup> It has been shown that urologists obtaining their own access is associated with lower complication rates and shorter hospitalization stays.<sup>10</sup> Reasons for this discrepancy can be attributed partly to the lack of training in residency, as well as the steep learning curve associated with mastering

## KEY MESSAGES

■ Obtaining percutaneous renal access is a challenging competency for urology trainees to obtain. Both the fluoroscopic techniques of bull's eye and triangulation are equally well taught and received by trainees.

■ Both techniques are associated with similar outcomes and complication rates; however, the bull's eye is associated with a faster operative time and fluoroscopic time compared to the triangulation technique.

such a technique.<sup>5,11</sup> A literature review of 14 articles determined that the learning curve for PCNL procedure plateaued at around 30–60 cases for operative time, 20–115 cases for fluoroscopic screen time, and 30–105 cases for stone-free rates. Successful puncture plateaued at 120 cases and radiation dose plateaued at 115 cases.<sup>12</sup> The introduction of simulators for surgical training allows trainees to address a portion of the discrepancy mentioned above by practicing in a stress-free and radiation-free environment.<sup>13</sup>

Prior to computerized simulators, biological and non-biological bench models, such as porcine or bovine kidneys and 3D printing, were used as training material; however, these methods did not negate the risk of radiation and body fluid exposure to the user.<sup>14,15</sup>

The PERC Mentor (<https://www.okbmedical.com/percmentor.html>) is currently the only virtual reality (VR) simulator that has been validated for training and assessment of PCA skills.<sup>16</sup> With its accurate visual imagery and realistic equipment replicating tactile feedback and movements associated with respiration, the PERC Mentor simulator has been used in many urology training programs to improve PCA skills.<sup>17,18</sup> The PERC Mentor is also capable of generating a performance report for each attempt, looking at operative time, fluoroscopy time, time spent introducing needle into the collecting system, number of attempts to puncture, complications and types, and amount of contrast injected.

PCNL global rating scale (GRS) is an evaluation system, modified from an endourological global rating scale that was validated to measure skill acquisition in ureteroscopy.<sup>13,19</sup> This PCNL-GRS rubric looks at five different parameters: anatomy identification, planned needle puncture, use of instrumentation, ability to per-

form tasks, and overall performance. It is a subjective assessment used in surgical education that has itself been validated to evaluate a trainee's performance.<sup>13,20</sup>

Significant improvement in all facets of the PCNL-GRS score was seen in subjects who underwent training, whereas improvement in only one parameter was seen in those who did not receive training. This demonstrates the ability of the PCNL-GRS score to quantify and differentiate between skill levels.<sup>19</sup>

When the PCNL-GRS score is combined with the objective data obtained by the PERC Mentor, an improvement in 11 out of 14 tested spheres was seen in trained subjects, whereas there was only an improvement in 1 of 14 elements in untrained subjects, reinforcing the pervasiveness of parameters tested when both subjective and objective data reports are combined.<sup>19</sup> Therefore, the PCNL-GRS score, along with the objective data obtained by the PERC Mentor, allows for a global evaluation of trainee skills and has been previously proven to be successful in differentiating competency between urology postgraduate trainees.<sup>19</sup>

Given the difficulty of mastering PCA and the existence of two different yet similarly effective methods of gaining PCA, the goal of our study was to determine which method — triangulation vs. bull's eye — is associated with an easier ability to learn on the PERC Mentor simulator, thus allowing urology trainees to gain competency faster. This is particularly essential in the Canadian urological residency training landscape, where a shift towards a Competency-by Design evaluation model outlines percutaneous renal access as a necessary skill to obtain during residency.<sup>21</sup> This is the first study of its kind comparing the ease of learning of both methods with simulation-naïve medical trainees on the validated PERC Mentor simulator. Our objective was to determine whether there is a significant difference between teaching both the triangulation and the bull's eye method in terms of objective and subjective assessment scores.

## METHODS

Fifteen second-year medical students from one Canadian, urban, medical university, participating in the Surgical Explorations and Discovery (SEAD) program were recruited. This two-week, voluntary program aimed to expose medical students early on to various surgical specialties. All participants were all simulator-naïve and had never performed any steps of a PCNL before. Each student was given access to written instructional material and videos outlining both the triangulation method and the bull's eye method

of obtaining PCA a few days prior to performing the procedure on the PERC Mentor simulator. Simplified written instructions were also provided to the students on the day of performance (Appendix available at [cuaj.ca](#)). They then observed an endourologist explain and demonstrate the steps of both methods on the simulator. Finally, they each were allowed one practice run using both techniques on the PERC Mentor to familiarize themselves with the simulator. Each trainee was observed by a postgraduate year (PGY) 5 urology resident and an endourology attending, who were available to guide the students and answer questions. This PCNL curriculum was designed by two endourologists practicing at academic hospitals affiliated with the trainees' medical school.

Each student then performed both the triangulation method and the bull's eye method of gaining PCA in the renal pole of their choice. Successful puncture was deemed by return of urine. Using a randomized, controlled, crossover study design, the participants were sorted into two groups. Group A got to perform the bull's eye method first, followed by the triangulation method, whereas group B performed the triangulation method first, followed by the bull's eye (Figure 1). This was done to ensure that differences seen between the two methods were not due to confounding factors, such as prior amount of exposure to the simulator. The techniques performed by each participant was compared to each other. The data was paired. Each participant was therefore also their own control.

Participants' performance was judged using the validated PCNL-GRS scoring system, as well as the PERC Mentor's registered performance results. The PCNL-

GRS is a subjective score given by the observing PGY-5 urology resident who is familiar with both techniques of obtaining PCA using a marking sheet. PCNL-GRS is comprised of five components looking at anatomy identification, planned needle puncture, use of instrumentation, ability to perform tasks, and overall performance. Each component is rated on a Likert scale from 1–5, up to a total of 25 points.<sup>20</sup>

The PERC Mentor performance sheet, generated by the simulator, records objective data, including total operative time, X-ray exposure time, time spent introducing needle into the collecting system, number of attempts to puncture the collecting system, number and type of complications, and total amount of retrograde contrast injected.

The results of the triangulation method and the bull's eye method were compared and evaluated using the IBM SPSS software. Paired Student's t-test was used to assess differences in total operative time, X-ray exposure time, time spent introducing needle into the collecting system, and total amount of retrograde contrast injected. Non-parametric Wilcoxon signed rank test was used to compare PCNL-GRS global score, as well as each individual component between the two methods. This test was also used to compare number of attempts to puncture the collecting system and number of complications. Additional statistics were done to assess whether there was an effect between PCA attempt number and group allocation (students being in group A vs. B) on objective and subjective performance results. A two-sided p-value <0.05 was considered significant.

## RESULTS

Fifteen medical students participated in the study; 60% were male and 40% were female. They were all in their second year of medical school.

All 15 participants performed both the triangulation method and the bull's eye method of obtaining PCA. The bull's eye method was associated with a statistically significant decreased fluoroscopy and operative time compared to the triangulation method. On average, the total operative time for the triangulation method was 128 seconds, while that for the bull's eye method was 91 seconds ( $p=0.03$ ). Fluoroscopy (X-ray) exposure time was also statistically significant between the two methods, with the triangulation method at 123 seconds and the bull's eye method at 87 seconds ( $p=0.03$ ).

There was no statistical difference in terms of PCNL-GRS total score, nor each individual component score between the two methods. As well, there was no dif-




Figure 1. Participant allocation methods.

ference in terms of number of attempts to puncture the collecting system, complication rates, and amount of contrast injected during the procedure between the two methods (Table 1).

Secondary analysis was done to see whether there was an appreciable learning effect that could be seen between a trainee's first and second attempt, regardless of the first technique performed. Within the objective measures, operative time and fluoroscopic time were analyzed, given that there were statistically significant differences observed in those two parameters. For subjective measures, only the total PCNL-GRS score was analyzed, given there was no statistical difference between any subcategory, nor total score (Figure 2). Though there was a trend toward improvement with increased PCNL-GRS scores and decreased operative and fluoroscopic times on the second attempt, none of the measures were statistically significant.

Additional analysis looked at whether the group allocation had an influence on the PCNL-GRS total score, as well as the operative time and fluoroscopic time for both the bull's eye and triangulation methods (Table 2). Within each technique, there was no difference between group A and group B for operative time, fluoroscopic time, and total PCNL-GRS score. The order in which each participant performed the technique had no impact on execution.

## DISCUSSION

Contrary to popular belief that teaching triangulation is more difficult than the bull's eye method for obtaining PCA, both methods were well learned by the trainees. Either method can be equally taught and learned without compromising outcomes. Our results, however, demonstrate that the bull's eye method in the hand of novice trainees is a faster and safer technique than the triangulation method. Due to the lesser need for C-arm manipulations, it is associated with a decrease in total operative time, as well as less fluoroscopy time, leading to less radiation exposure for both the patient and the trainee. There was, however, no difference between the two methods in terms of subjective performance, as demonstrated using the PCNL-GRS scoring system, nor in terms of complication rates, attempts to puncture, and amount of radiocontrast material used.

Given the similar success rates and adverse outcomes between both the triangulation and the bull's eye methods, our results show that the bull's eye method is likely an easier skill for trainees to acquire; however, either method can be learned without increased adverse events.


**Table 1. Objective and subjective results of various percutaneous nephrolithotomy methods**

	Percutaneous nephrolithotomy method		
	Triangulation (± SD)	Bull's eye (± SD)	p
<b>PCNL-GRS score</b>			
Total	18.00±4.94	19.38±4.80	0.474
Identify anatomy	4.87±0.52	4.53±0.83	0.102
Plan needle puncture	3.60±1.06	3.93±1.10	0.336
Use of instruments	3.47±1.19	3.79±1.25	0.314
Ability to perform task	2.67±1.59	3.20±1.26	0.2
Overall performance	3.40±1.24	3.67±1.05	0.248
<b>Objective data</b>			
Operative time (s)	128.07±35.53	91.33±40.99	0.033
Fluoroscopy time (s)	123.07±33.63	87.07±39.87	0.031
Time spent introducing needle into collecting system (s)	17.80±9.67	15.00±10.68	0.42
Attempts to puncture (#)	1.87±0.64	2.27±1.10	0.262
Complications (#)	1.73±0.96	1.73±0.80	0.914
Amount of contrast injected (mL)	46.33±11.72	41.00±11.21	0.22

SD: standard deviation.

We then compared the subjective and objective data between a trainee's first and second attempts to see if there was a cumulative learning effect that could be observed. The PCNL-GRS total score, operative time, and fluoroscopy time were the parameters used. Although there was a trend towards a mild improvement in all three facets, showing that learned skills were retained from attempt 1 to attempt 2, this was not statistically significant, likely due to the study being underpowered. Nonetheless, this information can be useful for educators and attendings in teaching hospitals; the bull's eye method could be used as the initial teaching method for novice learners, as it is associated with easier skill acquisition. Once the trainee feels

“ We found no difference in outcomes between the two PCA methods; however, the bull's eye technique, in the hand of novices, is associated with a faster skill acquisition. ”



**Figure 2.** Effect of percutaneous access (PCA) attempts on objective and subjective performance. PCNL: percutaneous nephrolithotomy

**Table 2. Effect of randomized PCA technique sequence on objective and subjective outcomes**

	Group allocation		
	Group 1 (± SD)	Group 2 (± SD)	p
<b>PCNL-GRS total score</b>			
Bull's eye	19.57±5.56	19.14±4.34	0.875
Triangulation	19.57±4.24	16.57±5.80	0.291
<b>Operative time (s)</b>			
Bull's eye	95.71±54.63	87.50±27.67	0.714
Triangulation	127.00±38.99	129.00±34.92	0.918
<b>Fluoroscopy time (s)</b>			
Bull's eye	92.14±53.41	82.63±26.19	0.662
Triangulation	124.14±35.72	122.13±34.14	0.913

PCA: percutaneous access; PCNL-GRS: percutaneous nephrolithotomy global rating scale; SD: standard deviation.

more at ease and has gained confidence in this technique, they can implement the triangulation method. Trainees will likely be more proficient than since they are already familiar with the basic fluoroscopic key steps. By using the PERC Mentor simulator, which has been accepted and validated as a surgical model for PCNL,<sup>20</sup> our results can be extrapolated to the operating room.

To our knowledge, this is the first study that has compared both methods using a virtual-reality model in simulator and PCA procedure-naive trainees. None of our participants have used the PERC Mentor or performed any steps of a PCNL in the operating room prior to the conduction of this study, thus allowing

our results to truly reflect the ease of learning of each method. Although studied on medical students, our results can be useful to guide urology residency curriculum, as all trainees start their learning journey with minimal knowledge. A urology resident's first exposure to percutaneous renal access and percutaneous nephrolithotomy is likely during residency.

A previous study conducted by Abdallah et al, using a bovine kidney as a biological model, compared the triangulation and bull's eye methods among trainees and concluded there was no statistically significant difference between the two models in terms of total operative time, although the bull's eye method used significantly less fluoroscopy time compared to the triangulation method.<sup>22</sup> These results are concordant with our findings.

Other studies have compared the two techniques in terms of adverse events, hospitalization time, and success (defined as stone-free rate) and shown no difference between the two methods in any of these domains.<sup>2,3</sup> The studies concluded that there is no preferred method to gaining PCA.

Our study shows similar results, with no difference in outcomes between the two methods; however, we were able to demonstrate that the bull's eye technique, in the hand of novices, is associated with a faster technical skills gain. Given the steep learning curve of PCNL skill acquisition<sup>9</sup> and limited residency training, being able to identify a technique that is faster to master while still providing the same outcomes is essential for skill acquisition.

**Limitations**

Our study has some limitations. We had a small sample size of 15 medical trainees from one institution; however, due to the crossover study design, each trainee was their own control, thus minimizing confounding factors. As shown, there was no difference between performance scores and group allocation, reinforcing the idea that the crossover design was able to avoid bias and wash out the experience gained from prior attempts at access. As well, all trainees had no experience performing any steps of PCNL and had never used the PERC Mentor simulator, therefore avoiding any prior knowledge bias.

We were, unfortunately, unable to report whether the trainees accessed the written instructional material and videos prior; however, the teaching was re-explained and demonstrated in person, where all trainees were in attendance.

Other limitations included having only one resident observe the trainees and give the PCNL-GRS score.

Therefore, we were unable to prove inter-observer reliability, although Matsumoto et al previously showed that the PCNL-GRS system has high inter-rater reliability, with two attendings and two fellows involved in their study.<sup>13</sup> Alternatively, having only one evaluator allowed consistency in the scoring metric. As well, the use of both subjective and objective performance evaluation methods allowed for a more global approach of each trainee's true competency level. This study did not look at a specific renal calyx of interest, which could be explored in future studies.

## CONCLUSIONS

Our study shows that both methods of obtaining PCA were well acquired by trainees; however, in the hand of novices, the bull's eye method of gaining PCA was associated with a shorter operative time, as well as shorter fluoroscopy exposure time by almost 30% compared to the triangulation method. The rate of adverse events for both methods were similar as well. Either method can be taught safely to novice trainees to acquire the necessary skills to perform PCNL without increased complication rates.

COMPETING INTERESTS: The authors do not report any competing personal or financial interests related to this work.

This paper has been peer-reviewed.

## REFERENCES

- Ng CF. Training in percutaneous nephrolithotomy: The learning curve and options. *Arab J Urol* 2014;12:54-7. <https://doi.org/10.1016/j.aju.2013.08.002>
- Tepeler AA, Armağan T, Akman et al. Impact of percutaneous renal access technique on outcomes of percutaneous nephrolithotomy. *J Endourol* 2012;26:828-33. <https://doi.org/10.1089/end.2011.0563>
- Budak S, Yucel C, Kisa E, et al. Comparison of two different renal access techniques in one-stage percutaneous nephrolithotomy: Triangulation vs. eye of the needle. *Ann Saudi Med* 2018;38:189-93. <https://doi.org/10.5144/0256-4947.2018.189>
- Stern J, Zeltser IS, Pearle MS. Percutaneous renal access simulators. *J Endourol* 2007;21:270-3. <https://doi.org/10.1089/end.2007.9981>
- Watterson JD, Soon S, Jana K. Access related complications during percutaneous nephrolithotomy: Urology vs. radiology at a single academic institution. *J Urol* 2006;176:142-5. [https://doi.org/10.1016/S0022-5347\(06\)00489-7](https://doi.org/10.1016/S0022-5347(06)00489-7)
- Kim SC, Kuo RL, Lingeman JE. Percutaneous nephrolithotomy: An update. *Curr Opin Urol* 2003;13:235-41. <https://doi.org/10.1097/00042307-200305000-00012>
- Gadzhiev N, Brovkin S, Grigoryev V, et al. Sculpturing in urology, or how to make percutaneous nephrolithotomy easier. *J Endourol* 2015;29:512-7. <https://doi.org/10.1089/end.2014.0656>
- Li H, Chen Y, Liu C, et al. Construction of a three-dimensional model of renal stones: Comprehensive planning for percutaneous nephrolithotomy and assistance in surgery. *World J Urol* 2013;31:1587-92. <https://doi.org/10.1007/s00345-012-0998-7>
- Sharma GR, Luitel B. Techniques for fluoroscopy-guided percutaneous renal access: An analytical review. *Indian J Urol* 2019;35:259-66. [https://doi.org/10.4103/iju.IJU\\_149\\_19](https://doi.org/10.4103/iju.IJU_149_19)
- Speed JM, Wang Y, Leow JJ, et al. The effect of physician specialty obtaining access for percutaneous nephrolithotomy on perioperative costs and outcomes. *J Endourol* 2017;31:1152-6. <https://doi.org/10.1089/end.2017.0441>
- Saluk J, Ebel J, Rose J, et al. Fellowship training in endourology: Impact on percutaneous nephrolithotomy access patterns. *Can Urol Assoc J* 2022;16:E76-81. <https://doi.org/10.5489/cuaj.7339>
- Quirke K, Aydin A, Brunckhorst O, et al. Learning curves in urolithiasis surgery: A systematic review. *J Endourol* 2018;32:1008-20. <https://doi.org/10.1089/end.2018.0425>
- Matsumoto ED, Hamstra SJ, Radomski SB, et al. A novel approach to endourological training: training at the Surgical Skills Center. *J Urol* 2001;166:1261-6. [https://doi.org/10.1016/S0022-5347\(05\)65749-7](https://doi.org/10.1016/S0022-5347(05)65749-7)
- Vijayakumar M, Balaji S, Singh A, et al. A novel biological model for training in percutaneous renal access. *Arab J Urol* 2019;17:292-7. <https://doi.org/10.1080/2090598X.2019.1642600>
- Karagozlu Akgul A, Unal D, Demirbas M, et al. A simple, non-biological model for percutaneous renal access training. *Urol J* 2018;15:1-5.
- Noureldin YA, Andonian S. Simulation for percutaneous renal access: Where are we? *J Endourol* 2017;31:510-9. <https://doi.org/10.1089/end.2016.0587>
- Papatsoris AG, Shaikh T, Patel D, et al. Use of a virtual reality simulator to improve percutaneous renal access skills: A prospective study in urology trainees. *Urol Int* 2012;89:185-90. <https://doi.org/10.1159/000337530>
- Zhang Y, Yu CF, Liu JS, et al. Training for percutaneous renal access on a virtual reality simulator. *Chin Med J (Engl)* 2013;126:1528-31.
- Noureldin YA, Fahmy N, Anidjar M, et al. Is there a place for virtual reality simulators in assessment of competency in percutaneous renal access? *World J Urol* 2016;34:733-9. <https://doi.org/10.1007/s00345-015-1652-y>
- Knudsen BE, Matsumoto ED, Chew BH, et al. A randomized, controlled, prospective study validating the acquisition of percutaneous renal collecting system access skills using a computer based hybrid virtual reality surgical simulator: Phase 1. *J Urol* 2006. 176:2173-8. <https://doi.org/10.1016/j.juro.2006.07.011>
- Urology Specialty Committee. EPA Guide: Urology. 2017. Available online at: <https://www.royalcollege.ca/rcsite/documents/cbd/epa-guide-urology-e.pdf>. Accessed February 2, 2023.
- Abdallah MM, Salem SM, Badreldin MR, et al. The use of a biological model for comparing two techniques of fluoroscopy-guided percutaneous puncture: A randomized crossover study. *Arab J Urol* 2013;11:79-84. <https://doi.org/10.1016/j.aju.2012.12.001>

CORRESPONDENCE: Dr. Nader Fahmy, Division of Urology, Department of Surgery, McGill University, Montreal, QC, Canada; [fahmynader@yahoo.com](mailto:fahmynader@yahoo.com)