

# Urinary complications after penile inversion vaginoplasty in transgender women

## Systematic review and meta-analysis

Christina Ding<sup>1</sup>, Adree Khondker<sup>1</sup>, Mitchell G. Goldenberg<sup>2</sup>, Jethro C.C. Kwong<sup>2</sup>, Katherine Lajkosz<sup>2</sup>, Emery Potter<sup>3</sup>, Alexandra Millman<sup>3,4</sup>, Yonah Krakowsky<sup>3,4\*</sup>, Nathan Perlis<sup>2\*</sup>

<sup>1</sup>Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada; <sup>2</sup>Division of Urology, Department of Surgery, University of Toronto, Toronto, ON, Canada; <sup>3</sup>Transition-Related Surgery Program, Women's College Hospital, University of Toronto, Toronto, ON, Canada; <sup>4</sup>Department of Surgery, Division of Urology, Women's College Hospital, University of Toronto, Toronto, ON, Canada

\*Co-senior authors

Cite as: Ding C, Khondker A, Goldenberg MG, et al. Urinary complications after penile inversion vaginoplasty in transgender women: Systematic review and meta-analysis. *Can Urol Assoc J* 2023;17(4):121-8. <http://dx.doi.org/10.5489/cuaj.8108>

Published online December 6, 2022

Appendix available at [cuaj.ca](http://cuaj.ca)

## ABSTRACT

**INTRODUCTION:** Penile inversion vaginoplasty (PIV) remains the gold standard technique for vaginoplasty, a gender-affirming feminizing surgery, but has been associated with urinary complications; however, there is little literature synthesizing urinary complications after PIV surgery, and there is a need to compile these complications to counsel patients pre- and postoperatively on managing surgical expectations. In this systematic review, we summarize the prevalence of urinary complications following PIV.

**METHODS:** We searched the MEDLINE, EMBASE, CINAHL, and Scopus databases in July 2020. The primary outcome was the prevalence of urinary and surgical complications in patients after penile inversion vaginoplasty. Pooled prevalence was determined from extrapolated data. ROBINS-I tool was used to assess study quality. The study was prospectively registered on PROSPERO (CRD42020204139).

**RESULTS:** Of 843 unique records, 27 articles were pooled for synthesis, with 3388 patients in total. Overall patient satisfaction ranged from 80–100%. The most common urological complications included poor/splayed stream (11.7%, 95% confidence interval [CI] 5.7–19.3), meatal stenosis (6.9%, 95% CI 2.7–12.7), and irritative symptoms (frequency, urgency, nocturia) (11.5%, 95% CI 2.6–25.1). Other urinary complications included retention requiring catheterization (5.1%, 95% CI 0.3–13.8), incontinence (8.7%, 95% CI 3.4–15.6), urethral stricture (4.6%, 95% CI 1.2–9.8), and urinary tract infection (5.6%, 95% CI 2.7–9.4). Most pooled studies had moderate risk of bias.

**CONCLUSIONS:** The available evidence suggests that there is a low prevalence of urinary complications following PIV. Overall, there is a need for standardization of data in transgender surgical care to better understand surgical outcomes and improve postoperative management.

## INTRODUCTION

Penile inversion vaginoplasty (PIV) is the gold standard and most common approach used among a spectrum of transition-related genital feminizing surgeries.<sup>1,2</sup> In PIV, the penis is disassembled and the corporal bodies are resected. A neo-clitoris is created from a portion of the glans and preserved neurovascular bundle; a vulva is created from penile and perineal skin; and the neovagina is created by dissecting into the perineal space, which is then lined with penile and scrotal skin. The native urethra is shortened, spatulated, and fixed to an anatomic position at the base of the vestibule between the clitoris and the vaginal introitus.<sup>3</sup> A urethral catheter is typically left in place for 3–7 days until the vaginal packing is removed; a suprapubic catheter may also be placed. Studies have demonstrated high satisfaction rates following PIV.<sup>4,5</sup>

The prevalence of urological postoperative complications in PIV are low.<sup>6,7</sup> Short-term urinary complications include urinary retention and urinary tract infections (UTI).<sup>7,8</sup> Patients may also experience complications such as meatal stenosis, urinary stream spraying, and recurrent UTIs. There are several reasons why the rates of urinary complications are largely unknown. Patients often travel great distances to specialty centers for surgery and are, therefore, unreachable for followup care.<sup>9</sup> Furthermore, some patients do not seek help for fear of experiencing transphobia when accessing care, and it is believed that many patients live with complications without seeking

help.<sup>10,11</sup> As transition-related surgeries become increasingly common, it is of timely importance for the general urological community to recognize and be aware of the common urological complications so that they can manage them appropriately. While previous reviews have examined outcomes following vaginoplasty, large-scale meta-analyses focusing on urinary complications are lacking.<sup>7</sup> Past systematic reviews often excluded clinically relevant outcomes, such as retention, splayed stream, incontinence, need for revision surgery, and patient satisfaction.<sup>12-14</sup> Herein, we present the largest systematic review and meta-analysis to date focusing on the prevalence of urinary complications after PIV.

## METHODS

### Search strategy

The following databases were searched from database inception through July 2020: MEDLINE, EMBASE, CINAHL, and Scopus. Search strategy included MESH terms and keywords, such as “sex reassignment surgery,” “vaginoplasty,” “gender-affirming surgery,” and “urinary complications,” as detailed in Table 1. The references of published studies and grey literature were manually searched to ensure articles were not missed. This systematic review and meta-analysis was reported in accordance with the Preferred Reporting items for Systematic Reviews and Meta-Analyses (PRISMA).<sup>15</sup>

### Eligibility criteria and data collection

Observational studies with data on urinary complications following PIV were included. Exclusion criteria included narrative reviews, case reports/series including less than five patients, trauma-indicated or congenital surgery, studies not involving urinary complications, and supplement articles. Articles containing alternate vaginoplasty surgery techniques, such as those involving bowel flaps or peritoneum, were excluded, as PIV involving penile and scrotal skin is more commonly performed in gender-affirming surgery and these alternative techniques have unique complications. References from prior systematic reviews were included in the title and abstract screening.

Titles and abstracts were screened independently by two authors for full-text review. Any disagreements were resolved by a third author. The following data were included: number of patients, age at surgery, length of followup, and patient satisfaction. Urological complications captured in our review included: UTI, urethral stricture, meatal stenosis, urethral injury and other surgical complications, urinary retention, incontinence, poor or splayed stream, irritative symptoms,

**Table 1. Search strategy from Embase Classic + Embase 1947 to July 17, 2020**

Search line	Search terms
1	exp Sex reassignment procedures/
2	Transsexualism/su
3	(Gender affirm* adj3 surger*).tw,kw.
4	((transit* or gender*) adj3 (surg* or procedure* or operation*)).ti,kw.
5	Sex change*.tw,kw.
6	Gender reassignment*.tw,kw.
7	Transsex*.tw,kw.
8	Gender change*.tw,kw.
9	Transgender*.tw,kw.
10	Transvestism*.tw,kw.
11	Gender identity disorder*.tw,kw.
12	Genitoplast*.tw,kw.
13	Vaginoplast*.tw,kw.
14	Neovagina*.tw,kw.
15	Vagina/su
16	(Penile inversion* adj3 vaginoplast*).tw,kw.
17	Scrotal inversion vaginoplasty.tw,kw.
18	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12
19	13 or 14 or 15
20	18 and 19
21	16 or 17 or 20

and revision surgeries for urinary complications. In the included studies, urological complications were either patient-reported outcomes or surgeon reports of outcomes (see Supplementary Table 1 for PICOTS criteria; available in the Appendix at [cuaj.ca](#)).

### Statistical analysis

DerSimonian-Laird random effects models were used to pool the Freeman-Tukey double-arcsine-transformed urinary complication percentages for each complication. The 95% confidence intervals (CI) were calculated using Wilson’s method. Summary statistics for the pooled cohort were estimated using weighted sampling based on the cohort size for each study. All statistical analyses were conducted using R version 4.1.1.

### Risk of bias

In accordance with Cochrane recommendations,<sup>16</sup> risk of bias for individual non-randomized studies was

appraised using all domains of the ROBINS-I tool.<sup>17</sup> Two authors independently assessed each study for risk of bias and any discrepancies were resolved by consensus. Studies with critical risk of bias were specified to be removed from quantitative synthesis. Funnel plots were generated to examine the risk of publication bias for all meta-analyses.

## RESULTS

### Search results and study characteristics

The PRISMA flow diagram for study selection is shown in Figure 1. A total of 843 unique records were screened after 776 duplicates were removed. After title and abstract screening, 133 articles moved onto full-text screening. The proportionate agreement was 86% in title and abstract screening. A third reviewer resolved any differences in title and abstract screening. Of these, 27 articles were pooled for quantitative synthesis comprising of 3388 patients.

### Risk of bias

The risk of bias for each study based on the ROBINS-I tool is provided in Supplementary Table 2 (available in the Appendix at *cuaj.ca*). Of the 27 articles, two were low risk, 21 were moderate risk, and four were severe risk of bias. The primary cause for at least moderate risk of bias was bias due to confounding factors and bias in measurement of outcomes.

### Urinary complications

Mean age at the time of vaginoplasty was  $36.2 \pm 4.5$  years (range 24–44 years). Mean followup was  $2.6 \pm 2.7$  years (range 0.15–7.8 years). Overall patient satisfaction ranged from 80–100% (median satisfaction 96.1%). Urological complications following PIV are summarized in Table 2. The three most common urological complications were poor/splayed stream, meatal stenosis, and irritative symptoms (frequency, urgency, nocturia) (Figures 2A–H).

Of the studies that reported UTI in PIV, the rates ranged from 0–32%, with 32% being an outlier (studies  $n=10$ ). The remaining reported rates of UTI ranged from 0–15%. The median was 4.4%, and the pooled average of UTIs was 5.6% (95% CI 2.7–9.4%).

Of the studies that reported urethral strictures ( $n=11$ ), the complications rate ranged from 0–18.3%. The median was 7.1%, and the pooled average of urethral strictures was 4.6% (95% CI 1.2–9.8%).

Meatal stenosis occurred from 1.1–40% after PIV (studies  $n=17$ ), with 40% as an outlier; the remaining

reported rates of meatal stenosis occurred at a rate of 1.8–14.8%. The median rate of meatal stenosis was 4.4%, and the pooled average of meatal stenosis was 6.9% (95% CI 2.7–12.7%).


Of the studies that reported urinary retention requiring catheterization ( $n=5$ ), the complications rate ranged from 0–13.9%. The median rate was 6.8%, and the pooled average was 5.1% (95% CI 0.3–13.8%).

Urinary incontinence can occur in 1.1–27.3% of patients after PIV, with median reported incidence of 9.1%, and pooled average of 8.7% (95% CI 3.4–15.6%). Urinary incontinence was reported in eight studies.

Poor or splayed stream occurs in 1.8–33.2%, with 33.2% being an outlier. The remaining reported rates of poor or splayed stream ranged from 0–22.5% (studies  $n=10$ ). The median in the data set for poor or splayed stream was 10.2%, and the pooled average was 11.7% (95% CI 5.7–19.3%).

### Surgical complications

Surgical complications are summarized in Supplementary Table 3 (available in the Appendix at *cuaj.ca*). There was no standardized surgical complication grading system used in the publications. Of the reported surgical complications, the most common surgical complications were wound dehiscence ( $n=9$  studies, range 0.6–33%, median 6.7%), vaginal stenosis ( $n=6$  studies, range 2.1–18%, median 3.05%), urethral injury ( $n=15$  studies, range 0.6–10.9%, median 5%), and rectal injury ( $n=11$  studies, range 1.1–7.5%, median 2.3%). Of the studies that reported neovaginal fistulas ( $n=13$ ), urethrovaginal



**Figure 1.** PRISMA 2020 flow diagram for new systematic reviews which included searches of databases and other sources. Adapted from Page MJ, et al. *BMJ* 2021;372:n71. For more information, visit: <http://www.prisma-statement.org/>.

**Table 2. Study characteristics on studies including the prevalence of urinary complications after penile inversion vaginoplasty**

Study	Patient characteristics			Urinary complications							Patient satisfaction
	n	Mean patient age at surgery (range)	Mean followup length in years (range)	UTI	Urethral stricture	Meatal stenosis	Retention	Incontinence	Poor or splayed stream	Irritative symptoms (frequency, urgency, nocturia)	
Pooled Statistics					5.7% (0–18.3%)	10.8% (1.1–40%)	8.6% (0–13.9%)	6.6% (1.1–27.3%)	11.9% (1.8–33.2%)	8.8% (7.8%, 19.3%)	
Amend 2013 <sup>22</sup>	24	39.1 (20–54)	3.3 (1.6–5.8)	0%	–	4.2%	–	8.3%	–	–	100%
Blanchard 1987 <sup>23</sup>	22	32.3 (22.8–59.5)	4.4 (0.5–11.8)	–	–	–	–	27.3%	9.1%	–	90.4%
Bouman 1988 <sup>24</sup>	55	28* (10–60)	2.33 (0.16–6.25)	–	9.1%	1.8%	–	–	–	–	98.2%
Buncamper 2016 <sup>2</sup>	475	38.6 <sup>†</sup> (18.1–70.8)	7.8 <sup>†</sup> (1.0–15.9)	4.4%	–	9.7%	13.9%	–	9.5%	–	–
Cristofari 2019 <sup>25</sup>	189	36.8 (19.9–64.7)	1.4 (12–59)	6.3%	–	1.6%	–	1.1%	–	–	98.4%
Eldh 1993 <sup>26</sup>	20	34 (20–67)	5.8 (0.5–30.0)	–	15%	–	–	–	–	–	–
Falcone 2017 <sup>27</sup>	69	33.5 <sup>†</sup> (SD 10.2)	5 <sup>†</sup> (SD 3.2)	–	–	13.0%	–	–	–	–	–
Gaither 2018 <sup>11</sup>	330	35 <sup>†</sup> (18–76)	0.25 <sup>†</sup> (0.3–6.1)	1.5%	0%	–	–	–	1.8%	–	–
Goddard 2007 <sup>10</sup>	180	41 <sup>†</sup> (19–76)	0.15 <sup>†</sup> (0.02–0.96)	–	18.3%*	4.4%*	–	–	5.6%*	–	80%
Hoebcke 2005 <sup>28</sup>	31	–	–	32.3%	–	–	–	19.3%	–	19.3%	–
Huang 1995 <sup>29</sup>	109	32.3 (18–71)	–	–	–	7.4%*	–	–	–	–	–
Ives 2019 <sup>30</sup>	101	42 (18–71)	0.25 (0.25–0.50)	–	–	–	–	–	10.9%	–	–
Karim 1995 <sup>31</sup>	200	–	–	–	3.5%	–	–	–	–	–	–
Krege 2001 <sup>32</sup>	66	36.8 <sup>†</sup> (20–57)	–	–	–	10.6%	–	–	–	–	–
Lawrence 2006 <sup>6</sup>	232	44 (18–70)	3 (1–7)	–	–	3.9%	–	–	33.2%	–	–
Levy 2019 <sup>33</sup>	240	33 (SD 23)	0.24 (SD 0.37)	–	–	2.9%	–	–	–	–	–
Loree 2020 <sup>34</sup>	30	37 (SD 5.4)	1.27 (SD 9.9)	–	0%	–	0%	–	–	–	92.0%
Massie 2018 <sup>35</sup>	117	38 (16–78)	1.75 (1.1–2.8)	6.8%	0%	–	8.5%	15.4%	–	94%	–
Neto 2012 <sup>36</sup>	332	36.7 (19–68)	–	–	6.0%	40.0%	5.0%	4%	–	7.8%	–

\*The full study population was not assessed for the outcome, and the population event rate is assumed from the sample event rate. \*\*Estimated based on data given. <sup>†</sup>Median value provided if mean was unavailable. SD: standard deviation.

**Table 2 (cont'd). Study characteristics on studies including the prevalence of urinary complications after penile inversion vaginoplasty**

Study	Patient characteristics			Urinary complications							Patient satisfaction
	n	Mean patient age at surgery (range)	Mean followup length in years (range)	UTI	Urethral stricture	Meatal stenosis	Retention	Incontinence	Poor or splayed stream	Irritative symptoms (frequency, urgency, nocturia)	
Opsomer 2018 <sup>37</sup>	161	38.2 (18–69)	2.4 (0.9–3.9)	–	5.6%	–	–	–	–	–	–
Papadopoulos 2017 <sup>38</sup>	40	38.6 (26–50)	0.5	15.0%	–	–	–	10.0%	22.5%	–	–
Perovic 2000 <sup>39</sup>	89	28 (18–56)	3.6 (0.3–6.0)	–	–	1.1%	–	–	–	–	–
Raigosa 2015 <sup>40</sup>	60	28 (19–50)	2.0 (1.2–3.8)	–	8.3%	–	–	–	–	–	–
Revol 2006 <sup>41</sup>	63	33 (22–56)	1.3 (0.2–6.6)	–	–	7.9%	–	–	–	–	–
Stein 1990 <sup>42</sup>	14	31.4 (23–49)	1.8 (0.4–4)	–	7.1%	–	–	–	–	–	100%
Tavakkoli-Tabassi 2014 <sup>43</sup>	112	25.8 (SD 3.3)	1.1 (0.5–2.0)	3.6%	–	3.6%	–	–	–	–	85.7%
van Noort 1993 <sup>44</sup>	27	24 (21–57)	2.58** (0.1–8.6)	3.7%	–	14.8%	–	3.7%	7.4%	–	100%

\*The full study population was not assessed for the outcome, and the population event rate is assumed from the sample event rate. \*\*Estimated based on data given. †Median value provided if mean was unavailable. SD: standard deviation.

fistulas ranged from 1.5–4.5% (median 2.55%) and rectovaginal fistulas ranged from 0.3–3.6% (median 1.15%). Many studies did not have any fistulas in their sample. The most common urinary revision surgeries were meatoplasty (n=10 studies, 0.4–15%, median 5.3%) and urethral revision (n=5 studies, 1.5–27%, median 7.1%).

## DISCUSSION

There is limited high-quality data available on the surgical risks and urinary complications of vaginoplasty, which makes it difficult to counsel patients and educate physicians on the expected postoperative course. We performed the largest systematic review and meta-analysis to date examining urinary complications after PIV; 27 studies were examined and analyzed for urinary complications.

Our systematic review, which included all studies reporting on the prevalence of urinary complications, suggests that the overall risk of complications is low and of minor severity. These complications could be managed by a general urologist in most cases. Of 3388 patients, the mean prevalence of reported urinary complications ranged from 5.0–11.9% in non-randomized studies, with an overall intermediate risk of bias. The

most common urological complications included poor/splayed stream (11.7%, 95% CI 5.7–19.3%), meatal stenosis (6.9%, 95% CI 2.7–12.7%), and irritative symptoms (frequency, urgency, nocturia) (11.5%, 95% CI 2.6–25.1%). Other urinary complications included retention requiring catheterization (5.1%, 95% CI 0.3–13.8%), incontinence (8.7%, 95% CI 3.4–15.6%), urethral stricture (4.6%, 95% CI 1.2–9.8%), and UTI (5.6%, 95% CI 2.7–9.4%). Further, many of these complications, like UTI, splayed stream, and meatal stenosis, are of minimal morbidity and can be readily treated with antibiotics or meatal dilators.<sup>3</sup> Urethral dilation is a short-term solution for an urgent situation of acute retention, and most patients with non-resolving meatal stenosis can be treated definitively with meatoplasty (5.3%), and urethral revision surgery (7.1%). Most of the pooled studies had moderate risk of bias.

There are multiple explanations for why there is a relatively low complication rate for this gender-affirming surgery. Firstly, after PIV, the distance from the urethral meatus to the bladder is still longer than a cis-female urethra, which may be a preventative factor against UTIs.<sup>18</sup> Second, the urethra is spatulated widely during the surgery to decrease the rate of retention and




Figure 2 (A-D). Urinary complications of penile inversion vaginoplasty. (A) Urinary tract infections; (B) urethral stricture; (C) meatal stenosis; and (D) poor/splayed stream.

Figure 2 (E-H). Urinary complications of penile inversion vaginoplasty. (E) Incontinence; (F) genital pain; (G) irritation; and (H) retention.

stenosis. Furthermore, if strictures are present, they can be corrected at the time of gender-affirming surgery. Finally, the prostate decreases in size from feminizing hormone therapy, which can lower the rates of retention and bladder outlet obstruction.<sup>19</sup>

There may be under-reported urinary complication rates, especially in studies from the earlier decades. Patients often travel for surgery and long-term, consistent followup may be limited.<sup>9</sup> Additionally, complications may be treated at local institutions and, therefore, are not

captured in study followup. Furthermore, there are a small number of transgender patients who are post-radiation or who have bladder outlet obstruction secondary to benign prostatic hyperplasia (BPH). They will not be captured in this review but will be at higher risk of complications and would need general urological management.

In our review, we summarized 27 studies and 3383 patients after PIV, with robust data of urinary complications due to the large sample size. Moreover, we conducted the largest systematic review and meta-

analysis of comprehensive urinary complications following female transgender surgery. Overall, these findings support the notion that PIV is associated with low rates of urinary complications, and most patients are satisfied after surgery; however, our review suggests that patients should be advised that urinary complications can occur and may need conservative, medical, or surgical management. Ultimately, our findings provide useful data summaries for patients and healthcare practitioners to aid in treatment decisions and managing realistic surgical expectations. General urologists can treat and manage most of these complications, and they should do so in a trans-inclusive environment. This includes referring to the patient by their name, which may be different from the legal name, avoiding questions around transition not relevant to their care, and including resources in the clinic where trans people are represented.<sup>20</sup>

### Limitations

The findings in this study should be interpreted in the light of the following methodological limitations.

First, in the studies included in this meta-analysis, the primary outcome of urinary complications was not reported in a standardized method. For example, urinary complication rates were reported from research questionnaires and patient-reported outcomes, which can introduce a risk of reporting bias and prohibits analysis of symptom severity; however, surgical correction rates had direct clinical indications, which was more objective than reported urinary symptoms. To account for this, we abstracted the surgical correction rate whenever possible. Future work should, therefore, implement standardized definitions and methods of reporting surgical and urinary complication, which would allow a better understanding of outcomes after PIV and other gender-affirming surgeries.<sup>21</sup>

Second, urinary complications were pooled based on the reported complication rate in the articles, and zero-event studies were excluded if the outcome was not mentioned. Given a potential of a zero-complication rate for a specific outcome, the actual urinary complication rate may be lower than what is reported in this review.

Finally, the overall study quality was medium (most studies had moderate risk of bias and four studies had a severe risk of bias). For this reason, randomized, large, or prospective studies are required for certainty with regards to these outcomes. Further studies are also needed to examine how to best manage the urinary complications.

### CONCLUSIONS

PIV is the gold standard vaginoplasty technique, with excellent patient outcomes. We have presented the largest meta-analysis on the urinary complications of PIV and have shown that complication rates are low. Complications that do arise are well within a general urologist's skillset to manage; however, there remains a need for standardization of data collection in vaginoplasty surgery and by improving data, we will better understand the true prevalence of urinary complications. As such, urologists and other healthcare practitioners can use this systematic review to counsel patients pre- and postoperatively on managing surgical expectations.

COMPETING INTERESTS: The authors do not report any competing personal or financial interests related to this work.

This paper has been peer-reviewed.

### REFERENCES

- Selvaggi G, Bellringer. Gender reassignment surgery: An overview. *Nat Rev Urol* 2011;8:274-81. <https://doi.org/10.1038/nrurol.2011.46>
- Buncamper ME, van der Sluis WB, van der Pas RSD, et al. Surgical outcome after penile inversion vaginoplasty: A retrospective study of 475 transgender women. *Plast Reconstr Surg* 2016;138:999-1007. <https://doi.org/10.1097/PRS.0000000000002684>
- Buncamper ME, van der Sluis WB, de Vries M, et al. Penile inversion vaginoplasty with or without additional full-thickness skin graft: To graft or not to graft? *Plast Reconstr Surg* 2017;139:649e-56e. <https://doi.org/10.1097/PRS.00000000000003108>
- Boas SR, Ascha M, Morrison SD, et al. Outcomes and predictors of revision labiaplasty and clitoroplasty after gender-affirming genital surgery. *Plast Reconstr Surg* 2019;144:1451-61. <https://doi.org/10.1097/PRS.00000000000006282>
- De Vries ALC, McGuire JK, Steensma TD, et al. Young adult psychological outcome after puberty suppression and gender reassignment. *Pediatrics* 2014;134:696-704. <https://doi.org/10.1542/peds.2013-2958>
- Lawrence AA. Patient-reported complications and functional outcomes of male-to-female sex reassignment surgery. *Arch Sex Behav* 2006;35:717-27. <https://doi.org/10.1007/s10508-006-9104-9>
- Horbach SER, Bouman MB, Smit JM, et al. Outcome of vaginoplasty in male-to-female transgenders: A systematic review of surgical techniques. *J Sex Med* 2015;12:1499-1512. <https://doi.org/10.1111/jsm.12868>
- Nassiri N, Maas M, Basin M, et al. Urethral complications after gender reassignment surgery: A systematic review. *Int J Impot Res* 2020;33:793-800. <https://doi.org/10.1038/s41443-020-0304-y>
- Aizura AZ. Feminine transformations: Gender reassignment surgical tourism in Thailand. *Med Anthropol* 2010;29:424-43. <https://doi.org/10.1080/01459740.2010.501314>
- Goddard JC, Vickery RM, Qureshi A, et al. Feminizing genitoplasty in adult transsexuals: Early and long-term surgical results. *BJU Int* 2007;100:607-13. <https://doi.org/10.1111/j.1464-410X.2007.07017.x>
- Gaither TW, Awad MA, Osterberg, et al. Postoperative complications following primary penile inversion vaginoplasty among 330 male-to-female transgender patients. *J Urol* 2018;199:760-5. <https://doi.org/10.1016/j.juro.2017.10.013>
- Dreher PC, Edwards D, Hager, S et al. Complications of the neovagina in male-to-female transgender surgery: A systematic review and meta-analysis with discussion of management. *Clin Anat* 2018;31:191-9. <https://doi.org/10.1002/co.23001>
- Bustos SS, Bustos VP, Mascaro A, et al. Complications and patient-reported outcomes in transfemale vaginoplasty: An updated systematic review and meta-analysis. *Plast Reconstr Surg Glob Open* 2021;9:e3510. <https://doi.org/10.1097/GOX.00000000000003510>
- Manrique OJ, Adabi K, Martinez Jorge J, et al. Complications and patient-reported outcomes in male-to-female vaginoplasty-where we are today: A systematic review and meta-analysis. *Ann Plast Surg* 2018;80:684-91. <https://doi.org/10.1097/SAP.0000000000001393>

15. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Int J Surg* 2021;88:105906. <https://doi.org/10.1016/j.ijsu.2021.105906>
16. Higgins J, Thomas J, Chandler J, et al. Cochrane handbook for systematic reviews of interventions version 6.1 (updated September 2020). Cochrane, 2020. Available at: [Training.cochrane.org/handb](https://www.training.cochrane.org/handb). Accessed December 6, 2022.
17. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355. <https://doi.org/10.1136/bmj.i4919>
18. Abelson B, Sun D, Que L, et al. Sex differences in lower urinary tract biology and physiology. *Biol Sex Differ* 2018;9:1-13. <https://doi.org/10.1186/s13293-018-0204-8>
19. Jin B, Turner L, Walters WAW, Handelsman DJ. Androgen or estrogen effects on human prostate. *J Clin Endocrinol Metab* 1997;82:413. <https://doi.org/10.1210/jc.81.12.4290>
20. Anderson K, Krakowsky Y, Potter E, et al. Adult transgender care: A review for urologists. *Can Urol Assoc J* 2021;15:345. <https://doi.org/10.5489/cuaj.6949>
21. Cacci A, Frediani D, Cacciamani GE, et al. Systematic review of studies reporting perioperative and functional outcomes following male-to-female gender assignment surgery (MtoF GAS): A call for standardization in data reporting. *Minerva Urol e Nefrol Ital J Urol Nephrol* 2019;71:479-86. <https://doi.org/10.23736/S0393-2249.19.03407-6>
22. Amend B, Seibold J, Toomey P, et al. Surgical reconstruction for male-to-female sex reassignment. *Eur Urol* 2013;64:141-9. <https://doi.org/10.1016/j.eururo.2012.12.030>
23. Blanchard R, Legault S, Lindsay WRN. Vaginoplasty outcome in male-to-female transsexuals. *J Sex Marital Ther* 1987;13:265-75. <https://doi.org/10.1080/00926238708403899>
24. Bouman FG. Sex reassignment surgery in male to female transsexuals. *Ann Plast Surg* 1988;21:526-31. <https://doi.org/10.1097/0000637-198812000-00006>
25. Cristofari S, Bertrand B, Leuzzi S, et al. Postoperative complications of male to female sex reassignment surgery: A 10-year French retrospective study. *Ann Chir Plast Esthet* 2019;64:24-32. <https://doi.org/10.1016/j.anplas.2018.08.002>
26. Eldh J. Construction of a neovagina with preservation of the glans penis as a clitoris in male transsexuals. *Plast Reconstr Surg* 1993;91:893-5. <https://doi.org/10.1097/00006534-199304001-00023>
27. Falcone M, Timpano M, Ceruti C, et al. A Single-center analysis on the learning curve of male-to-female penoscrotal vaginoplasty by multiple surgical measures. *Urology* 2017;99:234-9. <https://doi.org/10.1016/j.urol.2016.07.012>
28. Hoebeke P, Selvaggi G, Ceulemans P, et al. Impact of sex reassignment surgery on lower urinary tract function. *Eur Urol* 2005;47:398-402. <https://doi.org/10.1016/j.eururo.2004.10.008>
29. Huang TT. Twenty years of experience in managing gender dysphoric patients: I. Surgical management of male transsexuals. *Plast Reconstr Surg* 1995;96:921-30. <https://doi.org/10.1097/00006534-199509001-00023>
30. Ives GC, Fein LA, Finch L, et al. Evaluation of BMI as a risk factor for complications following gender-affirming penile inversion vaginoplasty. *Plast Reconstr Surg - Glob Open* 2019;7:1-8. <https://doi.org/10.1097/GOX.0000000000002097>
31. Karim RB, Hoge JJ, Bouman FG, et al. Refinements of pre-, intra-, and postoperative care to prevent complications of vaginoplasty in male transsexuals. *Ann Plast Surg* 1995;35:279-84. <https://doi.org/10.1097/0000637-199509000-00010>
32. Kreye S, Bex A, Lümmen G, et al. Male-to-female transsexualism: A technique, results and long-term follow-up in 66 patients. *BJU Int* 2001;88:396-402. <https://doi.org/10.1046/j.1464-410X.2001.02323.x>
33. Levy JA, Edwards DC, Cutruzzola-Dreher P, et al. Male-to-female gender reassignment surgery: An institutional analysis of outcomes, short-term complications, and risk factors for 240 patients undergoing penile-inversion vaginoplasty. *Urology* 2019;131:228-233. <https://doi.org/10.1016/j.urol.2019.03.043>
34. Loree JT, Burke MS, Rippe B, et al. Transfeminine gender confirmation surgery with penile inversion vaginoplasty: An initial experience. *Plast Reconstr Surg Glob Open* 2020;8:e2873. <https://doi.org/10.1097/GOX.0000000000002873>
35. Massie JP, Morrison SD, Van Maasdam J, et al. Predictors of patient satisfaction and postoperative complications in penile inversion vaginoplasty. *Plast Reconstr Surg* 2018;141:911e-21e. <https://doi.org/10.1097/PRS.0000000000004427>
36. Neto RR, Hintz F, Kreye S, et al. Gender reassignment surgery: A 13-year review of surgical outcomes. *Int Braz J Urol* 2012;38:97-107. <https://doi.org/10.1590/S1677-55382012000100014>
37. Opsomer D, Gast KM, Ramaut L, et al. Creation of clitoral hood and labia minora in penile inversion vaginoplasty in circumcised and uncircumcised transwomen. *Plast Reconstr Surg* 2018;142:729E-33E. <https://doi.org/10.1097/PRS.0000000000004926>
38. Papadopulos NA, Zavlin D, Lellé JD, et al. Combined vaginoplasty technique for male-to-female sex reassignment surgery: Operative approach and outcomes. *J Plast Reconstr Aesthetic Surg* 2017;70:1483-92. <https://doi.org/10.1016/j.bjps.2017.05.040>
39. Perovic SV, Stanojevic DS, Djordjevic MLJ. Vaginoplasty in male transsexuals using penile skin and a urethral flap. *BJU Int* 2000;86:843-50. <https://doi.org/10.1046/j.1464-410x.2000.00934.x>
40. Raigosa M, Arvedimento S, Yoon TS, et al. Male-to-female genital reassignment surgery: A retrospective review of surgical technique and complications in 60 patients. *J Sex Med* 2015;12:1837-45. <https://doi.org/10.1111/jsm.12936>
41. Reval M, Servant JM, Banzet P. Surgical treatment of male-to-female transsexuals: A 10-year experience assessment. *Ann Chir Plast Esthet* 2006;51:499-511. <https://doi.org/10.1016/j.anplas.2006.02.006>
42. Stein M, Tiefer L, Melman A. Followup observations of operated male-to-female transsexuals. *J Urol* 1990;143:1188-92. [https://doi.org/10.1016/S0022-5347\(17\)40221-7](https://doi.org/10.1016/S0022-5347(17)40221-7)
43. Tavakkoli Tabassi K, Djavan B, Hosseini J, et al. Fold-back perineoscrotal flap plus penile inversion vaginoplasty for male-to-female gender reassignment surgery in circumcised subjects. *Eur J Plast Surg* 2014;38:43-8. <https://doi.org/10.1007/s00238-014-1038-1>
44. van Noord DE, Nicolai JP. Comparison of two methods of vagina construction in transsexuals. *Plast Reconstr Surg* 1993;91:1308-15. <https://doi.org/10.1097/00006534-199306000-00018>

CORRESPONDENCE: Dr. Yonah Krakowsky, Murray Koffler Urologic Centre, Toronto ON, Canada; [yonah.krakowsky@sinahealth.ca](mailto:yonah.krakowsky@sinahealth.ca)

Visit <https://www.cua.org/UROpedia> to complete the multiple-choice questionnaire associated with this article. This program is an Accredited Self-Assessment Program (Section 3) as defined by the Maintenance of Certification Program of The Royal College of Physicians & Surgeons of Canada, and approved by the Canadian Urological Association. You may claim a maximum of 1 hour of credit.