

The role of quantitative MRI-based prostate zonal parameters in predicting clinically significant prostate cancer: A U.S. cohort

Christina Sze¹, Aaron Brant¹, Jeffery P. Johnson¹, Zorawar Singh², Gina DeMeo³, Joseph Moryousef⁴, Kevin C. Zorn⁵, Naeem Bhojani⁵, Dean S. Elterman⁶, Daniel Margolis⁷, Bilal Chughtai¹

¹Department of Urology, Weill Cornell Medicine/New York Presbyterian Hospital, New York, NY, United States; ²Northwell Health, New York, NY, United States; ³Weill Cornell Medical College, New York, NY, United States; ⁴McGill University, Montreal, QC, Canada; ⁵Division of Urology, Centre Hospitalier de l'Université de Montréal, Montreal, QC, Canada; ⁶Division of Urology, University Health Network, University of Toronto, Toronto, ON, Canada; ⁷Department of Radiology, New York Presbyterian Hospital, New York, NY, United States

Cite as: Sze C, Brant A, Johnson JP, et al. The role of quantitative MRI-based prostate zonal parameters in predicting clinically significant prostate cancer: A U.S. cohort. *Can Urol Assoc J* 2023 April 11; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.8195>

Published online April 11, 2023

Corresponding author: Dr. Bilal Chughtai, Department of Urology, Weill Cornell Medicine/New York Presbyterian Hospital, New York, NY, United States; bic9008@med.cornell.edu

ABSTRACT

Introduction: We aimed to investigate the clinical utility of quantitative prostatic zonal measurements on in multiparametric magnetic resonance imaging (mpMRI) for the predication of clinically significant prostate cancer (csPCa).

Methods: A retrospective, single-institution study included 144 men who underwent mpMRI from 2015–2017. Prostate zone parameters were measured on mpMRI. Correlation and multivariable analysis evaluated relationship between prostate zone parameters and the presence of csPCa.

Results: The mean age was 66.9±7.8 years old. The median (interquartile range [IQR]) prostate volume and prostate-specific antigen (PSA) were 51.6 ml (37.1–74.5) and 6.1 ng/ml (4.5–8.2), respectively. Men with csPCa had significantly

KEY MESSAGES

- Our work demonstrates that transition zone parameters, such as thickness and volume, correlated with urological symptom scores and the presence of clinically significant cancer.
- Intravesical prostatic protrusion significantly predicts symptom scores.
- The use of these parameters had area under the receiver operating curve similar to PSAD.
- It is our hope that this work will encourage the application of prostate zone on MRI in the diagnostic workup of elevated PSA.

smaller total prostate volume (TPV), transitional zone volume (TZV), and transitional zone thickness (TZT), and larger transitional zone density (TZD) compared to those without PCa; however, on multivariate variable analysis, only TZD maintained significance. TZD had a comparable area under the curve to PSA density (PSAD) and PSA (0.74 vs. 0.73 vs. 0.60, respectively). Subgroup analysis of men with PCa, PSAD and TZD were significantly higher in men with Gleason grade group (GG) ≥ 2 compared to those with GG < 2 ($p=0.002$); however, this significance is not maintained on logistic regression in predicting GG.

Conclusions: Quantitative features of prostate zones on MRI may aid in identifying better predictors of csPCa. Zonal-based PSA density (TZD) may be a useful marker in identifying csPCa. Further exploration is needed to understand the clinical application of larger TZV in men with csPCa compared to those with insignificant disease.

INTRODUCTION

Prostate cancer (PCa) and benign prostatic hyperplasia (BPH) are common pathologies of the prostate in older men¹. Therefore, in patients who presents with a history of lower urinary tract symptoms (LUTs) without evidence of infection and an elevated level of prostate-specific antigen (PSA), effective management is aimed at distinguishing between the two conditions, however, confirmatory tissue samples are often needed^{2,3}. Unfortunately, post-biopsy complications do arise such as discomfort, transient LUTs, rectal bleeding, and infectious complications that can lead to sepsis⁴.

The prostate is separated into three unique zones: a central zone (CZ), transition zone (TZ), and a peripheral zone (PZ). The majority of PCa cases originate in the PZ of the prostate (70%), but PCa can also arise in the TZ (20%) and in the CZ (10%)⁵. It is well documented that TZ volume (TZV) correlates with age and PSA and contributes to the overall increase in gland size with minimal changes in PZ volumes (PZV). Additionally, PZV is poorly correlated with age^{6,7}. Large series of radical prostatectomy specimens demonstrated that tumors from larger prostates have more favorable pathological features including Gleason grade group (GG), smaller tumor volume, and lower risk of extraprostatic extension or seminal vesicle invasion, suggesting that a larger prostate may have a protective effect against PCa by exerting a large compressive mechanical stress on the PZ, therefore, impeding tumor growth and thinning out the PZ⁸⁻¹¹. However, this inverse relationship between BPH and PCa is under continuous debate due to contradictory studies and lack of thorough exploration³. Further consideration of prostate zones may help to define this relationship.

Technological advances in magnetic resonance imaging (MRI) have improved the ability to localize PCa and visualize the different prostate zones¹². The clinical application of assessing prostate zones on MRI and as non-invasive marker for patient selection needs further exploration

¹³. We aim to investigate the clinical application of quantitative prostatic zonal measurements on MRI to predict clinically significant prostate cancer (csPCa).

METHODS

This was a retrospective single-institution study of biopsy-naïve patients who underwent a mpMRI and transrectal ultrasound guided (TRUS) prostate biopsy (2-core targeted biopsy per lesion and 12-core systematic biopsy) between 2015 and 2017. All patients were assessed for inclusion consecutively. Patients on BPH medications were included in the study. Patients who reported 5-alpha reductase (5ARI) use, PSA was adjusted by multiplying the level by 2. Patients with positive urine cultures, symptoms consistent with acute or chronic prostatitis, history of BPH procedures or prostate cancer were excluded. This study was approved by the Weil Cornell Medicine Institutional Review Board (IRB No. 1601016896A004).

Prostate volume measurements

Due to difficulty in distinguishing borders of the respective zones on MRI and to remain consistent with previously published urological literature, the “transition” zone in this paper represents the combined central and transition zones of the prostate ¹⁴. Total prostate volume (TPV) and transition zone volume (TZV) was measured on sagittal and axial T2 weighted (T2W) MRI images using the prostate ellipsoid formula (volume= 0.52 x length x width x height) ¹⁴.

The following parameters were calculated from prostate MRI images: peripheral zone volume (PZV) = TPV-TZV; transition zone index (TZI) = TZV/TPV; transition zone density (TZD) = PSA/TZV. Thickness of the transition zone (TZT) and peripheral zone (PZT) was measured on axial T2W MRI images as the maximal straight anterior-posterior distance between the outer and inner margins of the TZ or PZ. An example of prostate volume and the aforementioned calculation is provided in the Supplements (Supplementary Figures 1.1, 1.2 and 1.3).

Statistical methods

Distributions of variables were compared by the chi-squared test for categorical variables and the Mann–Whitney U test for continuous variables, which was not normally distributed. A nonparametric spearman correlation was performed to examine correlations between zonal measurements on MRI as well as clinical outcomes such as presence of PCa and csPCa. csPCa is defined as Gleason score ≥ 7 or GG ≥ 2 . Fisher-Z-Transformation was used to test the significance of correlations between groups. Logistic regression analysis in univariate and multivariate models (nominal for PCa and csPCa) to determine predictive factors for symptom scores and presence of PCa. Linear regression analysis was used to find predictors of AUASS and QoL. Receiver operating characteristic (ROC) curves and area under ROC curve (AUC) were also computed to assess the diagnostic ability of MRI characteristics compared to prostate specific antigen density (PSAD).

All statistical tests were 2- sided using $p < 0.05$ indicating statistical significance. All statistics were performed using the statistical package JMP® (JMP Pro, Version 16 Software, Microsoft® Windows® for x 64; SAS Institute Inc., Cary, North Carolina, USA, 1989-2019).

RESULTS

A total of 144 men were included in this study. The baseline demographic and clinical characteristics are summarized in Table 1. The median (interquartile range [IQR]) age was 67 (61-73) years old. White patients accounted for 59% of the patient cohort, and African Americans accounted for 6.3%. Median body mass index (BMI) was 26.2 kg/m^2 (24.2-29.2).

Median (IQR) prostate volume and PSA were 51.6ml (37.1-74.5) and 6.1ng/ml (4.5-8.2), respectively. PCa was detected on biopsy in 68 (47%) patients and csPCA was present in 36 (25%) patients. The majority (52%) of patients had \geq PIRADS 4 lesions on mpMRI. On mpMRI, most lesions were in the PZ (65%), and the rest were in the TZ (10%) or bordering both zones (19%).

Men with csPCA were older ($p=0.001$) compared to those without csPCA (Table 2). TPV, TZV, and TZT were significantly larger in those without csPCA ($p=0.01$). However, PZV and PZT were not significantly different between the groups. Additionally, PSA did not differ however PSAD and TZD were significantly higher in the csPCA ($p < 0.001$). On multivariate regression, older age, higher BMI, and larger TZD were associated with the presence of csPCA ($p < 0.001$, $p=0.003$, $p=0.01$, respectively) (Figure 1(a)). PSAD was not associated with csPCA ($p=0.11$).

AUC for TZD (0.74) was comparable to that of PSAD (0.73) (Figure 1(b)). The highest Youden's index was at TZD of 0.28. At this point, the diagnosis of csPCA has a 64% sensitivity and 76% specificity compared to PSAD of 0.18 (at the highest Youden's index) which has a sensitivity of 53% and specificity of 86%. In the 107 patients who did not have csPCA on biopsy, using TZD as a predictor of csPCA, 76% (82/107) of those biopsies could've been avoided but it would've missed 38% (14/37) of csPCA confirmed on biopsy. Additionally, in the 69 patients with \leq PIRADS 3 lesions on MRI, a TZD cut-off of 0.28, would've avoided 55/69 (80%) of biopsies.

Subgroup analysis of men with PCa, PSAD and TZD was significantly higher in men with $GG \geq 2$ compared to those with $GG < 2$ ($p=0.002$). However, this significance is not maintained on logistic regression in predicting GG.

Repeat biopsy was available for 75 patients with a median follow of 15 months (9-33). Of the 75 patients, 30 patients were upstaged on their second biopsy. Of the 30 patients, 2 patients underwent robotic radical prostatectomies, 1 underwent radiotherapy and 1 underwent cryotherapy.

DISCUSSION

There is mixed data describing the relationships between enlarged prostates and PCa. A potential relationship becomes more apparent when examining prostate parameters at a zonal level. We

examine our cohort of patients who underwent a mpMRI and prostate biopsy (target and systematic) to understand the significance of prostate zone parameters in differentiate BPH and PCa. It has been proposed that a large TZ (from BPH) is protective against PCa by exerting compressive force on the PZ therefore thinning it out however this is not entirely supported by this study. Our data demonstrated that TPV, TZV, and TZT were significantly larger in those with cisPCa compared to men with csPCa ($p=0.01$). However, PZV and PZT were not significantly different between the groups. Additionally on spearman correlation, TZV and PZV were directly correlated ($\rho=0.56$, $p<0.001$).

Compared to other prostate zones, the TZ parameters are most predictive of the presence of csPCa. TZV was trending towards significance ($p=0.05$). On logistic regression, TZD significantly predicted csPCa (OR 19.8, 95%CI 4.95-34.68). However, neither PZD nor PZT significantly predicted csPCa. Finding alternative predictors of csPCa without the consideration of prostate size is important because estimation of prostate volume relies on formulas that do not represent the irregular shape of the prostate. In fact, inter-observer calculation of the prostate volume can fluctuate up to 20%¹⁵. Jiang *et al.* looked at 691 patients who underwent mpMRI prior to TRUS and found that the transverse TZ sectional area was predictive of PCa (AUC 0.81) therefore a better predictor than PSAD (AUC 0.71)¹⁶. Using a cross-sectional area instead of volume reduces the possibility of erroneous calculations by a multiplicative factor and may be the reason for the increase in AUC. It also worth noting that our study showed TPV was not correlated with age which is consistent previous longitudinal population study which demonstrated no statistically significant correlation between age and the rate of prostate volume on MRI in a cohort of 278 men¹⁷.

Our data demonstrated that PSAD was not a predictor of csPCa, however TZD was and the AUC of TZD was similar to PSAD, consistent with findings from previous studies. Tanaka *et al.* reported both TPV and PSAD levels had greater AUC values than PSA in the detection of PCa (0.71 and 0.71 vs 0.52, respectively). These AUCs are nearly identical to our study, further validating our methodology (0.71 and 0.70 vs 0.52, respectively)¹⁸. When looking at csPCa, the AUC for PSA and PSAD improved to 0.62 and 0.73, respectively. This is slightly lower than that found in a recent cohort of 900 men (AUC 0.64 PSA and AUC 0.78 PSAD)¹⁹. In our cohort, the AUC of TZD is similar to the AUC of PSAD for csPCa (0.74 vs 0.73, respectively). The validity of TZD in the detection of csPCa has been reported in numerous studies²⁰⁻²², and has been shown to have a stronger correlation than PSAD for cancer de-differentiation and aggressivity²³. PSAD has been shown to be a robust predictor for the detection of csPCa however our data did not demonstrate this significance¹⁹. However, the PSAD values were significantly different between the no cancer and cancer group (Table 1). Our calculated 53% sensitivity and 86% specificity for PSAD was much lower than that found by Yusim *et al.* (70% sensitivity and 79% specificity)¹⁹. Our highest Youden's index for PSAD was 0.18 ng/ml^2 compared to theirs of 0.20 ng/ml^2 . This difference could be due to inadequate sampling from our smaller sample size. We acknowledge further studies with larger cohorts are needed to find the optimal cut-off point

for TDZ, however it does not underscore that TZD is a useful predictive tool that can be utilized clinically. In our multivariable regression, TZD but not PSAD predicted csPCa. Similar studies have demonstrated that TZD may have stronger predictive power to discriminate PCa from BPH^{18, 23}. Taken together, our study highlights more specific prostatic parameters that may help guide clinical decision-making following imaging.

Our study has certain limitations. Our cohort consists of mild-moderate sized prostates and is predominantly Caucasian. Additionally, our sample size is small and may not be sufficient to truly assess the significance of TZD in aiding cancer detection when considering predicted lesions scored by PIRADS. Due to the nature of a retrospective study, it was difficult to assess baseline patient characteristics including medical history, duration of medication usage and social history at the time of the MRI and biopsy. In our cohort with relatively large median prostate volumes, alpha-blocker usage was only 30% which seems lower. The cohort for symptom-specific analysis was smaller due to the limitation in the availability of clinical data and variability in administering symptoms scores. Additionally, as with any retrospective study, the risk of selection bias in assessing the prostate zone measurements was present. To address this, measurements were averaged from three independent readers under the guidance of a senior radiologist. Lastly, the entire study was conducted with subjects from one institution, however, targeted biopsy was performed by three urologists with distinct practices that represent the variability of experience in the community. Albeit these limitations, our study does favor a clinically applicable difference in prostate zone parameters between BPH and PCa.

CONCLUSIONS

Quantitative features of prostate zones on MRI may aid in identifying better predictors of csPCa. Zonal-based PSA density (TZD) may be a useful marker in identifying csPCa. As the usage of multiparametric MRI increases in the realm of prostate cancer, further studies are required to understand the potential clinical applications of larger TZV for men with csPCa.

REFERENCES

1. Egan, K. B.: The Epidemiology of Benign Prostatic Hyperplasia Associated with Lower Urinary Tract Symptoms: Prevalence and Incident Rates. *Urol Clin North Am*, **43**: 289, 2016
2. Gades, N. M., Jacobson, D. J., Girman, C. J. et al.: Prevalence of conditions potentially associated with lower urinary tract symptoms in men. *BJU Int*, **95**: 549, 2005
3. Alcaraz, A., Hammerer, P., Tubaro, A. et al.: Is there evidence of a relationship between benign prostatic hyperplasia and prostate cancer? Findings of a literature review. *Eur Urol*, **55**: 864, 2009
4. Loeb, S., Vellekoop, A., Ahmed, H. U. et al.: Systematic review of complications of prostate biopsy. *Eur Urol*, **64**: 876, 2013
5. McNeal, J. E., Redwine, E. A., Freiha, F. S. et al.: Zonal distribution of prostatic adenocarcinoma. Correlation with histologic pattern and direction of spread. *Am J Surg Pathol*, **12**: 897, 1988
6. Bura, V., Caglic, I., Snoj, Z. et al.: MRI features of the normal prostatic peripheral zone: the relationship between age and signal heterogeneity on T2WI, DWI, and DCE sequences. *Eur Radiol*, **31**: 4908, 2021
7. Oreste, M., Giuseppe, P. G., Gianna, P. et al.: Between-subject variations of transition zone epithelial volume and serum PSA levels in men with benign prostatic hyperplasia. *World J Urol*, **28**: 379, 2010
8. Lorenzo, G., Hughes, T. J. R., Dominguez-Frojan, P. et al.: Computer simulations suggest that prostate enlargement due to benign prostatic hyperplasia mechanically impedes prostate cancer growth. *Proc Natl Acad Sci U S A*, **116**: 1152, 2019
9. Chen, M. E., Troncoso, P., Johnston, D. et al.: Prostate cancer detection: relationship to prostate size. *Urology*, **53**: 764, 1999
10. Hong, S. K., Poon, B. Y., Sjoberg, D. D. et al.: Prostate size and adverse pathologic features in men undergoing radical prostatectomy. *Urology*, **84**: 153, 2014
11. Newton, M. R., Phillips, S., Chang, S. S. et al.: Smaller prostate size predicts high grade prostate cancer at final pathology. *J Urol*, **184**: 930, 2010
12. Rosenkrantz, A. B., Verma, S., Turkbey, B.: Prostate cancer: top places where tumors hide on multiparametric MRI. *AJR Am J Roentgenol*, **204**: W449, 2015
13. Guneyli, S., Ward, E., Thomas, S. et al.: Magnetic resonance imaging of benign prostatic hyperplasia. *Diagn Interv Radiol*, **22**: 215, 2016
14. Tian, Y., Liu, H. M., Yang, B. et al.: Prostatic anatomical parameters correlate with clinical characteristics suggestive of benign prostatic hyperplasia. *Asian J Androl*, **23**: 64, 2021
15. Choi, Y. J., Kim, J. K., Kim, H. J. et al.: Interobserver variability of transrectal ultrasound for prostate volume measurement according to volume and observer experience. *AJR Am J Roentgenol*, **192**: 444, 2009
16. Jiang, S., Huang, Z., Liu, B. et al.: MRI-Based Nomogram of Prostate Maximum Sectional Area and Its Zone Area for Prediction of Prostate Cancer. *Front Oncol*, **11**: 708730, 2021

17. Loeb, S., Kettermann, A., Carter, H. B. et al.: Prostate volume changes over time: results from the Baltimore Longitudinal Study of Aging. *J Urol*, **182**: 1458, 2009
18. Tanaka, N., Fujimoto, K., Chihara, Y. et al.: Prostatic volume and volume-adjusted prostate-specific antigen as predictive parameters for prostate cancer patients with intermediate PSA levels. *Prostate Cancer Prostatic Dis*, **10**: 274, 2007
19. Yusim, I., Krenawi, M., Mazor, E. et al.: The use of prostate specific antigen density to predict clinically significant prostate cancer. *Sci Rep*, **10**: 20015, 2020
20. Kikuchi, E., Nakashima, J., Ishibashi, M. et al.: Prostate specific antigen adjusted for transition zone volume: the most powerful method for detecting prostate carcinoma. *Cancer*, **89**: 842, 2000
21. Kang, S. H., Bae, J. H., Park, H. S. et al.: Prostate-specific antigen adjusted for the transition zone volume as a second screening test: a prospective study of 248 cases. *Int J Urol*, **13**: 910, 2006
22. Djavan, B., Zlotta, A. R., Remzi, M. et al.: Total and transition zone prostate volume and age: how do they affect the utility of PSA-based diagnostic parameters for early prostate cancer detection? *Urology*, **54**: 846, 1999
23. Schneider, A. F., Stocker, D., Hotker, A. M. et al.: Comparison of PSA-density of the transition zone and whole gland for risk stratification of men with suspected prostate cancer: A retrospective MRI-cohort study. *Eur J Radiol*, **120**: 108660, 2019

Competing interests: *Dr. Chughtai is a consultant for Allergan and Boston Scientific. All other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.*

FIGURES AND TABLES

Figure 1. Prostatic parameters to predict clinically significant prostate cancer; (A) multivariate regression model, (B) receiver operating characteristic for transition zone density (red) and PSA density (blue).

A

Variables	Regression coefficient	95% CI		p value
Age	0.21	0.10	0.32	0.0002
BMI (kg/m ²)	0.23	0.08	0.38	0.003
IPP	0.25	-1.36	1.85	0.76
TPV	-0.38	-0.83	0.08	0.11
TZV	0.52	0.00	1.04	0.05
PZV	0.19	-0.14	0.52	0.27
TZD	19.81	4.95	34.68	0.01
TZI	-2.25	-12.84	8.34	0.68
TZT	-0.70	-3.03	1.83	0.56
PZT	-0.02	-3.92	3.88	0.99
TZT/PZT	-0.69	-1.58	0.19	0.13
PSAD	-20.78	-46.13	4.56	0.11
PSA	-0.12	-0.25	0.14	0.07

CI = confidence interval; BMI = Body mass index; IPP = Intra-vesical prostatic protrusion; PSAD = Prostate serum antigen density; TPV = Total prostate volume; TZV = Transition zone volume; PZV = Peripheral zone volume; TZD = PSA density of the transition zone; TZI = Transition zone index; TZT = Transition zone thickness; PZT = peripheral zone thickness; PSA =

B




Table 1. Demographic and clinical characteristics of all patients who underwent mpMRI and targeted biopsy	
Parameters	
No. of patients	144
Age (years)	
Mean (SD)	66.9 (7.8)
Median (IQR)	67 (61–72.8)
BMI (kg/m ²)	
Mean (SD)	27.1 (4.5)
Median (IQR)	26.2 (24.2–29.2)
Race	
White	85 (59.0%)
Black	9 (6.3%)
Asian	7 (4.9%)
Latino	6 (16.0%)
Other	23 (9.7%)
Not reported	14
Prostate volume (cc)	
Mean (SD)	60.4 (33.9)
Median (IQR)	51.6 (37.1–74.5)
PSA (ng/dL)	
Mean (SD)	10.5 (37.6)
Median (IQR)	6.1 (4.5–8.2)
IPSS score, median (IQR)	9 (6–18)
IPSS categories	
Mild	27 (40.91%)
Moderate	25 (37.88%)
Severe	14 (21.21%)
Qmax (cc/sec), median (IQR)	9.6 (6.6–12.4)
PVR (cc), median (IQR)	41 (9–84)
QoL score, Median (IQR)	2 (1–3)
MRI characteristics	
PI-RADS 2	19 (13.19%)
PI-RADS 3	50 (34.72%)
PI-RADS 4	58 (40.28%)
PI-RADS 5	17 (11.81%)
BPH severity	
Mild	38 (26.39%)
Moderate	80 (55.56%)
Severe	24 (16.67%)
Not reported	1 (0.69%)
IPP, median (IQR)	0.67 (0.40–0.95)

Biopsy pathology	
No cancer	76 (52.78%)
GG1	31 (21.53%)
GG2	29 (20.14%)
GG3	2 (1.39%)
GG4	2 (1.39%)
GG5	4 (2.78%)
α -adrenergic antagonist use	51 (35.4%)
5- α reductase inhibitor use	16 (11.1%)

BMI: body mass index; GG: Gleason grade group; IPP: intraprostatic protrusion; IPSS: International Prostate Symptom Score; IQR: interquartile range; mpMRI: multiparametric magnetic resonance imaging; PI-RADS: Prostate Imaging-Reporting and Data System; PSA: prostate-specific antigen; PVR: postvoid residual; Qmax: maximal urinary flow rate;

Parameters	No csPCa	csPCa	p
Patients, n (%)	107	37	
Age ^a , years	65 (60–72)	71 (64.5–77)	0.001
BMI ^a , kg·m ⁻²	25.8 (23.6–28.5)	26.4 (24.5–30.4)	0.06
PSA ^a , ng·ml ⁻¹	5.8 (4.5–8.1)	6.8 (4.8–10.7)	0.09
Race, n (%)			0.86
White	62 (63.9%)	23(69.7%)	
Black	6 (6.2%)	3 (9.1%)	
Asian	6 (6.2%)	1 (3.0%)	
Latino	5 (5.2%)	1 (3.0%)	
Other	18 (18.6%)	5 (15.2%)	
TPV ^a , ml	55.2 (41–78.7)	39.8 (27.4–63)	0.01
TZV ^a , ml	31.2 (22.9–46.4)	20.7 (13.2–38.3)	0.01
PZV ^a , ml	22.7 (16.5–32.7)	22 (13.8–27.9)	0.33
TZI ^a	0.58 (0.50–0.67)	0.55 (0.43–0.64)	0.07
TZT ^a , cm	3.21 (2.7–3.8)	2.7 (2.2–3.4)	<0.001
PZT ^a , cm	0.84 (0.69–1.11)	0.88 (0.73–1.25)	0.22
PSAD ^a , ng·ml ²	0.11 (0.08–0.15)	0.18 (0.11–0.31)	<0.001
TZD ^a , ng·ml ²	0.19 (0.12–0.28)	0.32 (0.18–0.64)	<0.001

^aContinuous variables are shown as the median value and interquartile range, Wilcoxon/Kruskal-Wallis Test. OR: odds ratio; PCa: prostate cancer; PV: prostate volume; PSA: prostate-specific antigen; PSAD: PSA density; PZT: peripheral zone thickness; PZV: peripheral zone volume; TPV: total prostate volume; TZD: transitional zone density; TZI: transitional zone index; TZT: transitional zone thickness; TZV: transitional zone volume.

	IPSS		QoL		AUASS		IIEF		PCa QoL	
	ρ	p	ρ	p	ρ	p	ρ	p	ρ	p
LL1	0.09	0.47	-0.06	0.61	0.16	0.22	-0.09	0.75	0.10	0.39
IPP	0.20	0.10	0.26	0.04	0.26	0.04	-0.17	0.57	0.16	0.16
TZI	0.08	0.53	0.17	0.18	0.00	0.98	0.33	0.25	0.07	0.52
TZ/PZ	0.07	0.58	0.13	0.29	0.00	0.99	0.33	0.25	0.06	0.61
PZT	0.10	0.42	0.04	0.75	0.04	0.76	-0.22	0.46	-0.08	0.48
TZT	0.31	0.01	0.35	0.00	0.30	0.02	0.12	0.67	0.13	0.24
PSAD	-0.06	0.65	-0.20	0.11	0.02	0.85	-0.16	0.58	-0.13	0.26
α -adrenergic antagonist use	0.16	0.21	0.25	0.05	0.11	0.42	0.16	0.21	0.25	0.03
5- α reductase inhibitor use	0.13	0.30	0.23	0.07	0.13	0.34	0.13	0.30	-0.01	0.95

AUASS American Urological Association symptoms score; IIEF International Index of Erectile Function; IPP: intraprostatic protrusion; IPSS: International Prostate Symptom Score; PCaQoL: expanded prostate cancer index composite for clinical practice and prostate cancer quality of life; PSAD: prostate-specific antigen density; PZ: peripheral zone; PZT: peripheral zone thickness; QoL: quality of life; TZ: transitional zone; TZI: transitional zone index; TZT: transitional zone thickness.