

Prevalence of secondary erythrocytosis in men receiving testosterone therapy

A matched-cohort analysis of intranasal gel, injections, and pellets

Rohit Reddy^{1*}, Parris Diaz^{1*}, Ruben Blachman-Braun¹, Justin Loloi³, Farah Rahman¹, Jesse Ory², Alexandra Dullea¹, Isaac Zucker¹, Daniel C. Gonzalez¹, Eliyahu Kresch¹, Ranjith Ramasamy¹

¹Desai Sethi Urology Institute, Miller School of Medicine, University of Miami, Miami, FL, United States; ²Department of Urology, Dalhousie University, Halifax, NS, Canada; ³Department of Urology, Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY, United States

*Equal contributors

Cite as: Reddy R, Diaz R, Blachman-Braun R, et al. Prevalence of secondary erythrocytosis in men receiving testosterone therapy: A matched cohort analysis of intranasal gel, injections, and pellets. *Can Urol Assoc J* 2023;17(7):E202-7. <http://dx.doi.org/10.5489/cuaj.8210>

Published online April 11, 2023

ABSTRACT

INTRODUCTION: Increased hematocrit (HCT) is a common adverse effect in men on testosterone therapy (TTh). We aimed to uncover differences in HCT changes among men receiving different forms of TTh.

METHODS: We conducted a single-center, retrospective, matched-cohort study of patients treated for testosterone deficiency (TD) to investigate the effect of three TTh regimens on HCT. We included men who received intranasal testosterone (NT), intramuscular testosterone (TC), or subcutaneous testosterone pellet (TP) regimens between January 2011 and December 2020. We matched treatment cohorts 1:1:1 for age, body mass index (BMI), and history of obstructive sleep apnea (OSA). Those taking TTh for <16 weeks were excluded. Comparison between groups was performed with Mann-Whitney U test, Student's t-test, ANOVA, or Kruskal-Wallis test as appropriate.

RESULTS: Seventy-eight matched-cohort individuals with TD received either NT, TC, or TP. The most common TD symptoms prior to initiation of TTh were erectile dysfunction (38%), low libido (22%), and lack of energy (17%). Baseline serum testosterone and HCT were higher in NT recipients ($p < 0.05$). As expected, all men receiving TTh were found to have increased serum testosterone levels at followup ($p < 0.001$). Relative to their respective baselines, men receiving TC experienced the greatest increase in serum testosterone (240.8 ng/dL to 585.5 ng/dL), followed by NT (230.3 ng/dL to 493.5 ng/dL) and TP (210.8 ng/dL to 360.5 ng/dL) (all $p < 0.001$). TC and TP were associated with significant increases in HCT (4.4% and 1.7%) while NT was associated with a decrease in HCT (-0.8%) at 16-week followup.

CONCLUSIONS: When controlled for age, BMI, and OSA, men receiving NT experienced decreased HCT compared to TC or TP at 16-week followup. Intranasal testosterone, while able to increase serum testosterone levels to reference range, does not appear to have a significant impact on HCT compared to the longer-acting forms of TTh.

INTRODUCTION

Testosterone therapy (TTh) has been regarded as the mainstay of treatment for symptomatic testosterone deficiency (TD) for the last 60 years.^{1,2} One of the adverse effects of TTh is increased hematocrit (HCT). Many forms of TTh can lead to increased HCT.^{3,4} The underlying mechanism of testosterone-induced rise in HCT is unclear, but it may involve increases in erythropoietin, decreases in hepcidin, or hypoxia from worsening obstructive sleep apnea (OSA). Newer theories have considered estradiol playing a causative role for increased HCT through the stimulation of hematopoietic stem cells.⁵ It is also unclear whether route of administration, the peak serum testosterone level, or the area under the therapeutic curve of men receiving TTh contributes to changes in HCT. Regardless of the underlying etiology, it appears that men who develop erythrocytosis (HCT >54%) within the first year of receiving TTh may have an increased risk of major adverse cardiovascular events.^{5,6}

To our knowledge, there are no direct studies evaluating differential changes in HCT between intranasal, intramuscular, and subcutaneous testosterone regimens in a matched cohort. We hypothesized that short-acting TTh would more closely resemble physiological endogenous testosterone release and would have less of an effect on HCT relative to longer-acting formulations. This study aimed to quantify the potential differences in HCT change

and erythrocytosis prevalence between intramuscular testosterone cypionate (TC), testosterone pellet (TP), and intranasal testosterone (NT). Ultimately, the results from this study may be used to better characterize the adverse effects of TTh regimens and to guide the decision-making process between patients and clinicians.

METHODS

We conducted a single-center, retrospective, matched-cohort analysis of patients treated for TD to investigate the effect of various TTh regimens on HCT. This study was reviewed and approved by our institutional review board.

Our health system database was searched for men receiving TTh between January 1, 2011, and December 31, 2020. Filtered patients completed a washout period of four weeks for gels and injection-based therapies and 16 weeks for subcutaneous pellets and reported rigorous adherence to their respective treatment schedule. We included men with a morning serum total testosterone (T) under 300 ng/dL; and hypogonadal symptoms, including erectile dysfunction (ED), sleep disturbances, decreased energy, low libido, premature ejaculation (PE), and depressed mood. Included men had a baseline HCT and were started on either TP, NT, or TC. Furthermore, we included men who stayed on TTh for 16 weeks or more and had subsequent bloodwork (testosterone and HCT, at a minimum) at that time point. We excluded men if they used more than one type of TTh during the targeted period.

A minimum of 16-week followup from treatment initiation was chosen, as one cycle of erythropoiesis is approximately 120 days.⁷ The dosages and frequency of administration were as follows: for TP, 800 mg every five months; for NT, 11 mg three times a day; and for TC, 100 mg every week.^{8,9}

We performed a retrospective chart review to characterize the distribution of presenting hypogonadal symptoms (listed above) of men diagnosed with TD (Figure 1). Among cohorts, we were able to match for age within three years, established history of OSA, and body mass index (BMI) within 4%.

Outcomes

We collected laboratory results, including HCT, serum total testosterone (T), estradiol (E), and prostate-specific antigen (PSA). The primary outcomes were changes in HCT and T following at least 16 weeks of TTh. We also recorded changes in E and PSA during the same 16-week period. In accordance with American Urological Association (AUA) guidelines, erythrocytosis was defined

as HCT >54%, and successful treatment of TD was defined as T level of ≥ 450 ng/dL.¹⁰ All T assays were collected using liquid chromatography-mass spectrometry.

Power calculation

The sample size of the study was calculated a priori based on previously published data for NT (+1% HCT; standard deviation [SD] 4%), as well as the Beggs et al study that demonstrated a 5% increase in HCT in men using weekly injections of TC.^{11,12} We set α to 0.05 and statistical power to 0.8. The two-tailed sample size calculated for an adequate power was 25 patients for each cohort.

Statistical analysis

We performed a matched-cohort study within the University of Miami database. The exposure was defined as use of TTh. Matched cohorts were constructed with patients placed in a 1:1:1 ratio for age, BMI, and OSA. Patients satisfying the inclusion and exclusion criteria were artificially compared in “triplets” for matching characteristics. Of all eligible patients, cohorts were constructed based on age, BMI, and OSA values or histories. Constraints for age was within three years, BMI within 4%, and prevalence of OSA was maintained in all three treatment cohorts.^{13,14}

Data analysis was performed with SPSS version 28. Continuous variables were presented as mean \pm SD or median (interquartile range [IQR] 25–75) in accordance with data distribution, and comparison between groups was performed with the Mann Whitney U-test, Student's t-test, ANOVA, or Kruskal-Wallis test as required. Differential prevalence of OSA among groups was analyzed with the Chi-squared test. Changes in T, E, and PSA over time within groups were evaluated with the Wilcoxon test. Absolute HCT change (from baseline to followup) was also reported. A p-value of <0.05 was considered statistically significant.

RESULTS

Seventy-eight men receiving either NT, TC, or TP (26 men in each treatment cohort) within the University of Miami Health System were “triplet” grouped into cohorts for appropriate matching of age, BMI, and OSA. The most common hypogonadal symptoms prior to initiation of TTh were ED (38%), low libido (22%), and lack of energy (17%) (Figure 1). Overall, mean age was 44.6 ± 10.6 years and mean BMI was 30.5 ± 5.8 kg/m², with a baseline T of 223.5 (191.1–260.4) ng/dL, PSA of 0.7 (0.4–1.15) ng/dL, and HCT of 44% (40.8–46.3). Age, T, E, and PSA at baseline were




Figure 1. Percent distribution of hypogonadal symptoms prior to initiation of testosterone therapy (TTh) in testosterone deficient (TD) men. *Estimates are based on retrospective medical chart review of patients. ED: erectile dysfunction; PE: premature ejaculation.

similar among groups ($p > 0.05$). At baseline, HCT and BMI were higher in men in the NT group ($p < 0.05$). All three TTh regimen groups had significantly different levels of T, HCT, and E at followup ($p < 0.05$) (Table 1).

On followup, there was no patient with erythrocytosis, that is, $HCT \geq 52\%$. The overall absolute median HCT change during the study period was 1.5 (-0.9–4.0) (1.5 ± 3.9). The median HCT change for the TP group was 1.7 (-0.05–3.7) (1.7 ± 3.2), for the NT group -0.8 (-2.3–1.1) (1.2 ± 2.8), and for the TC group 4.4 (1.6–6.9) (4.1 ± 3.8) (Figure 2A). Within each group, all TTh formulations successfully increased T levels to reference range ($p < 0.001$), with TP from 210.8 (145–234.4) ng/dL to 360.5 (250.8–486) ng/dL; NT from 230.3 (200.3–261.6) ng/dL to 493.5 (391–698.3) ng/dL; and TC from 240.8 (193.4–288.4) ng/dL to 584.5 (349.5–802.3) ng/dL (Figure 2B).

When considering the other serologic markers, there was no significant change in PSA (TP: $p = 0.147$; NT: $p = 0.057$; and TC: $p = 0.336$) and E (TP: $p = 0.909$; NT: $p = 0.915$; and TC: $p = 0.149$) during followup.

Overall, 12 (15.4%) men had OSA at baseline, and OSA prevalence was similar among the three groups ($p = 0.7440$). Baseline T, E, PSA, and HCT were similar among men with OSA and those without OSA ($p > 0.05$). On followup, patients with and without OSA had similar E, PSA, and HCT levels ($p > 0.05$); however, T levels were lower in patients with OSA (without

Table 1. Clinical and biochemical characteristics of the analyzed patients and comparison between TTh groups

	Overall N=78 (100%)	TP n=26 (33.3%)	NT n=26 (33.3%)	TC n=26 (33.3%)	p
Age (years)	44.6±10.6	43.7±7.4	43.7±10.2	46.5±13.5	0.550
BMI (kg/m ²)	30.5±5.8	29.5±7.2	32.7±4.7	29.1±4.5	0.042
OSA					
No (%)	66 (84.6%)	23 (88.5%)	22 (84.6%)	21 (80.8%)	0.744
Yes (%)	12 (15.4%)	3 (11.5%)	4 (15.4%)	5 (19.2%)	
Before treatment					
Testosterone (ng/dL)	223.5 (191.1–260.4)	210.8 (145–234.4)	230.3 (200.3–261.6)	240.8 (193.4–288.4)	0.090
PSA (ng/dL)	0.70 (0.40–1.15)	0.60 (0.40–0.82)	0.84 (0.40–1.53)	0.70 (0.50–1.30)	0.450
Hematocrit (%)	44 (40.8–46.3)	43.8 (40.7–46.9)	45.2 (44.3–46.7)	41.6 (38.9–44.5)	0.010
Estradiol (pg/mL)	23 (17.1–29)	22.2 (13.2–25.3)	22 (17.3–28.9)	23.1 (18.5–29.8)	0.522
After treatment					
Testosterone (ng/dL)	469 (327–662.5)	360.5 (250.8–486)	493.5 (391–698.3)	584.5 (349.5–802.3)	0.011
PSA (ng/dL)	0.70 (0.50–1.20)	0.65 (0.45–1.03)	1.0 (0.48–1.50)	0.68 (0.50–1.10)	0.402
Hematocrit (%)	45.5 (43.5–47.4)	45.5 (43.8–47.2)	44.3 (42.7–46.5)	47.1 (44.2–48.7)	0.033
Estradiol (pg/mL)	24.6 (18.1–34)	21.5 (9.1–25.7)	25.5 (18.3–33.8)	30 (19.3–54.4)	0.038

Mean ± standard deviation; median (interquartile range 25th–75th). BMI: body mass index; PSA: prostate-specific antigen; NT: intranasal testosterone; TC: testosterone cypionate; TP: subcutaneous testosterone pellets.




Figure 2. Box plot figure showing the changes within testosterone therapy (TTh) formulation groups of serum (A) change in hematocrit; and (B) testosterone levels.

OSA: 480 [342–675.3] ng/dL vs. with OSA: 338.5 [183–527] ng/dL, $p=0.048$) (Table 2).

DISCUSSION

Testosterone therapy has remained the mainstay of treatment for men with symptomatic TD; often, patients require lifelong supplementation to abate hypogonadal symptoms.¹⁵ Increased HCT is a common adverse effect of TTh, with the potential to cause serious adverse cardiovascular and thromboembolic effects.¹⁵ Previous studies have demonstrated that TTh formulations, such as NT, can closely mimic the physiological release of endogenous testosterone by virtue of short-acting properties and daily dosage requirements.^{16,17} Given previous studies demonstrating the capability of short-acting TTh to maintain sex hormones and sperm parameters, we hypothesized that NT would have less of an effect on HCT than longer-acting formulations.

To our knowledge, this is the first study to compare the changes in HCT between intranasal, intramuscular, and subcutaneous testosterone regimens in a matched cohort. We found that long-acting TTh is associated with significant increases in HCT compared to short-acting formulations. NT does not appear to have a significant impact on HCT compared to the longer-acting formulations while simultaneously increasing T levels to reference range in most patients.

The role of TTh in erythropoiesis has yet to be fully elucidated. Rather than having an indirect role in increasing erythropoietin, testosterone appears to act directly on hematopoietic stem cells to stimulate red blood cell synthesis.¹⁸ Several key mechanisms have been implicated, including iron incorporation into red

Table 2. Clinical and biochemical characteristics of the analyzed patients and comparison between men with OSA and those without OSA			
	No OSA n=66 (84.6%)	With OSA n=12 (15.4%)	p
Age (years)	44±10.4	48.1±11.6	0.221
BMI (Kg/m ²)	29.8±5.1	33.9±7.9	0.024
TRT			
TP (%)	23 (88.5%)	3 (11.5%)	
NT (%)	22 (84.6%)	4 (15.4%)	
TC (%)	21 (80.8%)	5 (19.2%)	0.744
Before treatment			
Testosterone (ng/dL)	226 (193.8–261.6)	208.9 (190–248.3)	0.346
PSA (ng/dL)	0.60 (0.40–1.20)	0.70 (0.48–1.15)	0.700
Hematocrit (%)	44 (40.8–46.3)	44.1 (39.7–47.7)	0.938
Estradiol (pg/mL)	22.5 (16.7–29)	23 (18–29)	0.747
After treatment			
Testosterone (ng/dL)	480 (342–675.3)	338.5 (183–527)	0.048
PSA (ng/dL)	0.70 (0.50–1.23)	0.73 (0.50–1.28)	0.883
Hematocrit (%)	45.4 (43.3–47.2)	46.9 (43.6–47.6)	0.432
Estradiol (pg/mL)	24.8 (19–33.2)	19.5 (15–36)	0.397

Mean ± standard deviation; median (interquartile range 25th–75th). BMI: body mass index; OSA: obstructive sleep apnea; PSA: prostate-specific antigen; NT: intranasal testosterone; TC: testosterone cypionate; TP: subcutaneous testosterone pellets.

cells.¹⁹ The effect of TTh on red blood cell production is substantiated by the development of anemia in men undergoing androgen deprivation therapy, which can

be corrected with androgen replacement or cessation of androgen deprivation therapy.^{20,21}

Studies directly comparing different TTh modalities with regards to HCT changes are lacking. From individual trials, rates of erythrocytosis have been reported as 1.3% in men using NT, 10.4% in men using TP, and 11.2% in men using TC.^{12,22,23} According to AUA guidelines, men on TTh who develop erythrocytosis may require dose adjustment, temporary discontinuation, or referral to a hematologist for phlebotomy. Unfortunately, while these interventions may help mitigate the adverse effects associated with erythrocytosis, patients may experience recurring hypogonadal symptoms for an indeterminate time. These implications highlight the importance of determining the “erythropoietic profiles” of different TTh delivery methods.

When examining the secondary outcomes of this study, it is important to note that aside from PSA, many differences were seen among the various TTh delivery methods. No TTh regimen significantly changed PSA at followup, reinforcing previous evidence that TTh likely has minimal effect on the prostate.^{21,22} Among the three cohorts, the TC group had the highest mean T levels (584.5 ng/dL), followed by the NT group (493.5 ng/dL) and the TP group (360.5 ng/dL). Our results indicate that many patients using TP were unable to reach a T level of 450 ng/dL, the benchmark of therapeutic success.²⁴ These results directly contrast with those of previous studies demonstrating that TP can sufficiently increase serum T to therapeutic levels.^{23,25}

Interestingly, despite patients with OSA having similar HCT as those without OSA on followup, the former was found to have significantly lower T levels, raising the question of whether patients with concomitant TD and OSA have a blunted response to TTh. Patients with OSA may inherently be subjected to low T levels, potentially due to OSA-induced decreased pituitary gonadal function.²⁶ Although OSA has been associated with polycythemia in the setting of TTh,²⁷ it remains to be seen whether OSA at varying severities modulates the response to TTh and if so, what the underlying mechanisms are.

In the present study, we were able to create matched cohorts and simultaneously control for age, BMI, and OSA in all three treatment groups. This improves on the existing literature, as most studies fail to adequately account for OSA, a known contributor to increased HCT.^{28,29} After matching, we observed an increased BMI among men in the NT arm. As lower HCT was seen in men of this treatment arm, we believe this value is not clinically significant despite statistical significance.

Secondly, recruitment took place in a diverse metropolitan area with a unique patient population, many of whom may be traditionally underrepresented in clinical research.³⁰ All serologic testing was done at outside institutions, such as LabCorp or Quest, that used liquid chromatography-mass spectrometry, the gold standard for T blood testing.

Limitations

Limitations of this study include unknown timing of blood tests relative to the TTh dosing schedule. This may have led to unreliable T levels due to the peaks and troughs associated with each TTh modality. Although patients switching TTh modality received routine wash-out periods, the degree of adherence is not directly observable. Additionally, as NT was not released until 2016, there is a time frame difference for recruitment, which could potentially affect the results. Our timeline to understand HCT changes was within 16 weeks; while this follows the laboratory evaluation guidelines under the American Society of Andrology, it prevents evaluation of HCT changes that may have occurred upon longer-term followup. Finally, some patients were previously enrolled in ongoing clinical trials and may have possibly benefited from additional visits, closer followup, and improved availability of medication, which may have led to enhanced medication adherence relative to the broader population. Further investigation of TTh-induced HCT changes should evaluate genetic and/or environmental dispositions to this phenomenon, how to properly trend individual patient changes in HCT, or even the long-term effect of NT on HCT in larger sample sizes.

CONCLUSIONS

Short-acting intranasal testosterone appears to adequately treat TD without significantly affecting HCT or risk of erythrocytosis. Men on longer-acting TTh modalities experienced an increase in HCT. To the best of our knowledge, our study is the first matched-cohort analysis to examine the differential HCT changes between intranasal, injectable, and subcutaneous pellet forms of TTh. Our findings add to the existing evidence on TTh adverse effect profiles and may subsequently influence the shared decision-making process for the optimal TTh regimen.

COMPETING INTERESTS: The authors do not report any competing personal or financial interests related to this work.

This paper has been peer-reviewed.

REFERENCES

- Jockenhovel F. Testosterone therapy — what, when, and to whom? *Aging Male* 2004;7:319-24. <https://doi.org/10.1080/13685530400016557>
- Nackeeran S, Kohn T, Gonzalez D, et al. The effect of route of testosterone on changes in hematocrit: A systematic review and Bayesian network meta-analysis of randomized trials. *J Urol* 2022;207:44-51. <https://doi.org/10.1097/JU.0000000000002188>
- Walker RF, Zakai NA, MacLehose RF, et al. Association of testosterone therapy with risk of venous thromboembolism among men with and without hypogonadism. *JAMA Intern Med* 2020;180:190-7. <https://doi.org/10.1001/jamainternmed.2019.5135>
- Nguyen CP, Hirsch MS, Moeny D, et al. Testosterone and “age-related hypogonadism” — FDA concerns. *N Engl J Med* 2015;373:689-91. <https://doi.org/10.1056/NEJMp1506632>
- Ory J, Nackeeran S, Balaji NC, et al. Secondary polycythemia in men receiving testosterone therapy increases risk of major adverse cardiovascular events and venous thromboembolism in the first year of therapy. *J Urol* 2022;101097JU0000000000002437. <https://doi.org/10.1097/JU.0000000000002437>
- White J, Petrella F, Ory J. Testosterone therapy and secondary erythrocytosis. *Int J Impot Res* 2022;34:693-7. <https://doi.org/10.1038/s41443-021-00509-5>
- Kautz L, Nemeth E. Molecular liaisons between erythropoiesis and iron metabolism. *Blood* 2014;124:479-82. <https://doi.org/10.1182/blood-2014-05-516252>
- Testosterone nasal gel (Natesto) for hypogonadism. *Med Lett Drugs Ther* 2015;57:73-4.
- McCullough A. A review of testosterone pellets in the treatment of hypogonadism. *Curr Sex Health Rep* 2014;6:265-9. <https://doi.org/10.1007/s11930-014-0033-7>
- Mullhall JP, Trost LW, Brannigan RE, et al. Evaluation and management of testosterone deficiency: AUA guideline. *J Urol* 2018;200:423-32. <https://doi.org/10.1016/j.juro.2018.03.115>
- Beggs LA, Yarrow JF, Conover CF, et al. Testosterone alters iron metabolism and stimulates red blood cell production independently of dihydrotestosterone. *Am J Physiol Endocrinol Metab* 2014;307:E456-61. <https://doi.org/10.1152/ajpendo.00184.2014>
- Rogol AD, Tkachenko N, Bryson N. NatestoTM, a novel testosterone nasal gel, normalizes androgen levels in hypogonadal men. *Andrology* 2016;4:46-54. <https://doi.org/10.1111/andr.12137>
- McClintock TR, Valovska M-TI, Kwon NK, et al. Testosterone replacement therapy is associated with an increased risk of urolithiasis. *World J Urol* 2019;37:2737-46. <https://doi.org/10.1007/s00345-019-02726-6>
- Katsoularis I, Fonseca-Rodríguez O, Farrington P, et al. Risk of acute myocardial infarction and ischemic stroke following COVID-19 in Sweden: A self-controlled case series and matched cohort study. *Lancet* 2021;398:599-607. [https://doi.org/10.1016/S0140-6736\(21\)00896-5](https://doi.org/10.1016/S0140-6736(21)00896-5)
- Kovac JR, Rajanahally S, Smith RP, et al. Patient satisfaction with testosterone replacement therapies: The reasons behind the choices. *J Sex Med* 2014;11:553-62. <https://doi.org/10.1111/jsm.12369>
- Ramasamy R, Masterson TA, Best JC, et al. Effect of Natesto on reproductive hormones, semen parameters, and hypogonadal symptoms: A single-center, open-label, single-arm trial. *J Urol* 2020;204:557-63. <https://doi.org/10.1097/JU.0000000000001078>
- Hellstrom WJG, Soubra A. Re: Effect of Natesto on reproductive hormones, semen parameters, and hypogonadal symptoms: A single-center, open-label, single-arm trial. *Eur Urol* 2021;79:890-1. <https://doi.org/10.1016/j.eururo.2021.02.012>
- Rochira V, Zirilli L, Madeo B, et al. Testosterone action on erythropoiesis does not require its aromatization to estrogen: Insights from the testosterone and estrogen treatment of two aromatase-deficient men. *J Steroid Biochem Mol Biol* 2009;113:189-94. <https://doi.org/10.1016/j.jsbmb.2008.12.007>
- Shahani S, Braga-Basaria M, Maggio M, et al. Androgens and erythropoiesis: Past and present. *J Endocrinol Invest* 2009;32:704-16. <https://doi.org/10.1007/BF03345745>
- Snyder PJ. Effects of testosterone replacement in hypogonadal men. *J Clin Endocrinol Metab* 2000;85:2670-7. <https://doi.org/10.1210/jc.85.8.2670>
- Bhasin S, Woodhouse L, Casaburi R, et al. Testosterone dose-response relationships in healthy young men. *Am J Physiol Endocrinol Metab* 2001;281:E1172-81. <https://doi.org/10.1152/ajpendo.2001.281.6.E1172>
- Rotker KL, Alavian M, Nelson B, et al. Association of subcutaneous testosterone pellet therapy with developing secondary polycythemia. *Asian J Androl* 2018;20:195-9. https://doi.org/10.4103/aja.aja_51_17
- Wheeler KM, Smith RP, Kumar RA, et al. A comparison of secondary polycythemia in hypogonadal men treated with domiphen citrate vs. testosterone replacement: A multi-institutional study. *J Urol* 2017;197:1127-31. <https://doi.org/10.1016/j.juro.2016.10.068>
- Fernández-Balsells MM, Murad MH, Lane M, et al. Clinical review 1: Adverse effects of testosterone therapy in adult men: A systematic review and meta-analysis. *J Clin Endocrinol Metab* 2010;95:2560-75. <https://doi.org/10.1210/jc.2009-2575>
- Ohlander SJ, Varghese B, Pastuszak AW. Erythrocytosis following testosterone therapy. *Sex Med Rev* 2018;6:77-85. <https://doi.org/10.1016/j.sxmr.2017.04.001>
- Kim SD, Cho KS. Obstructive sleep apnea and testosterone deficiency. *World J Mens Health* 2019;37:12-8. <https://doi.org/10.5534/wjmh.180017>
- Lundy SD, Parekh NV, Shoskes DA. Obstructive sleep apnea is associated with polycythemia in hypogonadal men on testosterone replacement therapy. *J Sex Med* 2020;17:1297-303. <https://doi.org/10.1016/j.jsxm.2020.03.006>
- Hanafy HM. Testosterone therapy and obstructive sleep apnea: Is there a real connection? *J Sex Med* 2007;4:1241-6. <https://doi.org/10.1111/j.1743-6109.2007.00553.x>
- Payne K, Lipshultz LI, Hotaling JM, et al. Obstructive sleep apnea and testosterone therapy. *Sex Med Rev* 2021;9:296-303. <https://doi.org/10.1016/j.sxmr.2020.04.004>
- Wilson JJ, Mick R, Wei SJ, et al. Clinical trial resources on the internet must be designed to reach underrepresented minorities. *Cancer J* 2006;12:475-81. <https://doi.org/10.1097/00130404-200611000-00007>

CORRESPONDENCE: Dr. Ranjith Ramasamy, Desai Sethi Urology Institute, Miller School of Medicine, University of Miami, Miami, FL, United States; Ramasamy@miami.edu