

**Prognostic model using postoperative normalization of C-reactive protein levels in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy**

Jun Teishima, Junichiro Hirata, Takuya Toge, Riku Uematsu, Yoshie Mita, Takahiko Yoshii, Ichiro Nakamura

Department of Urology, Kobe City Hospital Organization, Kobe City Medical Center West Hospital, Kobe, Japan

**Cite as:** Teishima J, Hirata J, Toge T, et al. Prognostic model using postoperative normalization of C-reactive protein levels in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy. *Can Urol Assoc J* 2023 November 23; Epub ahead of print.

<http://dx.doi.org/10.5489/cuaj.8393>

Published online November 23, 2023

**Corresponding author:** Dr. Jun Teishima, Department of Urology, Kobe City Hospital Organization Kobe City Medical Center West Hospital, Kobe, Japan; @med.kobe-u.ac.jp

\*\*\*

**ABSTRACT**

**Introduction:** To improve the prediction of outcomes in patients who will undergo radical nephroureterectomy (RNU) for upper tract urothelial carcinoma (UTUC), we investigated the preoperative prognostic factors and developed a risk classification model.

**Methods:** A total of 144 patients who underwent RNU with history of neither neoadjuvant nor adjuvant chemotherapy between 2008 and 2022 were retrospectively reviewed. Associations between perioperative/clinicopathological factors and outcomes, including cancer-specific survival (CSS), were assessed. We specifically focused on preoperative serum C-reactive protein (CRP) and its postoperative normalization.

**Results:** Non-normalization of postoperative serum CRP level and pathological T3 stage were identified as independent predictive factors of shorter CSS in univariate and multivariate analysis ( $p=0.0150$  and  $0.0037$ , hazard ratio: 3.628 and 4.470, respectively). We classified the patients into three groups using these factors and found that five-year CSS was 88, 42.5, and 0% in the low-risk group (0 factors), intermediate-risk group (one factor), and high-risk group (two factors), respectively ( $p<0.0001$ ).

**Conclusions:** Non-normalization of postoperative serum CRP level and pathological T stage were identified as independent postoperative prognostic factors in patients with UTUC who underwent RNU. These factors can stratify three prognostic groups and may help urologists in clinical decision-making for adjuvant therapy.

## INTRODUCTION

Upper tract urothelial carcinoma (UTUC) accounts for 5 to 10% of urothelial carcinoma [1]. Radical nephroureterectomy (RNU) has been a standard surgical option for UTUC free from metastasis. Five-year cancer-specific survival (CSS) is less than 50% for cases with pathological T2 and T3 (pT2/3) stages and less than 10% for those with pT4 stage [2]. Randomized control trials (RCTs) demonstrated that adjuvant systemic therapy using combination chemotherapy or an immune-checkpoint inhibitor (ICI) improved survival after RNU with high risk [3][4]. With such effective adjuvant therapies being reported, it is becoming increasingly important to predict the postoperative prognosis of UTUC and identify high-risk UTUCs for which adjuvant therapy is warranted. Inflammatory responses are known to reflect the grade of malignancy in various cancers, and previous reports have shown that inflammation-related markers, such as C-reactive protein (CRP) and Neutrophil-to-lymphocyte ratio, can be useful in predicting cancer prognosis and treatment response [5][6]. CRP is a representative inflammation-related marker and has been reported as a biomarker for urological malignancies [7]. The goal with this study was to identify factors that can predict the postoperative prognosis of UTUC, focusing on CRP, one of the representative inflammation-related markers, and its postoperative changes and develop a risk-classification model.

## METHODS

### Patients

This retrospective study was approved by the Ethics Committee of Kobe City Medical Center West Hospital (authorization number: 22-019). The medical records of patients who underwent RNU for unilateral UTUC at Kobe City Medical Center West Hospital between January 2009 and December 2019 were retrospectively reviewed. RNU was conducted using a laparoscopic approach with open distal ureteric excision in all patients. Patients who received neoadjuvant (NAC) or adjuvant chemotherapy (AC) were excluded from this study. Regional lymphadenectomy was conducted at the discretion of the surgeon, and the template of lymphadenectomy was used as described by Kondo et al. [8]. Using these criteria, 144 patients were included in this study. We obtained relevant clinicopathological data from medical records, including age, sex, tumor location, pathological TNM stage, tumor grade, concomitant carcinoma in situ and lymphovascular invasion. We focused on preoperative serum CRP and its postoperative normalization. Preoperative C-reactive protein was measured one month before surgery, along with other preoperative screening tests. Although some patients developed infections or other problems underwent multiple measurements of CRP, the postoperative CRP value was defined as the value measured one month after surgery. Patients whose preoperative CRP level was 0.3 mg/dL or lower were classified in the “low-CRP group”. and others were in the “higher CRP group”. The higher CRP group was further classified into two groups; those whose serum CRP level decreased to 0.3 mg/dL or lower at one month after RNU as the

“normalized CRP group” and those whose CRP level was higher than 0.3 mg/dl at one month after RNU was defined as the “non-normalized CRP group.”

### Followup regimen

Our follow-up protocol consisted of urine analysis and chest–abdomen–pelvis CT scans, with or without contrast, every 3 to 6 months for at least 5 years. Cystoscopy and urinary cytology were conducted at 3 months as the screening for intravesical recurrence. If negative, cystoscopy and cytology were repeated every 3 months for 2 years, every 6 months thereafter until 5 years, then annually. Disease progression was defined as local failure at the operative site, regional lymph node metastasis, or distant metastasis. Intravesical recurrence was not considered as disease progression.

### Statistical analysis

Differences in the distribution of variables among groups were analyzed by conducting a chi-square test for categorical variables and Mann Whitney test for continuous variables. Recurrence-free survival (RFS) and CSS probabilities were estimated using the Kaplan-Meier method, and differences between groups were assessed using the log-rank testing. The Cox proportional hazards regression model was used for multivariate analyses. All statistical analyses were conducted using the Statview 5.0 software (Abacus Concepts, Inc., Berkeley, CA, USA), and p values less than 0.05 were determined as statistically significant.

## RESULTS

This study cohort consisted of 144 patients undergoing RNU for UTUC. The characteristics of these patients are listed in Table 1. The median age of patients was 76 (50–93). Ninety-eight (68.1%) and 46 (31.9%) patients were categorized in the lower CRP and higher CRP group, respectively. There was no significant difference in patient background between these two groups. Of the patients in the higher CRP group, 27 (58.7%) and 19 (41.3%) were classified in the normalized CRP and non-normalized CRP group, respectively. There were no significant differences in patient background between these two groups (Table 2).

The CSS and RFS curves after RNU for the entire cohort of patients are shown in Fig. 1. The median of the observation period was 38.6 months, and 118 of the 144 patients (81.9%) survived. The 1-, 2- and 3-year CSS rates for the entire cohort were 97.7, 89.3, and 83.3%, respectively. The 1-, 2- and 3-year RFS rates for the entire cohort were 88.8, 81.8, and 81.8%, respectively. There was no significant difference in the CSS for the lower CRP group compared with those for the normalized CRP group ( $p=0.9406$ ) (Fig. 2a), while the CSS rates for the non-normalized postoperative CRP group were significantly worse than those for the others ( $p=0.0107$ ) (Fig. 2b). Both the CSS and RFS rates for the group with pT3/4-stage group were significantly worse than those with pT1/2 stages ( $p<0.0001$ ) (Fig. 2c, 3c), while there was no significant difference in RFS rates between the group with non-normalized postoperative CRP and those with others ( $p=0.1210$ ) (Fig. 3a, b). Non-normalization of postoperative serum CRP

level and pT3 stage were identified as independent predictive factors of shorter CSS in univariate and multivariate analysis (Table 3). We classified into four groups based on these two factors; no applicable factor, CRP non-normalization only, pT3/4 only, and both factors, so that the difference in weight of the two factors can be seen (Fig 4a). As the results showed no significant difference in the CSS for the two groups with only one applicable factor ( $p=0.3650$ ), we classified the patients into three groups using the number of applicable factors and found that 5-year CSS was 88, 42.5, and 0% in the low-risk group (0 factors), intermediate-risk group (one factor), and high-risk group (two factors), respectively. ( $p<0.0001$ ) (Fig. 4b).

## DISCUSSION

We determined the predictive factors for the prognosis in patients with UTUC undergoing RNU then developed a prognostic model on the basis of the involvement of risk factors, pT and normalization of CRP. To the best of our knowledge, this is the first report on a prognostic model of UTUC after RNU on the basis of these two factors.

UTUC includes cases with rapid progression and poor prognosis after surgery and require some optional treatments such as AC or NAC. For muscle-invasive bladder cancer (MIBC), NAC has been shown to improve prognosis in the previous RCT for MIBC [9], and NAC also may be an option for UTUC, considering the decline in renal function after RNU [10]. While there have been efforts to improve the accuracy of the staging examination of UTUC [11], it is difficult to accurately assess the risk of UTUC by preoperative imaging. Therefore, to avoid overtreatment for UTUC in addition to RNU, adjuvant therapy for selective patients based on postoperative information is appropriate to improve prognosis [12]. However, even AC for UTUC was controversial. Certain studies demonstrated the effect of AC on reducing distant metastases [13] and intravesical recurrence [14], while other retrospective studies reported no impact on prognosis [15, 16]. An RCT reported the results of postoperative adjuvant therapy using an ICI for locally advanced UC [3]. That study demonstrated the benefit mainly for patients with MIBC who often receive NAC. Another RCT, however, showed the benefit of AC with gemcitabine and cisplatin anticancer chemotherapy for patients with UTUC who were not treated with NAC [4]. Advances in postoperative systemic therapy for UTUC is expected to improve oncological outcomes after RNU. To decrease the risk of adverse events caused by such modalities, however, the choice of patients who undergo adjuvant therapy is an important issue. Therefore, the goal with this study was to analyze prognostic factors after RNU to identify UTUC patients with high risk of postoperative recurrence and/or progression. The association between the prognosis of UTUC and inflammatory response has been demonstrated [17, 18]. These inflammation-related factors can reflect the presence or absence of disease progression and may be prognostic factors by comparing before and after surgery [19]. The results indicate that in addition to pT3 stage or higher, postoperative CRP non-normalization was an independent poor prognostic factor (Table 3). The finding that high CRP levels after RNU surgery are associated with poor prognosis regardless of preoperative CRP levels is consistent with the results of our previous study [20]. The association between inflammatory response and disease

activity has already been reported previously in UTUC, and high CRP levels are thought to reflect inflammatory cytokine production from the tumor. The present study focuses on elevated C-reactive protein levels after radical surgery, which may reflect the presence of micrometastases [21]. Other studies demonstrated several prognostic factors including pathological lymph node metastasis [21], histological high grade [22], lymphovascular invasion [23, 24], and hydronephrosis [17]. However, these were not determined as independent prognostic factors by multivariate analysis (Table 3). Therefore, our prognostic model uses these two factors, showing that it is possible to stratify prognosis on the basis of the number of factors. Only patients who had not undergone preoperative treatment were included, and the results should provide useful information for predicting prognosis and selecting adjuvant therapy.

There are several limitations to this study. One is that the volume of the study was small and retrospective based on the real-world data. And some other well-known predictors of poor prognosis, such as lymph node metastasis, did not carry in the present model. In this study, lymph node dissection was performed in only 47 cases (32.7% of the total), and it is thought that there were cases with pathological lymph node metastasis among those who did not undergo lymph node dissection. Although this study provides important hypothesis, external validation is required by further study in the future. While the perioperative change in CRP was a significant predictor of CSS, there was no statistical significance, only a tendency in RFS. Possible reasons for this result may be the effect of small volume of the study or variety of postoperative choice of therapeutic option. For UC with metastasis, chemotherapy and subsequent therapy using ICI or antibody-drug conjugate have been a standard option [25, 26]. Inflammatory response has been reported to be associated with the efficacy of such regimens of systemic therapy for UC [27, 28]. The inclusion of postoperative CRP as a parameter in our model may also reflect its association with the effect of systemic therapy after recurrence and may have been statistically significant in CSS rather than RFS. Another limitation is that our prognostic model is based on postoperative information and cannot predict prognosis at the preoperative time point. The indication and extent of lymph node dissection (LND) were also determined by each surgeon. Although the indications for LND remains unestablished, its impact on prognosis in patients with UTUC undergoing RNU needs to be considered. It is hoped that the results of this study will be evaluated and confirmed in well-designed, larger prospective studies.

## CONCLUSIONS

We presented a prognostic model for UTUC that takes into account the inflammatory response. With the introduction of more sophisticated and accurate imaging and new preoperative treatments, prognostic prediction will become even more important to enable individualization of treatment.

## REFERENCES

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2022. *CA Cancer J Clin* 2022;71:7-33. [https://doi: 10.3322/caac.21708](https://doi.org/10.3322/caac.21708). Epub 2022 Jan 12.
2. EAU Guidelines. Edn. presented at the EAU Annual Congress Amsterdam, 2022. <https://d56bochluxqnz.cloudfront.net/documents/full-guideline/EAU-Guidelines-on-Upper-Tract-Urothelial-Carcinoma-2022>.
3. Bajorin DF, Witjes JA, Gschwend JE, et al. Adjuvant nivolumab versus placebo in muscle-invasive urothelial carcinoma. *N Engl J Med* 2021;384:2102-14. [https://doi: 10.1056/NEJMoa2034442](https://doi.org/10.1056/NEJMoa2034442).
4. Birtle A, Johnson M, Chester J, et al. Adjuvant chemotherapy in upper tract urothelial carcinoma (the POUT trial): a phase 3, open-label, randomised controlled trial. *Lancet* 2020;395:1268-77. [https://doi: 10.1016/S0140-6736\(20\)30415-3](https://doi.org/10.1016/S0140-6736(20)30415-3). Epub 2020 Mar 5.
5. Templeton AJ, McNamara MG, Seruga B, et al. Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic review and meta-analysis. *J Natl Cancer Inst* 2014;106: e124. [https://doi: 10.1093/jnci/dju124](https://doi.org/10.1093/jnci/dju124). Print 2014 Jun.
6. Grivennikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell* 2010;140:883-99. [https://doi: 10.1016/j.cell.2010.01.025](https://doi.org/10.1016/j.cell.2010.01.025).
7. Saito K, Kihara K. C-reactive protein as a biomarker for urological cancers. *Nat Rev Urol* 2011;8:659-66. [https://doi: 10.1038/nrurol.2011.145](https://doi.org/10.1038/nrurol.2011.145).
8. Kondo T, Hara I, Takagi T. Therapeutic role of template-based lymphadenectomy in urothelial carcinoma of the upper urinary tract. *World J Clin Oncol* 2015;6:237-51. [https://doi: 10.5306/wjco.v6.i6.237](https://doi.org/10.5306/wjco.v6.i6.237).
9. Grossman HB, Natale RB, Tangen CM, et al. Neoadjuvant chemotherapy plus cystectomy compared with cystectomy alone for locally advanced bladder cancer. *N Engl J Med* 2003;349:859-66. [https://doi: 10.1056/NEJMoa022148](https://doi.org/10.1056/NEJMoa022148).
10. Matin SF, Margulis V, Kamat A, et al. Incidence of downstaging and complete remission after neoadjuvant chemotherapy for high-risk upper tract transitional cell carcinoma. *Cancer* 2010;116:3127-34. [https://doi: 10.1002/cncr.25050](https://doi.org/10.1002/cncr.25050).
11. Honda Y, Goto K, Sentani K, et al. T categorization of urothelial carcinomas of the ureter with CT: preliminary study of new diagnostic criteria proposed for differentiating T2 or lower from T3 or higher. *AJR Am J Roentgenol* 2015;204:792-7. [https://doi: 10.2214/AJR.14.13167](https://doi.org/10.2214/AJR.14.13167).
12. Alva AS, Matin SF, Lerner SP, et al. Perioperative chemotherapy for upper tract urothelial cancer. *Nat Rev Urol* 2012;9:266-73. [https://doi: 10.1038/nrurol.2012.57](https://doi.org/10.1038/nrurol.2012.57).
13. Bamias A, Deliveliotis Ch, Fountzilas G, et al. Adjuvant chemotherapy with paclitaxel and carboplatin in patients with advanced carcinoma of the upper urinary tract: a study by the Hellenic Cooperative Oncology Group. *J Clin Oncol* 2004;22:2150-4. [https://doi: 10.1200/JCO.2004.09.043](https://doi.org/10.1200/JCO.2004.09.043).
14. Soga N, Arima K, Sugimura Y. Adjuvant methotrexate, vinblastine, adriamycin, and cisplatin chemotherapy has potential to prevent recurrence of bladder tumors after surgical removal of upper urinary tract transitional cell carcinoma. *Int J Urol* 2008;15:800-3. [https://doi: 10.1111/j.1442-2042.2008.02114.x](https://doi.org/10.1111/j.1442-2042.2008.02114.x). Epub 2008 Jul 10.


15. Lee SE, Byun SS, Park YH, et al. Adjuvant chemotherapy in the management of pT3N0M0 transitional cell carcinoma of the upper urinary tract. *Urol Int* 2006;77:22-6. [https://doi: 10.1159/000092930](https://doi.org/10.1159/000092930).
16. Hellenthal NJ, Shariat SF, Margulis V, et al. Adjuvant chemotherapy for high risk upper tract urothelial carcinoma: results from the Upper Tract Urothelial Carcinoma Collaboration. *J Urol* 2009;182:900-6. [https://doi: 10.1016/j.juro.2009.05.011](https://doi.org/10.1016/j.juro.2009.05.011).
17. Bajorin DF, Witjes JA, Gschwend JE, et al. Adjuvant nivolumab versus placebo in muscle-invasive urothelial carcinoma. *N Engl J Med* 2021;384:2102-14. [https://doi: 10.1056/NEJMoa2034442](https://doi.org/10.1056/NEJMoa2034442).
18. Kohada Y, Hayashi T, Goto K, et al. Preoperative risk classification using neutrophil-lymphocyte ratio and hydronephrosis for upper tract urothelial carcinoma. *Jpn J Clin Oncol* 2018;48:841-50. [https://doi: 10.1093/jjco/hyy084](https://doi.org/10.1093/jjco/hyy084).
19. Obata J, Kikuchi E, Tanaka N, et al. C-reactive protein: a biomarker of survival in patients with localized upper tract urothelial carcinoma treated with radical nephroureterectomy. *Urol Oncol* 2013;31:1725-30. [https://doi: 10.1016/j.urolonc.2012.05.008](https://doi.org/10.1016/j.urolonc.2012.05.008). Epub 2012 Nov 8.
20. Teishima J, Ohara S, Shinmei S, et al. Normalization of C-reactive protein levels following cytoreductive nephrectomy in patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitors is associated with improved overall survival. *Urol Oncol* 2018;36:339.e9-15. [https://doi: 10.1016/j.urolonc.2018.04.008](https://doi.org/10.1016/j.urolonc.2018.04.008).
21. Tanaka N, Kikuchi E, Shirotake S, et al. The predictive value of C-reactive protein for prognosis in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy: a multi-institutional study. *Eur Urol* 2014;65:227-34. [https://doi:10.1016/j.eururo.2012.11.050](https://doi.org/10.1016/j.eururo.2012.11.050). Epub 2012 Dec 1.
22. Chen IA, Chang CH, Huang CP, et al. Factors predicting oncological outcomes of radical nephroureterectomy for upper tract urothelial carcinoma in Taiwan. *Front Oncol* 2022;11:766576. [https://doi: 10.3389/fonc.2021.766576](https://doi.org/10.3389/fonc.2021.766576). eCollection 2021.
23. Yamada Y, Nakagawa T, Miyakawa J, et al. Subclassification of pT3 upper tract urothelial carcinoma: a multicenter retrospective study. *World J Urol* 2023;41:767-76. [https://doi: 10.1007/s00345-023-04300-7](https://doi.org/10.1007/s00345-023-04300-7). Epub 2023 Feb 4.
24. Takemoto K, Hayashi T, Hsi RS, et al. Histological variants and lymphovascular invasion in upper tract urothelial carcinoma can stratify prognosis after radical nephroureterectomy. *Urol Oncol* 2022;40: 539.e9-16. [https://doi: 10.1016/j.urolonc.2022.08.010](https://doi.org/10.1016/j.urolonc.2022.08.010). Epub 2022 Oct 13.
25. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as second-line therapy for advanced urothelial carcinoma. *N Engl J Med* 2017;376:1015-26. [https://doi: 10.1056/NEJMoa1613683](https://doi.org/10.1056/NEJMoa1613683). Epub 2017 Feb 17.
26. Rosenberg JE, O'Donnell PH, Balar AV, et al. Pivotal trial of enfortumab vedotin in urothelial carcinoma after platinum and anti-programmed death 1/programmed death ligand 1 therapy. *J Clin Oncol* 2019;37:2592-600. [https://doi: 10.1200/JCO.19.01140](https://doi.org/10.1200/JCO.19.01140). Epub 2019 Jul 29.
27. Kijima T, Yamamoto H, Saito K, et al. Early C-reactive protein kinetics predict survival of patients with advanced urothelial cancer treated with pembrolizumab. *Cancer Immunol Immunother* 2021;70:657-65. [https://doi: 10.1007/s00262-020-02709-2](https://doi.org/10.1007/s00262-020-02709-2). Epub 2020 Sep 2.

28. Saito K, Urakami S, Komai Y, et al. Impact of C-reactive protein kinetics on survival of patients with advanced urothelial carcinoma treated by second-line chemotherapy with gemcitabine, etoposide and cisplatin. *BJU Int* 2012;110:1478-84. [https://doi: 10.1111/j.1464-410X.2012.11153.x](https://doi.org/10.1111/j.1464-410X.2012.11153.x). Epub 2012 Apr 23.


DRAFT

FIGURES AND TABLES


**Figure 1.** a) Cancer-specific survival (CSS); and b) recurrence-free survival (RFS) in entire cohort.




**Figure 2.** Cancer-specific survival (CSS) stratified by a) lower, normalized, and non-normalized CRP group; b) non-normalized c-reactive protein (CRP) group and others; and c) pathological (p) T1/2 for a) and pT3/4 for b).



**Figure 3.** Recurrence-free survival (RFS) stratified by a) lower, normalized, and non-normalized CRP group; b) non-normalized CRP group and others; and c) pT1/2 and pT3/4, respectively.



**Figure 4.** a) Cancer-specific survival (CSS) stratified to four groups; no applicable factor, CRP non-normalization only, pT3/4 only, and both factors; and b) those stratified by the number of applicable risk factors.



<b>Table 1. Characteristics of patients</b>				
	<b>Lower CRP (n=98)</b>	<b>Higher CRP (n=46)</b>		<b>Total (n=144)</b>
Age, n (%)				
≤ 75 years	47 (48.0)	23 (50.0)	0.8193	70 (48.6)
> 75 years	51 (52.0)	23 (50.0)		74 (51.4)
Sex, n (%)				
Male	56 (57.1)	28 (60.9)	0.6723	84 (58.3)
Female	42 (42.9)	18 (39.1)		60 (41.7)
Tumor location, n (%)				
Renal pelvis	49 (50.0)	25 (54.3)	0.6265	74 (51.4)
Ureter	49 (50.0)	21 (45.7)		70 (48.6)
Tumor grade, n (%)				
G1/2	74 (75.5)	36 (78.3)	0.7171	110 (76.4)
G3	24 (24.5)	10 (21.7)		34 (23.6)
Pathological T stage, n (%)				
pT1/2	75 (76.5)	32 (69.6)	0.3724	107 (74.3)
pT3/4	23 (23.5)	14 (30.4)		37 (25.7)
Lymphovascular invasion, n (%)				
Absent	81 (82.7)	38 (82.6)	0.9548	119 (82.6)
Present	17 (17.3)	8 (17.4)		25 (17.4)
Lymph node stage, n (%)				
pN0	28 (28.6)	13 (28.3)	0.4123 *	41 (28.5)
pN1/2	5 (5.1)	1 (2.2)		6 (4.2)
pNx	65 (66.3)	32 (69.6)		97 (67.4)
Margin status, n (%)				
Negative	94 (95.9)	42 (91.3)	0.2597	136 (94.4)
Positive	4 (4.1)	4 (8.7)		8 (5.6)
Hydronephrosis, n (%)				
Absent	57 (58.2)	22 (47.8)	0.2451	79 (54.9)
Present	41 (41.8)	24 (52.2)		65 (45.1)
Surgical approach, n (%)				
Laparoscopic	88 (89.8)	44 (95.7)	0.2358	132 (91.7)
Open	10 (10.2)	2 (4.4)		12 (8.3)
Laterality, n (%)				
Right	44 (44.9)	13 (28.3)	0.0570	57 (39.6)
Left	54 (55.1)	33 (71.7)		87 (60.4)

\*pN0 and pNx versus pN1/2. CRP: C-reactive protein.

	<b>Normalized CRP (n=27)</b>	<b>Non-normalized CRP (n=19)</b>		<b>Total (n=46)</b>
Age, n (%)				
≤75 years	11 (40.7)	12 (63.2)	0.1848	23 (50.0)
>75 years	16 (59.3)	7 (36.8)		23 (50.0)
Sex, n (%)				
Male		14 (73.7)	0.1852	28 (60.9)
Female	13 (48.1)	5 (26.3)		18 (39.1)
Tumor location, n (%)				
Renal pelvis	15 (55.6)	10 (52.6)	0.8446	25 (54.3)
Ureter	12 (44.4)	9 (47.4)		21 (45.7)
Tumor grade, n (%)				
G1/2	20 (74.1)	16 (84.2)	0.4118	36 (78.3)
G3	7 (25.9)	3 (15.8)		10 (21.7)
Pathological T stage, n (%)				
pT1/2	18 (66.7)	14 (73.7)	0.6105	32 (69.6)
pT3/4	9 (33.3)	5 (26.3)		14 (30.4)
Lymphovascular invasion, n (%)				
Absent	22 (81.5)	16 (84.2)	0.8100	38 (82.6)
Present	5 (18.5)	3 (15.8)		8 (17.4)
Lymph node stage, n (%)				
pN0	7 (25.9)	6 (31.6)	0.3964*	13 (28.3)
pN1/2	1 (3.7)	0 (0)		1 (2.2)
pNx	19 (70.4)	13 (68.4)		32 (69.6)
Margin status, n (%)				
Negative	26 (96.3)	16 (84.2)	0.1520	42 (91.3)
Positive	1 (3.7)	3 (15.8)		4 (8.7)
Hydronephrosis, n (%)				
Absent	14 (51.9)	8 (42.1)	0.5147	22 (47.8)
Present	13 (48.1)	11 (57.9)		24 (52.2)
Surgical approach, n (%)				

Laparoscopic	26 (96.3)	18 (94.7)	0.7984	44 (95.7)
Open	1 (3.7)	1 (5.3)		2 (4.4)
Laterality. n (%)				
Right	9 (33.3)	4 (21.1)	0.3624	13 (28.3)
Left	18 (66.7)	15 (78.9)		33 (71.7)

\*pN0 and pNx versus pN1/2. CRP: C-reactive protein.

	Univariate HR	95% CI	p	Multivariate HR	95% CI	p
Age						
<75	Reference		0.1549			
≥75	1.998	0.770– 5.185				
Sex						
male	Reference		0.9181			
female	1.044	0.458– 2.383				
Tumor location						
Renal pelvis	Reference		0.3786			
Ureter	1.449	0.634– 3.310				
Tumor grade						
G1/2	Reference		0.1069			
G3	2.069	0.849– 5.040				
Pathological T stage						
pTa/1/2	Reference		<0.0001	Reference		0.0033
pT3/4	7.120	3.060– 16.568		4.470	1.648– 12.121	
Lymphovascular invasion						
(-)	Reference		<0.0001	Reference		0.0598
(+)	6.010	2.600– 13.894		2.581	0.962– 6.930	
Margin status						

Negative	Reference		0.2642			
Positive	1.998	0.593– 6.734				
Pathological N stage						
pN0 or pNx	Reference		0.9722			
pN1	0.965	0.130– 7.184				
Hydronephrosis						
(-)	Reference		0.0632	Reference		0.3315
(+)	2.221	0.957– 5.156		1.526	0.650– 3.578	
CRP normalization						
Normalized	Reference		0.0163	Reference		0.0150
Non-normalized	3.430	1.254– 9.381		3.628	1.284– 10.254	

CI: confidence interval; CRP: C-reactive protein; HR: hazard ratio,

DRAFT