

Pediatric testicular torsion management practices

A survey of Canadian urologists

Wyatt MacNevin¹, Morgan MacDonald¹, Dawn L. MacLellan^{1,2}, Daniel T. Keefe^{1,2}

¹Department of Urology, Queen Elizabeth II Health Sciences Centre, Halifax, NS, Canada; ²Division of Pediatric Urology, IWK Health Centre, Halifax, NS, Canada

Cite as: MacNevin W, MacDonald M, MacLellan DL, et al. Pediatric testicular torsion management practices: A survey of Canadian urologists. *Can Urol Assoc J* 2024;18(6):201-7. <http://dx.doi.org/10.5489/cuaj.8644>

Published online April 2, 2024

ABSTRACT

INTRODUCTION: Pediatric testicular torsion (TT) is a urologic emergency that may result in testicular loss if left untreated. Testicular salvage is dependent on prompt intervention, and thus delays in diagnosis and management may threaten testicular viability. Knowledge of real-world Canadian practice patterns for pediatric TT will allow optimization of practices based on resource availability and geographic limitations to improve care.

METHODS: An electronic survey on pediatric TT management was distributed to Canadian urologists. Descriptive statistics were performed on respondent demographic factors, hospital policies and barriers to care, surgical approaches, and transfer practices. Respondent practice patterns were analyzed based on geographic location and training.

RESULTS: Thirty-four urologists responded, with most respondents operating a community practice. Ultrasonography (US) was frequently used to support TT diagnosis. Despite this, poor US access was often cited as a barrier to care, with particular impact on rural urologists. Neonatal patients and <10 years old were commonly transferred to a pediatric hospital for definitive management due to surgeon discomfort and hospital policies. Reported transport methods commonly included use of the patient's own vehicle or ambulance based on availability and timing.

CONCLUSIONS: Neonatal patients and patients under 10 years old are most commonly reported to be transferred to pediatric hospitals for TT management. Patients located in rural locations and at centers with limited US access may be at risk for delayed diagnosis and treatment. Pathways for prompt management of suspected TT may better serve these younger pediatric patients.

INTRODUCTION

Testicular torsion (TT) is a urologic emergency associated with sudden and severe testicular pain that may lead to testicular atrophy or loss if left untreated.¹ TT is most common in the pediatric patient population, with an incidence of 3.8–5.9 per 100 000 patients <17 years of age.^{1,2} Expedient diagnosis and surgical management of pediatric TT is crucial for maintaining testicular viability, preserving fertility, and reducing pain and distress.³ Despite the importance of prompt intervention, practice patterns in Canada can vary based on institutional and geographical factors, as well as surgeon preference and comfort.⁴ These differences in practice patterns directly lead to variability in outcomes and the quality of care that patients receive.⁴⁻⁹

Testicular salvage is directly correlated with time to intervention, thus delays in diagnosis, barriers to appropriate imaging, and transfer to a pediatric hospital when necessary, are all factors that may delay treatment and contribute to increased risk of testicular loss.^{10,11} This risk is accentuated by the extensive geographical size of Canada in relation to the number of pediatric hospitals, urologists comfortable treating pediatric TT, and limited anesthesia services for pediatric patients in rural settings. Despite this variability in TT management, there is limited data on practice patterns by Canadian urologists. With knowledge of real-world practice patterns for pediatric TT, treatment approaches can be optimized based on resource availability and geographic limitations.^{7,12}

KEY MESSAGES

- Transfer to pediatric hospital for management of testicular torsion is reported to be common for neonatal patients and those <10 years old.
- Despite ultrasound supporting most pediatric testicular torsion diagnoses, access remains a barrier to care.
- Pediatric urologists reported significantly greater comfort with the surgical management of testicular torsion compared to non-pediatric urologists for neonatal torsion and torsion in patients <10 years old.
- There were no differences in self-reported comfort in managing torsion in children 10–18 years old.

The primary objective of this study was to identify the variability in pediatric TT management based on practice type and geographic location of the treating urologist. Secondary objectives included determining the barriers to patient care based on hospital type and resource availability. Furthermore, the surgical approach for management of pediatric TT and the comfort level of urologists in treating torsion based on patient age group was explored.

METHODS

After research and ethics approval was obtained, an electronic survey was developed and distributed in English to Canadian urologists through UroComm, a monthly correspondence email sent on behalf of the Canadian Urological Association, and through advertising via social media (i.e., Twitter/X). Data was collected and stored anonymously using Opinio (Object Plant, Oslo, Norway), with responses being accepted for eight weeks. No compensation was provided for completing the survey.

Survey structure

A 30-question survey was developed based on review of the current literature on pediatric TT management.^{7,12,13} The survey consisted of questions on respondent demographic information (e.g., training, years in practice, location, and practice type), hospi-

tal-based policies and barriers in the management of pediatric TT, assessment and surgical approach to TT management, and current practices related to transferring patients to a pediatric hospital for definitive surgical management.

Statistical methods and data analysis

Survey data was exported into IBM Statistical Package for the Social Sciences (SPSS) software (version 28). Descriptive statistics were performed on respondent demographic factors, hospital policies and barriers to care, surgical approaches, and transfer practices. Chi-squared analysis and independent sample T-test was performed to analyze practice patterns based on respondent location and fellowship training. Incomplete responses were excluded from analysis. A 95% confidence interval (CI) was used for determining statistical significance. The significance threshold was set at $p=0.05$.

RESULTS

Demographics

Thirty-four urologists responded to the survey, with most respondents being located in Ontario (29.4%, $n=10/34$) and Nova Scotia (20.6%, $n=7/34$). As there are approximately 730 active Canadian urologists, this represents 4.7% of the Canadian urology community.¹⁴ Most respondents held a community practice (70.6%, $n=24/34$) and 70.6% of respondents ($n=24/34$) reported that pediatric elective cases were performed by the respondent at their hospital of practice. When asked about geographic location, there was an approximately equal representation of respondents in terms of distance of their practice to the nearest pediatric hospital (Table 1).

The majority of respondents underwent fellowship training (67.6%, $n=23/34$), with 23.5% ($n=8/34$) completing fellowship training in pediatric urology. Similarly, 38.2% ($n=13/34$) identified general urology (non-sub-specialty) as their primary specialty of practice, followed by pediatric urology (20.6%, $n=7/34$) (Table 2).

Hospital policy

When examining hospital policies and cutoffs for surgical management of pediatric TT, 35.0% ($n=9/26$) of community urologists reported that there were no age-based cutoffs or policies for pediatric patients requiring an operation at their center. Similarly, American Society of Anesthesiologists (ASA) score was not used by 35.5% ($n=12/34$) of respondents for pediatric patients. Only

Table 1. Respondent geographic and practice demographics

	Frequency (%)
Practice Location - Province/territory	
Nova Scotia	7 (20.6)
New Brunswick	6 (17.6)
Prince Edward Island	1 (2.9)
Quebec	5 (14.7)
Ontario	10 (29.4)
Manitoba	1 (2.9)
Saskatchewan	1 (2.9)
Alberta	1 (2.9)
British Columbia	1 (2.9)
Newfoundland	1 (2.9)
Northwest Territories	0 (0)
Nunavut	0 (0)
Yukon	0 (0)
Practice type (multiple selections allowed)	
Pediatric hospital	5 (14.7)
Academic - adult hospital	6 (17.6)
Academic - adult and combined pediatric hospital	3 (8.8)
Community practice	24 (70.6)
Community locum practice	1 (2.9)
Pediatric elective cases performed at site	
Yes	24 (70.6)
No	8 (23.5)
Other - low-risk (e.g., circumcision only)	2 (5.9)
Distance to nearest pediatric hospital, km	
N/A	5 (15.2)
<20	4 (10.3)
20-50	4 (10.3)
51-100	4 (10.3)
101-200	7 (21.2)
201-400	3 (9.1)
>400	6 (15.4)

Table 2. Respondent training and experience

	Frequency (%)
Years in practice	
<5 years	9 (26.5)
5-10 years	7 (20.6)
11-15 years	7 (20.6)
16-20 years	7 (20.6)
21-25 years	2 (5.9)
>25 years	2 (5.9)
Fellowship training	
Pediatric urology	8 (23.5)
Endourology and stone disease	4 (13.8)
Minimally invasive surgery, robotics	5 (17.2)
Transplant urology	2 (6.9)
Oncology	1 (3.4)
Functional and reconstructive urology	3 (10.3)
N/A	11 (37.9)
Primary subspecialty of practice	
Pediatric urology	7 (21.2)
Endourology and stone disease	6 (18.2)
Minimally invasive surgery, robotics	2 (6.1)
Transplant urology	1 (3.0)
Oncology	1 (3.0)
Functional and reconstructive urology	2 (6.1)
Andrology	1 (3.0)
N/A - general urology	13 (39.4)

Barriers to care

Access to appropriate ultrasonography (US) was reported as a barrier by 25.0% (n=8/32) of respondents (Table 3). Despite 41.2% (n=14/34) of respondents reporting that scrotal/testicular US is available all of the time at their institution, 58.8% (n=20/34) reported US access being only available during daytime hours and on special request after normal working hours, which posed as a barrier to diagnosis and care. US access being a barrier to care was more common in respondents who were located >100 km from a pediatric hospital, with 75% (n=12/16) of respondents reporting reduced access compared to 25% (n=3/12) of those located within 100 km (p=0.009).

7.7% (n=2/26) of community urologists indicated pubertal status was used for operative planning at their hospital. There were no significant differences in hospital policies regarding cutoffs for care when comparing pediatric urologists with non-pediatric urologists (p>0.05).

Table 3. Barriers to care, evaluation, and surgical management of testicular torsion by pediatric and adult urologists

	Pediatric urologist, frequency (%)	Adult urologist, frequency (%)	p
Reduced ultrasound access			
Yes	1 (14.3)	7 (28.0)	0.503
No	6 (85.7)	18 (72.0)	
Anesthesia discomfort			
Yes	0 (0.0)	3 (8.00)	0.512
No	6 (100.0)	22 (88.0)	
Hospital policies			
Yes	0 (0.0)	3 (8.00)	0.512
No	6 (100.0)	22 (88.0)	
Surgeon discomfort			
Yes	0 (0.0)	1 (4.00)	0.618
No	6 (100.0)	24 (96.0)	
Evaluation and surgical management			
Use of TWIST score routinely			
Yes	3 (42.9)	2 (8.33)	0.02
No	4 (57.1)	24 (91.7)	
Use of TWIST when diagnosis unclear			
Yes	0	0	–
No	7 (100)	26 (100)	
Use of TWIST with low suspicion			
Yes	1 (14.3)	5 (19.2)	0.624
No	6 (85.7)	21 (80.8)	
Fixation method			
1-point	1 (14.2)	1 (3.84)	0.489
2-point	3 (42.9)	9 (34.6)	
3-point	3 (42.9)	16 (61.5)	
Contralateral fixation			
Yes	6 (85.7)	1 (3.84)	0.865
No	0 (0)	21 (80.8)	
Only if proven ipsilateral torsion	1 (14.3)	4 (15.4)	

Torsion assessment and management

US use in suspected TT was always used by 67.6% (n=23/34) of respondents, with only 26.5% (n=9/34) using US solely in cases of diagnostic uncertainty. Non-pediatric urologists relied on US for diagnosis (76.9%, n=20/26) more than pediatric urologists (42.9%, n=3/4) (p=0.05). There was no difference in reliance on US for TT diagnosis based on respondent distance from a pediatric hospital for non-pediatric urologists (p=0.5). US was primarily performed by radiologists or radiology technicians (97.1%, n=33/34). Furthermore, the use of the Testicular Workup for Ischemia and Suspected Torsion (TWIST) scoring system for suspected TT was only used routinely by 15.1% (n=5/33) of respondents, with 45.5% (n=15/33) reporting that they were not familiar with the TWIST score for TT assessment.

During orchidopexy, 57.6% (n=19/33) perform three-point fixation and 36.4% (n=12/33) use two-point fixation. There were no significant differences between fixation methods between pediatric urologists and non-pediatric urologists (two-point fixation: 42.9%, n=3/7 vs. 34.6%, n=9/26; three-point fixation: 42.9%, n=3/7 vs. 61.5%, n=16/26, p=0.48). Geographic location had no significant impact on fixation method used by non-pediatric urologists (p=0.309).

Fixation of the contralateral testicle was often performed by pediatric urologists, with 85.7% (n=6/7) always performing fixation during time of surgery. Of this group, 14.3% (n=1/7) perform fixation only if there are proven signs of torsion intraoperatively. Non-pediatric urologists (80.8%, n=21/26) routinely fixate the contralateral testicle during orchidopexy for TT (p=0.865).

Urologist comfort level

Pediatric urologists self-reported significantly higher comfort levels with managing TT compared to non-pediatric urologists for both neonatal patients (4.86±0.38 vs. 1.65±0.85, p<0.01) and patients <10 years of age (5.00±0.0 vs. 4.00±1.02, p=0.015) (Table 4). There was no difference in self-reported comfort level in surgical management of TT in patients aged 10–18 years old between pediatric urologists and non-pediatric urologists (5.00±0.00 vs. 4.81±0.49, p=0.314).

Transfer practices

Surgeon discomfort was reported as a barrier to treatment by 73.1% (n=19/26) of non-pediatric urologists compared to 0% of pediatric urologists (p=0.001). Similarly, 76.9% (n=20/26) of non-pediatric urologists reported anesthesia discomfort being a barrier to treatment (p=0.016).

For patients <10 years of age and non-neonates, local treatment of torsion was most common (80.8%, n=21/26), with only 19.2% (n=5/26) transferring patients to pediatric urology. The main reasons for transfer included surgeon discomfort (15.4%, n=4/26), anesthesia discomfort (15.4%, n=4/26), and hospital policy (11.5%, n=3/26) (Table 5).

All respondents (n=26/26) would treat TT in patients aged 10–18 years old locally without transfer to pediatric urology. If transfer was required, reasons were attributed to anesthesia discomfort (11.5%, n=3/26) and hospital policy (3.8%, n=1/26).

Transfer logistics

For all patient age groups, there were no significant differences in reasons to transfer based on rural location with a distance >100 km from a pediatric hospital. When patients are transferred to pediatric urology, the patient's own vehicle (57.7%, n=15/26) and ambulance (53.8%, n=14/26) are commonly used options, with air transfer being less common (7.7%, n=2/26). The reason provided for using the patient's own transportation was increased wait time if waiting for an ambulance for transport. When estimating transfer times, 32.3% (n=11/34) of respondents reported that patients arrive at the pediatric hospital in <2 hours, on average. Approximate transfer time was <1 hour (50%, n=6/12) for patients located <100 km from a pediatric hospital. Only 25% (n=4/16) of respondents located >100 km from a pediatric hospital reported transfer times between 1–2 hours, with the majority being >3 hours (31%, n=5/16).

DISCUSSION

Barriers to care in the management of pediatric TT exist in Canada and vary based on geographic location.⁴ US access was the most cited barrier to TT care, with over 50% of respondents indicating that after-hours US access was limited and only available in special cases. Furthermore, US access was more limited in rural centers located further from a pediatric center. This is particularly important to consider, as the majority of respondents indicated that US is required prior to surgical intervention in their practice. With reduced US access and the reliance on US prior to surgical intervention, patients who present with TT after-hours or at centers with reduced US access may face prolonged time to intervention. This may lead to increased rates of orchiectomy, as emergency department (ED) wait times >1 hour for patients with TT is associated with increased orchiectomy rates.¹⁰ This risk may be further increased if patients require trans-

fer to a pediatric hospital, which may subsequently prolong care.⁸

Children aged 10 years or younger were most commonly transferred to a pediatric hospital for definitive care, with an average transfer time of two hours for those located >100 km away. As duration of symptoms <6 hours remains a significant factor associated with testicular salvage, these patients are at risk for testicular loss.¹⁵ This is in agreement with a recent study demonstrating that additional transfer time was associated with increased orchiectomy rates in pre-pubertal patients.^{10,16} To further complicate care for this patient group, US access was more limited in rural centers in which patients face increased transfer times, which may further increase the risk of testicular loss.^{8,11}

Surgeon discomfort, anesthesia discomfort, and hospital policies were all significant factors leading to transfer of patients <10 years of age with TT. When examining hospital policies, ASA score and age, and not pubertal status, were more commonly used, which aligns with current decision tools used for both the general pediatric and adult operative population.¹⁷ Self-reported confidence in treatment of TT in neonatal patients and patients <10 years of age was higher in pediatric urologists, in keeping with the reported

Table 4. Self-reported comfort in surgical management of testicular torsion by age group

	Pediatric urologist, mean ± SD	Adult urologist	p
Neonatal	4.86±0.378	1.65±0.846	<0.0001
<10 years old	5.00±0.00	4.00±1.020	0.015
10–18 years old	5.00±0.00	4.81±0.491	0.315

SD: standard deviation.

Table 5. Reasons for patient transfer by patient age group

	Pediatric urologist, frequency (%)	Adult urologist, frequency (%)	p
Patient age <10 years old			
Surgeon discomfort	0/7 (0)	4/26 (15.4)	0.01
Anesthesia discomfort	0/7(0)	4/26 (15.4)	0.01
Hospital policy	0/7(0)	3/26 (11.5)	0.190
Patient age 10–18 years old			
Surgeon discomfort	0/7 (0)	0/26 (0)	–
Anesthesia discomfort	0/7(0)	3/26 (11.5)	0.190
Hospital policy	0/7 (0)	1/26 (3.8)	0.532

transfer practices. For patients ≥ 11 years of age, there were no significant differences in self-reported comfort and preference for transfer to a pediatric center. Therefore, efforts to expedite diagnosis of TT presentations in patients ≤ 10 years of age may be a potential area for improvement, as these patients are more likely to be transferred to a pediatric hospital.^{8,10} This is especially true for neonatal torsion, which should exclusively be managed by pediatric urologists.¹⁸ Additionally, to improve urologist comfort in treating pediatric TT, more comprehensive exposure and training in residency should be considered to reduce the need for transfer to a pediatric center.

Treatment of pediatric TT is complicated for a number of reasons, including its time-sensitive nature, use of imaging to aid diagnosis, and factors such as surgeon/anesthetist discomfort and hospital policies, which may require patient transfer for treatment. Proposed pathways to improve pediatric TT management must be inclusive of all these factors.¹⁹⁻²¹

Based on the barriers to care identified in this study, ED pathways can be better optimized for children

“ It is critical to optimize ED pathways for children such that testicular exams are routinely performed on patients presenting with abdominal or groin pain so as not to delay TT diagnosis. ”

such that testicular exams are routinely performed on patients presenting with abdominal or groin pain to not misdiagnose or delay diagnosis of torsion.²² Additionally, when patients are noted to have groin/testicular pain at triage, pathways should be developed to facilitate more urgent assessment by an emergency physician, with a streamlined protocol to facilitate concurrent US assessment and urology referral to minimize ED wait time, with particular attention on younger patients who are more likely to require transfer.^{19,23}

The use of the TWIST score, a risk-based score system using features of clinical history and physical examination, should further be used to streamline ED TT care pathway.²⁴ Further education may be required for emergency physicians to become comfortable and competent in using the TWIST scoring system, which may reduce the need for US and expedite care.

To address reduced US access, advocacy efforts should be made for acquiring portable US machines in rural locations with limited US access and radiologist availability.^{5,14} This should be combined with mandatory training for urologists and emergency physicians on scrotal US in instances where radiology is not accessible.^{10,23} Dedicated point-of-care US (POCUS) training for urologists would expedite care and work to improve patient outcomes.¹⁰

To optimize patient transfer for TT treatment, site-specific guidelines must be developed based on available resources (i.e., US and radiologist availability, ambulance availability) and geographic distance to the nearest pediatric hospital. A thorough review of each center's current process for diagnosis and treatment/transfer will allow for identification of areas in TT treatment that contribute to delay, which can then be targeted for improvement. As surgeon and anesthetist discomfort were identified as areas leading to patient transfer, continuing medical education initiatives on pediatric TT management, along with more emphasis on pediatric torsion during residency training, may reduce the need for transfer and expedite care.

Lastly, population-based efforts to increase knowledge of TT is important to support the prompt presentation to the ED when patients are experiencing symptoms.^{25,26}

To our knowledge, this is one of the only Canadian studies to investigate real-world pediatric TT practices with consideration of geographic barriers and identification of potential areas of improvement for TT management. Of note, the majority of respondents in our study were community urologists with varied geographic location and practice experience, reflecting the majority of practicing urologists in Canada.¹⁴

Limitations

The limitations of this study include the low response rate of 4.7% and the higher percentage of respondents having completed fellowship training in pediatric urology (approximately 20%) than expected, likely due to the subject of this study and its broad distribution to the Canadian urology membership. This may bias the results due to the over-representation of pediatric urologists. Additionally, the low response rate means that the TT practice patterns of most community urologists remain uncaptured.

Furthermore, confidence in treatment of TT was self-reported, with no objective outcomes measured with respect to complication rate and testicular salvage, therefore, it is possible scores may not reflect

actual competence. Future studies comparing outcomes between pediatric urologists and non-pediatric urologists performing orchidopexy for pediatric TT may provide more strength to this finding.

Additionally, this study used a survey approach eliciting respondent's perspectives of their practices and did not quantify pediatric TT volume at their center, number of transfers, and/or transfer time on a case-by-case basis. Furthermore, with respect to TT evaluation, the use of US in the survey did not explicitly include POCUS, which may be helpful in qualifying POCUS use as an area of improvement. Future studies are planned to quantify these practices while examining testicular salvage rates based on ED triage and transfer times.

CONCLUSIONS

Pediatric TT requires prompt investigation and surgical management. Results from this study have identified that patients <10 years of age are more commonly transferred to pediatric urology for intervention. Furthermore, patients in rural locations located further away from a pediatric hospital, and those located at centers with reduced access to US imaging may be further at risk for testicular death and require additional efforts to improve US access and prompt transfer when necessary. With improved understanding of the real-world practices for pediatric TT in Canada, guidelines and practices may be improved to better serve these at-risk patients.

COMPETING INTERESTS: The authors do not report any competing personal or financial interests related to this work.

This paper has been peer-reviewed.

REFERENCES

1. Zhao LC, Loutz TB, Meeks JJ, et al. Pediatric testicular torsion epidemiology using a national database: Incidence, risk of orchiectomy and possible measures toward improving the quality of care. *J Urol* 2011;186:2009-13. <https://doi.org/10.1016/j.juro.2011.07.024>
2. Greear GM, Romano MF, Katz MH, et al. Testicular torsion: Epidemiological risk factors for orchiectomy in pediatric and adult patients. *Int J Impot Res* 2021;33:184-90. <https://doi.org/10.1038/s41443-020-0331-8>
3. Osumah TS, Jimbo M, Granberg CF, et al. Frontiers in pediatric testicular torsion: An integrated review of prevailing trends and management outcomes. *J Pediatr Urol* 2018;14:394-401. <https://doi.org/10.1016/j.jpuro.2018.07.002>
4. Chan EP, Wang PZ, Myslik F, et al. Identifying systems delays in assessment, diagnosis, and operative management for testicular torsion in a single-payer health-care system. *J Pediatr Urol* 2019;15:251-7. <https://doi.org/10.1016/j.jpuro.2019.03.017>
5. Abdelhalim A, Chamberlin JD, McAleer IM. A Survey of the current practice patterns of contralateral testis fixation in unilateral testicular conditions. *Urology* 2018;116:156-60. <https://doi.org/10.1016/j.urology.2018.03.008>
6. Sugrue DD, O'Connor E, Davis N. Testicular torsion in Ireland: A 10-year analysis of incidence and risk of orchiectomy. *Ir J Med Sci* 2022;191:2267-74. <https://doi.org/10.1007/s11845-021-02816-8>

7. Ryan KA, Folkard SS, Bastianpillai C, et al. The management of testicular torsion in the UK: How can we do better? A national quantitative and qualitative analysis of the factors affecting successful testicular salvage. *J Pediatr Urol* 2020;16:815. <https://doi.org/10.1016/j.jpuro.2020.08.018>
8. Preece J, Ching C, Yackey K, et al. Indicators and outcomes of transfer to tertiary pediatric hospitals for patients with testicular torsion. *J Pediatr Urol* 2017;13:388.e1-6. <https://doi.org/10.1016/j.jpuro.2017.03.034>
9. Guerra LA, Wiesenthal J, Pike J, et al. Management of neonatal testicular torsion: Which way to turn? *Can Urol Assoc J* 2013;2:376-9. <https://doi.org/10.5489/cuaj.792>
10. Romao RLP, Anderson KH, MacLellan D, et al. Point-of-care influences orchiectomy rates in pediatric patients with testicular torsion. *J Pediatr Urol* 2019;15:367. <https://doi.org/10.1016/j.jpuro.2019.04.014>
11. Steeman A, Ngatchou W, Ramadan AS, et al. Impact of treatment delays on outcome of acute testicular torsion: A 15-year retrospective study. *Acta Chir Belg* 2022;122:116-22. <https://doi.org/10.1080/00015458.2021.1883391>
12. Van Doeveren T, Somani BK, Hoen LA, et al. Real-world management of testicular torsion: Level of adherence to the EAU Pediatric Urology guidelines. *J Pediatr Urol* 2023;19:117-24. <https://doi.org/10.1016/j.jpuro.2022.09.020>
13. Pinar U, Duquesne I, Lannes F, et al. The use of doppler ultrasound for suspected testicular torsion: Lessons learned from a 15-year multicentre retrospective study of 2922 patients. *Eur Urol Focus* 2022;8:105-11. <https://doi.org/10.1016/j.euf.2021.02.011>
14. Canadian Medical Association, Specialty Profile: Urology. Canadian Medical Association. Accessed: November 29, 2023. Available: <https://www.cma.ca/sites/default/files/2019-01/urology-e.pdf>
15. Mellick LB, Sinex JE, Gibson RW, et al. A systematic review of testicle survival time after a torsion event. *Pediatr Emer Care* 2019;35:821-5. <https://doi.org/10.1097/PEC.0000000000001287>
16. Ramachandra P, Palazzi K, Holmes N, et al. Factors influencing rate of testicular salvage in acute testicular torsion at a tertiary pediatric center. *West J Emerg Med* 2015;16:190-4. <https://doi.org/10.5811/westjem.2014.11.22495>
17. Aplin S, Baines D, and De Lima J. Use of the ASA physical status grading system in pediatric practice. *Paediatr Anaesth* 2007;17:216-22. <https://doi.org/10.1111/j.1460-9592.2006.02094.x>
18. Monteilh C, Calixte R, and Burjonrappa S. Controversies in the management of neonatal testicular torsion: A meta-analysis. *J Pediatr Surg* 2020;55:815-9. <https://doi.org/10.1016/j.jpedsurg.2018.07.006>
19. Zee RS, Bayne CE, Gomella PT, et al. Implementation of the accelerated care of torsion pathway: A quality improvement initiative for testicular torsion. *J Pediatr Urol* 2019;15:473-9. <https://doi.org/10.1016/j.jpuro.2019.07.011>
20. Winters DA, Mehmi AK, Menzies-Wilson R, et al. Supporting a learning system in pediatric emergency pathways: Using organizational comparisons, serious incidents, and near-miss events to improve the diagnosis and treatment of testicular torsion. *Pediatr Emer Care* 2022;38:1402-7. <https://doi.org/10.1097/PEC.0000000000002766>
21. Arevalo MK, Sheth KR, Menon VS, et al. Straight to the operating room: An emergent surgery track for acute testicular torsion transfers. *J Pediatr* 2018;192:178-83. <https://doi.org/10.1016/j.jpeds.2017.09.009>
22. S. Vasconcelos-Castro and M. Soares-Oliveira, Abdominal pain in teenagers: Beware of testicular torsion. *J Pediatr Surg* 2020;55:1933-5. <https://doi.org/10.1016/j.jpedsurg.2019.08.014>
23. Heckscher D, Jalfon M, Buck MB, et al. Implementation of a health system intervention to reduce time from presentation to surgical intervention for pediatric testicular torsion. *J Pediatr Urol* 2023; S1477-5131(23)00498-9. <https://doi.org/10.1016/j.jpuro.2023.10.042>
24. Sheth KR, Keays M, Grimsby G, et al. Diagnosing testicular torsion before urological consultation and imaging: Validation of the TWIST score. *J Urol* 2016;195:1870-6. <https://doi.org/10.1016/j.juro.2016.01.101>
25. Alsarlar CE, Ryan SL, Donel E, et al. Standardized process to improve patient flow from the emergency room to the operating room for pediatric patients with testicular torsion. *J Pediatr Urol* 2016;12:233. <https://doi.org/10.1016/j.jpuro.2016.04.019>
26. Friedman AA, Ahmed H, Gitlin JS, et al. Standardized education and parental awareness are lacking for testicular torsion. *J Pediatr Urol* 2016;12:166. <https://doi.org/10.1016/j.jpuro.2016.01.008>

CORRESPONDENCE: Dr. Daniel T. Keefe, Division of Pediatric Urology, IWK Health Centre, Halifax, NS, Canada; Daniel.keefe@iwnshealth.ca