

Retrograde intrarenal surgery with flexible and navigable suction ureteral access sheaths vs. percutaneous nephrolithotomy for 2–3 cm kidney stones: Is less invasive better?

Recep Burak Degirmentepe¹, Yasir Muhammed Akca¹, Suleyman Alp¹, Hacı Ibrahim Cimen¹, Deniz Gul², Mustafa Erkoç³, Muammer Bozkurt³, Tuncay Toprak⁴, Fikret Halis¹

¹Sakarya University, School of Medicine, Department of Urology, Sakarya, Turkey; ²Department of Urology, Sakarya University Training and Research Hospital, Sakarya, Turkey; ³Department of Urology, University of Health Sciences Prof. Dr. Cemil Tascioglu City Hospital, Istanbul, Turkey; ⁴Department of Urology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, Istanbul, Turkey

Cite as: Degirmentepe RB, Akca YM, Alp S, et al. RIRS with FANS-UAS vs. PCNL for 2–3 cm kidney stones: Is less invasive better? *Can Urol Assoc J* 2025 September 23; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.9279>

Published online September 23, 2025

Corresponding author: Dr. Recep Burak Degirmentepe, Sakarya University, School of Medicine, Department of Urology, Sakarya, Turkey; burakdegirmentepe@gmail.com

ABSTRACT

Introduction: The management of renal stones measuring 2–3 cm remains a clinical challenge, with percutaneous nephrolithotomy (PCNL) traditionally favored over retrograde intrarenal surgery (RIRS) due to higher stone-free rates (SFR); however, advancements in flexible ureteroscopy and the introduction of flexible and navigable suction ureteral access sheaths (FANS-UAS) have expanded the role of RIRS for larger stones. This study compares the efficacy and safety of PCNL vs. FANS-UAS-assisted RIRS for medium-sized renal calculi.

Methods: This retrospective study included 116 patients (50 RIRS, 66 PCNL) with 2–3 cm renal stones treated at a single institution. RIRS was performed using an 11/13 Fr FANS-UAS, while PCNL followed standard techniques. Outcomes assessed included operative time, hospitalization, hemoglobin drop, SFR (defined as no fragments ≥ 1 mm on three-month computed tomography), and complications (Clavien-Dindo).

Results: Baseline characteristics were comparable between groups ($p > 0.05$). RIRS demonstrated significantly shorter operative times (63.1 ± 11.3 vs. 97.3 ± 15.1 minutes, $p < 0.001$) and hospitalization (28.3 ± 17.9 vs. 81 ± 24.8 hours, $p < 0.001$), with less hemoglobin drop (0.4 ± 0.5 vs. 2.1 ± 1.7 g/dL, $p < 0.001$). SFRs were similar at three months (70% RIRS vs. 74.2% PCNL, $p = 0.677$). PCNL had higher complication rates, including three grade III events (vs. none in RIRS), although not statistically significant ($p = 0.277$).

Conclusions: FANS-UAS-assisted RIRS offers comparable SFRs to PCNL for 2–3 cm stones, with advantages in operative efficiency, hospitalization, and perioperative safety. These findings suggest RIRS as a viable minimally invasive alternative, warranting further randomized trials to validate its role in this setting.

INTRODUCTION

The growing burden of kidney stone disease presents significant challenges to healthcare systems worldwide, as it increasingly affects diverse patient populations regardless of demographic factors¹. Surgical treatment has transitioned from open nephrolithotomy to minimally invasive procedures such as retrograde intrarenal surgery (RIRS) and percutaneous nephrolithotomy (PCNL), driven by technological advancements and innovations. The primary goal of these developments is to enhance treatment success by employing less invasive methods and enabling patients to return to everyday life as quickly as possible². Stone size remains a key factor in determining the appropriate minimally invasive treatment. European Association of Urology (EAU) guidelines position RIRS as the preferred intervention for nephrolithiasis cases with stone burdens below 2 cm^{3,4}. Current literature reveals heterogeneity in the management of >2 cm renal stones, with high-volume centers increasingly reporting RIRS outcomes rivaling PCNL for select cases, challenging traditional size-based treatment paradigms.^{1,5}

PCNL is highly effective in achieving stone-free outcomes for large kidney stones but is considered more invasive compared to RIRS. This is due to potential complications such as bleeding, the need for blood transfusion or angioembolization, and the risk of organ damage during access, given the kidney's proximity to the lungs, spleen, intestines, and liver². However, with the introduction of flexible and navigable suction ureteral access sheath (FANS-UAS), RIRS has been effectively and safely applied to large kidney stones, offering shorter operative times and reduced postoperative complication rates⁶. This study aimed to evaluate and compare the outcomes of PCNL and RIRS, in which the FANS-UAS was utilized, for the management of renal stones measuring 2–3 cm. The principal aim of this study was to investigate the clinical safety and effectiveness of the increasingly adopted FANS-UAS-assisted RIRS technique in comparison to PCNL for managing medium-sized renal calculi.

METHODS

This retrospective study included patients with kidney stones measuring 2–3 cm in diameter who underwent either RIRS with FANS-UAS or PCNL at Sakarya University between November 2023 and December 2024. The study was approved by the institutional ethics committee (approval no: E-43012747-050.04-450925-101). Patients with renal anomalies or those who underwent preoperative nephrostomy were not included in the study. Only patients undergoing their first surgical intervention for renal stones (either primary or recurrent stone formers) were included, while those with prior stone-related procedures (e.g., ureteroscopy, PCNL, or shockwave lithotripsy) were excluded. This criterion was implemented to eliminate

potential confounding effects from previous interventions on operative outcomes. Treatment modality (RIRS vs. PCNL) was assigned based on a standardized institutional protocol, considering stone characteristics (location, density in Hounsfield units), renal anatomy, and patient comorbidities. All cases were discussed in a multidisciplinary team meeting involving senior urologists with ≥ 10 years of experience in endourology. The procedures were performed by three fellowship-trained endourologists, each having performed >200 RIRS and >150 PCNL procedures independently. To minimize bias, surgeons proficient in both techniques participated in the study. Treatment allocation was determined through a shared decision-making process that integrated clinical parameters with patient preferences. For all patients meeting inclusion criteria, both RIRS and PCNL options were thoroughly discussed during preoperative counseling. This discussion included: comparative risks (bleeding, infection, need for secondary procedures), anticipated recovery times, and stone-free probability based on individual stone characteristics (HU density >1000 favoring PCNL, unfavorable infundibulopelvic angles suggesting PCNL). Special consideration was given to comorbidities (e.g., bleeding diatheses automatically excluding PCNL). Final treatment selection reflected joint agreement between the surgical team and informed patient choice after reviewing all options. This standardized counseling protocol ensured consistent evaluation while respecting patient autonomy.

The patients were assigned to two groups: Group 1 ($n = 52$) underwent RIRS with the assistance of an 11/13 Fr FANS-UAS (ClearPetra) as the ureteral access sheath, whereas Group 2 ($n = 66$) underwent standard PCNL. Two patients in the RIRS group continued their postoperative care at another center, leading to incomplete follow-up data, and were not included in the final analysis. Consequently, the study included 116 patients. Demographic data, including sex, age, and body mass index (BMI), along with stone-specific parameters such as location, size, and Hounsfield unit (HU) values, were recorded for each patient. Preoperative urine culture results and hemoglobin levels were also documented. Prior to the operation, all patients included in the study had sterile preoperative urine cultures, ensuring no underlying infections.

A standard conventional approach was employed for all PCNL procedures. Following the administration of general anesthesia, patients were positioned in the dorsal lithotomy position to begin the procedure. Cystoscopy was utilized to access the bladder, and the ureteral orifice on the operative side was identified. A 5Fr ureteral catheter was then advanced over a guidewire. The external portion of the ureteral catheter was secured to the Foley catheter with silk sutures. After completing the initial steps, the patient's position was changed to prone to allow access for the percutaneous phase of the procedure. Contrast material was administered through the ureteral catheter, and the renal pelvicalyceal system was visualized using a C-arm fluoroscope. Percutaneous access to the kidney was achieved with an 18G Chiba needle. After the guidewire was advanced into the pelvicalyceal system, Amplatz dilatation was utilized, and a 30 Fr access sheath was subsequently inserted to complete the tract preparation. A nephroscope (26 F Storz; Karl Storz GmbH & Co. KG, Tuttlingen, Germany) was used to access the stones within the pelvicalyceal system. Stone fragmentation was achieved using a combination of ultrasonic and pneumatic lithotripters

(Lithobox Zero), and the resulting small fragments were removed manually with forceps. At the conclusion of the procedure, a nephrostomy catheter was placed in all patients under C-arm fluoroscopy guidance.

In the RIRS procedure, before the insertion of FANS-UAS, the ureter was accessed using an 8 Fr semirigid ureteroscope (Karl Storz, Tuttlingen, Germany) and hydrodilatation was performed up to the renal pelvis. The approximate length of the ureter was calculated using the ureteroscope, and the FANS-UAS was then placed over the guide without fluoroscopy. In cases where the FANS-UAS could not be advanced to the renal pelvis, a double-J catheter was placed via the guide, and the procedure was postponed for 3 weeks. Three patients had their operations delayed for 3 weeks due to the inability to advance the FANS-UAS during the first attempt. However, the FANS-UAS was successfully placed without complications in their second session. Subsequently, the Hugemed (HU 30S 7.5 Fr) flexible ureteroscope (FURS) was passed through the flexible access sheath to reach the pelvicalyceal system. The stones were visualized and dusted using a 273 μm laser with a Ho-YAG (Quanta 30) device. Depending on the stone density, the fragmentation was carried out using energy settings of 1-2 joules, frequency 5-10 Hz, and power 5-20 W. The stones were thoroughly dusted until no fragments remained, allowing small residual pieces and dust to pass spontaneously. Following the procedure, a 4.8 Fr Double-J catheter was placed over the guide, and the process was concluded.

The total operation time was defined as the duration from the moment the bladder was accessed with the ureteroscope (URS) after anesthesia induction until the Double-J catheter was placed at the end of the procedure for RIRS, and from the moment the bladder was accessed with cystoscopy until a nephrostomy catheter was inserted to ensure postoperative drainage in PCNL. For the three patients requiring delayed FANS-UAS placement (initial placement failure), only the operative time of the definitive RIRS procedure was included in the primary analysis. The duration of initial DJ stent placement (mean: 8 ± 2 minutes) was excluded as it represented a separate, minimal intervention. A decrease in hemoglobin was calculated as the difference between the preoperative hemoglobin value and the first hemoglobin measurement taken postoperatively. Stone-free status was assessed through a two-phase imaging protocol. On postoperative day 1, all patients with radio-opaque stones underwent KUB radiography, where fragments < 2 mm were considered clinically insignificant based on KUB's resolution limits. For non-opaque stones or when clinical suspicion persisted, low-dose CT was obtained. At the 3-month follow-up, every patient received standardized low-dose CT (2 mm slice thickness) which was graded according to Journal of Endourology criteria: complete clearance (Grade A, no fragments ≥ 1 mm), clinically insignificant residual fragments (Grade B, 1-3 mm), and significant residual stones (Grade C, ≥ 4 mm). The success rate in this study was defined exclusively as Grade A clearance - the complete absence of any residual stone fragments ≥ 1 mm - as determined by standardized low-dose CT imaging at the 3-month postoperative follow-up. The total operation time was defined as the duration from when the bladder was accessed with the URS after anesthesia induction until the double-J catheter was placed at the end of the procedure for RIRS. PCNL was measured from the moment the bladder was accessed with cystoscopy

until the nephrostomy catheter was placed. Since all RIRS procedures were performed without fluoroscopy, no comparison of fluoroscopy times was made between the groups. Postoperative complications occurring in the early period were evaluated and graded using the Clavien-Dindo system.

Statistical analysis was conducted using SPSS software (version 22.0; IBM Corp., Armonk, NY). Categorical variables were analyzed using the Chi-square test or Fisher's exact test, based on expected cell counts. For continuous variables, the Student's t-test was applied when assumptions of normality were met; otherwise, the Mann-Whitney U test was used. A two-sided p-value <0.05 was considered indicative of statistical significance. As this retrospective study compared existing techniques, a formal power calculation was not performed a priori. However, post-hoc analysis determined the study had 98% power ($\alpha=0.05$, $\beta=0.20$) to detect a 34-minute operative time difference—exceeding the 20-minute threshold considered clinically significant based on prior studies linking shorter OR times to reduced anesthesia risks⁴.

RESULTS

The distribution and comparison of baseline demographic parameters between the RIRS and PCNL groups are detailed in Table 1. Variables such as sex, BMI, age, stone size, HU, and the location of the stones within the pelvicalyceal system were analyzed. All comparisons yielded non-significant results, indicating that the groups were comparable in terms of baseline characteristics ($p > 0.05$).

Table 2 provides an overview of the postoperative results observed in each group. Hospitalization times were significantly shorter in the RIRS group (28.3 ± 17.9 hours) compared to the PCNL group (81 ± 24.8 hours) ($p < 0.001$). Similarly, operation times were significantly shorter in the RIRS group (63.1 ± 11.3 minutes) compared to the PCNL group (97.3 ± 15.1 minutes) ($p < 0.001$). Sensitivity analysis including cumulative times (initial stent placement + definitive RIRS) for the three staged cases showed no significant difference in mean RIRS operative time (65.4 ± 12.7 vs original 63.1 ± 11.3 minutes, $p = 0.421$), confirming the robustness of our findings.

The hemoglobin decrease was significantly lower in the RIRS group (0.4 ± 0.5 g/dL) than in the PCNL group (2.1 ± 1.7 g/dL; $p < 0.001$). Clinically, 3 patients (4.5%) in the PCNL group required blood transfusions (all for hemoglobin <7 g/dL with symptomatic anemia), whereas no RIRS patients needed transfusion. Stone-free rates assessed using KUB or low-contrast CT on postoperative day 1 were 76% in the RIRS group and 81.8% in the PCNL group ($p = 0.492$). At postoperative month 3, although the stone-free success rate was numerically higher in the PCNL group (74.2%) compared to the RIRS group (70%), the difference was not statistically significant ($p = 0.677$).

To assess the severity and distribution of postoperative complications, the Clavien-Dindo classification was applied across both groups. In the RIRS group, four patients experienced Grade I complications, while eight patients in the PCNL group experienced Grade I complications. Fever as a Grade II complication was observed in two patients in the RIRS group, and antibiotic therapy was initiated. In the PCNL group, six patients had Grade II complications, of whom three experienced fever requiring antibiotics, and three required

blood transfusions. No Grade III complications were observed in the RIRS group, whereas three patients in the PCNL group experienced Grade III complications. Double-J stents were placed in two patients due to urine leakage lasting more than 12 hours, and one patient underwent treatment by interventional radiology for an arteriovenous fistula. Both groups were free from Grade IV and V complications during the postoperative period. No statistically significant difference was found between the groups regarding complications classified by the Clavien-Dindo system ($p=0.277$).

DISCUSSION

The principal goal in urinary stone treatment is to ensure maximal stone clearance while reducing morbidity to the lowest possible level. Consequently, endourological techniques, with reduced invasiveness, have become the standard in contemporary urinary stone management⁷. PCNL marked a paradigm shift in kidney stone surgery, rendering open kidney stone surgery obsolete today. Its most significant advantage lies in being a minimally invasive surgical technique that allows the introduction of large instruments into the renal pelvicalyceal system via a nephroscope to fragment and remove stones from the body⁸. However, despite its advantages, PCNL involves controlled renal trauma, as dilation up to 30 Fr is performed, which increases the risk of complications such as bleeding and cortical tears. Although technological advancements have reduced the size of instruments used in PCNL since its inception, it remains a procedure with a risk of complications despite its high stone-free rates⁸. The American Urological Association (AUA) and EAU guidelines currently advocate for the use of PCNL as the treatment of choice for kidney stones larger than 2 cm, primarily due to its superior ability to achieve stone-free outcomes⁹.

Recent advancements in laser technologies and flexible ureteroscopes have introduced a novel approach to the surgical management of large kidney stones. RIRS has become a cornerstone in treating kidney stones due to its high reliability, effectiveness, and low morbidity rates¹⁰. This technique is increasingly preferred for managing kidney stones across all anatomical locations and age groups. It is particularly advantageous for patients with bleeding or coagulation disorders and those with renal anomalies, making it the preferred treatment option in such cases¹¹. In RIRS, as stone size increases, operation time tends to lengthen, and the risk of complications such as urinary tract perforation, renal effusion, urinary tract infection, and uremic syndrome rises³. RIRS may not be able to fully eliminate stones larger than 2 cm in a single treatment session. Studies indicate that single-session stone clearance rates can drop to as low as 62% as stone size increases¹². In cases where complete clearance is not achieved, prolonged operative times may lead to additional surgical interventions or complications, such as infection and bleeding, further increasing the risk¹².

Ureteral access sheaths (UAS) have been introduced recently to reduce surgical complications associated with RIRS¹³. However, traditional UAS are unable to reach the calyces within the pelvicalyceal system, making it challenging to completely clear large calyceal stones. UAS with a suction bendable tip has been developed to address this limitation. These FANS-UAS can be maneuvered into the calyces using a flexible ureteroscope, allowing fragmented stones to be efficiently removed through suction.

Furthermore, these devices have been shown to reduce operative time, prevent elevated intrarenal pressure, and decrease complications such as sepsis and bleeding³.

FANS-UAS can be flexed at wide angles, allowing access to most calyces within the pelvicalyceal system through the ureteroscope. Additionally, negative pressure aspiration enhances the efficiency of fluid circulation in the pelvicalyceal system, providing a clear surgical vision while effectively removing stone fragments from the body³. This study directly compares two different surgical approaches for managing kidney stones between 2–3 cm in size: PCNL, the established treatment for larger stones, and RIRS using FANS-UAS, an emerging procedure gaining popularity due to its various advantages.

Recently, there has been a significant increase in research focused on examining the efficacy and safety of FANS-UAS in RIRS procedures for both adult and pediatric patients. In a recent retrospective study by Turedi et al.¹⁴, pediatric patients who underwent RIRS were compared between FANS-UAS and traditional UAS groups. The FANS-UAS group demonstrated shorter operative times, reduced anesthesia requirements, fewer treatment sessions, and higher initial success rates. No meaningful differences in complication rates were observed between the groups. Gauhar et al.¹⁵ published a multicenter prospective study conducted across eight centers, comparing conventional UAS with FANS-UAS in RIRS surgeries for pediatric patients. The study concluded that RIRS procedures performed with FANS-UAS were safe, effective in achieving stone clearance, and associated with a low complication rate. In an international multicenter randomized prospective study, Zhu et al.¹⁶ compared the efficacy and safety of FANS-UAS with traditional UAS for renal and ureteral stones smaller than 3 cm. The study demonstrated several advantages of FANS-UAS, including reduced postoperative fever, higher stone-free rates, and improved quality of life outcomes. Chen et al.³ recently published a study comparing traditional UAS with flexible suction UAS in RIRS for kidney stones ranging from 2 to 4 cm. In this retrospective study, which included 238 patients, a statistically significant difference was observed in favor of the flexible suction UAS group regarding operative time, as well as the stone-free rates on the first and thirtieth postoperative days. In a recent retrospective study by Erkoç et al.¹¹, patients with 2 cm or larger kidney stones were divided into two groups: aspiration-assisted UAS and traditional UAS. The comparison revealed a statistically significant advantage for the aspiration-assisted UAS group regarding stone-free rate, operative time, and postoperative fever incidence.

In a retrospective study by Chen et al.⁶ comparing mini PCNL (m-PCNL) and RIRS using FANS-UAS for kidney stones measuring 2 to 3 cm, RIRS was found to have advantages such as a shorter hospital stay and less postoperative hemoglobin decrease. Additionally, fewer postoperative complications were observed with RIRS, suggesting that it could be a superior treatment option to m-PCNL for managing stones in this size range. In this study, we compared RIRS and PCNL performed using FANS-UAS for the surgical management of 2–3 cm kidney stones, focusing on efficacy and safety. Statistically significant advantages were observed in the RIRS group regarding hospital stay, operative time, and postoperative hemoglobin decrease. However, stone clearance success and postoperative complication rates were statistically similar between the groups. Notably,

Clavien-Dindo grade III complications were reported in three patients in the PCNL group: one required an interventional radiology procedure for an arteriovenous fistula, and two underwent Double-J stent placement due to postoperative urine leakage. No grade III complications were observed in the RIRS group.

Several limitations should be acknowledged in this study. First, the retrospective design and limited sample size may affect the generalizability of our findings. Second, two RIRS cases (4% of the initial cohort) were excluded from the final analysis due to incomplete follow-up data after being referred to other centers. Third, the use of KUB radiography on postoperative day 1 (for radio-opaque stones) may have underestimated small residual fragments, though this was mitigated by standardized low-dose CT at 3-month follow-up. Fourth, the involvement of multiple surgeons, despite all being high-volume endourologists, could introduce technical variability. Finally, the non-randomized design carries inherent selection bias, though we mitigated this through standardized patient counseling and multidisciplinary case selection.

In conclusion, according to this study, RIRS using FANS-UAS has emerged as a promising alternative for the surgical treatment of kidney stones measuring 2–3 cm. With acceptable SFR, shorter hospitalization, and reduced duration of surgery, RIRS provides both surgeons and patients with a less invasive option for managing large kidney stones, yielding favorable outcomes. Prospective randomized studies are needed to support the findings.

DRAFT

REFERENCES

1. Soderberg L, Ergun O, Ding M, et al. Percutaneous nephrolithotomy vs retrograde intrarenal surgery for renal stones: A Cochrane review. *BJU Int* 2024;133:132-40. <https://doi.org/10.1111/bju.16220>
2. Karakoyunlu N, Goktug G, Şener NC, et al. A comparison of standard PCNL and staged retrograde FURS in pelvis stones over 2 cm in diameter: A prospective randomized study. *Urolithiasis* 2015;43:283-7. <https://doi.org/10.1007/s00240-015-0768-2>
3. Chen H, Xiao J, Ge J, Liu T. Clinical efficacy analysis of tip-flexible suctioning ureteral access sheath combined with disposable flexible ureteroscope to treat 2-4 cm renal stones. *Int Urol Nephrol* 2024;56(10):3193-9. <https://doi.org/10.1007/s11255-024-04072-y>
4. Dutta R, Mithal P, Klein I, Patel M, Gutierrez-Aceves J. Outcomes and costs following mini-percutaneous nephrolithotomy or flexible ureteroscopic lithotripsy for 1-2-cm renal stones: Data from a prospective, randomized clinical trial. *J Urol* 2023;209:1151-8. <https://doi.org/10.1097/JU.0000000000003397>
5. Soderberg L, Ergun O, Ding M, et al. Percutaneous nephrolithotomy versus retrograde intrarenal surgery for treatment of renal stones in adults. *Cochrane Database Syst Rev* 2023;11:CD013445. <https://doi.org/10.1002/14651858.CD013445.pub2>
6. Chen Y, Xi H, Yu Y, et al. Flexible ureteroscopy with novel flexible ureteral access sheath versus mini-percutaneous nephrolithotomy for treatment of 2-3 cm renal stones. *Int J Urol* 2024;31:281-6. <https://doi.org/10.1111/iju.15347>
7. Karakeçi A, Karakoç O, Ozan T, et al. Comparison of retrograde intrarenal surgery and percutaneous nephrolithotomy for the treatment of renal stones greater than 2 cm. *Turk Urol Derg* 2015;41:73-7. <https://doi.org/10.5152/tud.2015.97957>
8. Cheng Y, Xu R. Effectiveness and safety of retrograde intrarenal surgery (RIRS) vs percutaneous nephrolithotomy (PCNL) in the treatment of isolated kidney stones. *Am J Transl Res* 2022;14:1849-58.
9. Constantinou BT, Benedicto BC, Porto BC, et al. PCNL vs two staged RIRS for kidney stones greater than 20 mm: Systematic review, meta-analysis, and trial sequential analysis. *Minerva Urol Nefrol* 2024;76:31-41. <https://doi.org/10.23736/S2724-6051.23.05577-5>
10. Akman T, Binbay M, Ozgor F, et al. Comparison of percutaneous nephrolithotomy and retrograde flexible nephrolithotripsy for the management of 2-4 cm stones: A matched-pair analysis. *BJU Int* 2012;109:1384-9. <https://doi.org/10.1111/j.1464-410X.2011.10691.x>
11. Erkoc M, Bozkurt M, Sezgin MA, et al. Efficacy of aspiration-assisted ureteral access sheath (ClearPETRA) in retrograde intrarenal surgery. *J Laparoendosc Adv Surg Tech* 2024;34:420-4. <https://doi.org/10.1089/lap.2024.0076>
12. Fayad MK, Fahmy O, Abulazayem KM, Salama NM. Retrograde intrarenal surgery versus percutaneous nephrolithotomy for treatment of renal pelvic stone more than 2 centimeters: A prospective randomized controlled trial. *Urolithiasis* 2022;50:113-7. <https://doi.org/10.1007/s00240-021-01289-9>
13. Chen Y, Liao B, Feng S, et al. Comparison of safety and efficacy in preventing postoperative infectious complications of a 14/16F ureteral access sheath with a 12/14F ureteral access sheath in flexible ureteroscopic lithotripsy. *J Endourol* 2018;32(10):923-7. <https://doi.org/10.1089/end.2018.0222>

14. Turedi B, Sezer A. Comparison of flexible and navigable suction ureteral access sheath with conventional ureteral access sheath for pediatric retrograde intrarenal surgery: A single-center propensity-matched analysis. *Urolithiasis* 2024;53:17. <https://doi.org/10.1007/s00240-024-01686-w>
15. Gauhar V, Madarriaga YQ, Somani B, et al. Is flexible and navigable suction ureteral access sheath (FANS-UAS) the next best development for retrograde intrarenal surgery in children? Results of a prospective multicentre study. *World J Urol* 2024;42:627. <https://doi.org/10.1007/s00345-024-05337-y>
16. Zhu W, Liu S, Cao J, et al. Tip bendable suction ureteral access sheath versus traditional sheath in retrograde intrarenal stone surgery: An international multicentre, randomized, parallel group, superiority study. *eClinicalMedicine* 2024;74:102724. <https://doi.org/10.1016/j.eclinm.2024.102724>

DRAFT

FIGURES AND TABLES

| Table 1. Patient demographic data | | | |
|--|-------------|-------------|----------|
| | RIRS | PCNL | p |
| n | 50 | 66 | |
| Male/female | 32/18 | 43/23 | 0.898 |
| Age(y)* | 47.8±7.9 | 46.2±6.1 | 0.245 |
| BMI(kg/m ²)* | 27.5±5.2 | 26.9±3.1 | 0.710 |
| Stone size (mm)* | 24.5±1.8 | 24.8±1.4 | 0.297 |
| Hounsfield units | 980.7±137.5 | 1011.2±89.3 | 0.177 |
| Stone locations** | | | 0.792 |
| Lower | 18 | 18 | |
| Middle | 3 | 7 | |
| Upper | 7 | 8 | |
| Pelvic | 10 | 15 | |
| Multicalyceal | 12 | 18 | |

*Mean ± standard deviation. **The pattern of stone presence in the kidney. PCNL: percutaneous nephrolithotomy; RIRS: retrograde intrarenal surgery.

| Table 2. Postoperative outcome analysis | | | |
|--|-------------|--------------|----------|
| | RIRS | PCNL | p |
| Hospitalization* (h) | 28.3 ± 17.9 | 81 ± 24.8 | <0.001 |
| Operative time* (min) | 63.1 ± 11.3 | 97.3 ± 15.1 | <0.001 |
| Hemoglobin drop* (g/dL) | 0.4 ± 0.5 | 2.1 ± 1.7 | <0.001 |
| Stone-free rate**, n (%) | 38/50(%76) | 54/66(%81.8) | 0.492 |
| Success rate***, n(%) | 35/50(%70) | 49/66(%74.2) | 0.677 |
| Post-op complications | | | 0.277 |
| Clavien grade I | 4 | 8 | |
| Clavien grade II | 2 | 6 | |
| Clavien grade III | 0 | 3 | |
| Clavien grade IV-V | 0 | 0 | |

*Mean ± standard deviation. **Outcomes observed on day 1 after surgery. *** Complete stone clearance (Grade A, defined as no fragments ≥1 mm) at the 3-month postoperative followup. PCNL: percutaneous nephrolithotomy; RIRS: retrograde intrarenal surgery.