

APPENDIX

Supplementary Table 1. Additional perioperative outcomes by type of VTE prophylaxis received at discharge			
Variable		Enoxaparin (n=102)	Apixaban (n=83)
Intra-op transfusion, n (%)	PRBC	5 (4.9)	5 (6.0)
	PRBC and albumin	8 (7.8)	8 (9.6)
	Albumin	11 (10.8)	9 (10.8)
Postop transfusion, n (%)	PRBC	22 (21.6)	10 (12.0)
	PRBC and albumin	10 (9.8)	7 (8.4)
	Albumin	18 (17.6)	10 (12.0)
Median (IQR) pre-op Hb		129.5 (115.0, 142.8)	124.0 (113.5, 138.5)
Median (IQR) POD1 Hb		93.0 (84.0, 107.0)	96.0 (86.5, 106.0)
Median (IQR) Hb at discharge		98.0 (85.5, 110.8)	100.0 (90.0, 109.5)
Clavien-Dindo pre-discharge grade, n (%)	1, 2	52 (51.0)	42 (50.6)
	3, 4	11 (10.8)	3 (3.6)
Median (IQR) LOS, days		7.0 (6.0, 7.0)	6.0 (5.0, 7.0)

Hb: hemoglobin; IQR: interquartile range; LOS: length of stay; PRBC: packed red blood cells.