

Case - Hematuria and uroepithelial thickening: Anticoagulation-induced suburothelial hemorrhageSepehr Niakani¹, Adel Arezki², Giovanni Artho³, Alexis Rompré-Brodeur²¹Faculty of Medicine and Health Sciences, McGill University, Montreal, QC, Canada; ²Division of Urology, Department of Surgery, McGill University Health Center, Montreal, QC, Canada; ³Division of Abdominal Radiology, Department of Radiology, McGill University Health Center, Montreal, QC, Canada**Cite as:** Niakani S, Arezki A, Artho G, et al. Case - Hematuria and uroepithelial thickening: Anticoagulation-induced suburothelial hemorrhage. *Can Urol Assoc J* 2026 March 16, 2026; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.9319>

Published online March 16, 2026

Corresponding author: Sepehr Niakani, Faculty of Medicine and Health Sciences, McGill University, Montreal, QC, Canada; sepehr.niakani@mail.mcgill.ca**INTRODUCTION**

Antopol and Goldman discovered lesions mimicking renal neoplasms in seven patients who had undergone nephrectomies as a result, and subsequently named them Antopol-Goldman Lesion (AGL).¹

AGL is a subepithelial hemorrhage typically affecting the renal pelvis^{1,2}, and usually presents as hematuria and flank pain, most commonly in the elderly.¹⁻⁶ Since AGL masquerades as renal neoplasms, it was mainly diagnosed histologically from specimens collected after nephrectomies.³ With the advent of advanced imaging modalities and diagnostic tools such as flexible ureteroscopy, nephrectomies are avoided, and patients are treated conservatively.^{3,5,7,8}

In this report, we present the case of a 52-year-old female with AGL occurring in the context of supratherapeutic international normalized ratio (INR) level secondary to warfarin therapy. We demonstrate that conservative management by temporary cessation of warfarin and optimizing anticoagulant dosing led to the full resolution of symptoms.

KEY MESSAGES

- Supratherapeutic warfarin therapy is a risk factor for Antopol-Goldman lesion occurrence.
- Diagnosis relies on imaging, typically CT urogram showing suburothelial hemorrhage.
- Conservative management is effective in stable patients without concerning features.
- Follow-up may include cystoscopy, repeat imaging, and INR monitoring based on individual risk.

CASE REPORT

A 52-year-old female presented to the Emergency Department (ED) with painless gross hematuria persisting for three days, accompanied by mild left lower quadrant (LLQ) pain radiating from the left flank. Her medical history was significant for hypertension and a mechanical mitral valve replacement performed two months prior for severe mitral regurgitation. She was maintained on warfarin for anticoagulation, alongside metoprolol and aspirin.

Two days prior to the ED presentation, she attended a routine hematology clinic visit, where her INR was found to be 4.6 (therapeutic target for mechanical mitral valves: 3.0–3.5⁹). In response, the hematologist advised holding her next warfarin dose. However, by that time, she had already been experiencing gross hematuria for one day. Despite the temporary discontinuation of warfarin, her hematuria persisted and was followed by worse LLQ pain, prompting her presentation to the ED the following day.

On initial assessment, her vital signs were stable. Physical examination was unremarkable except for mild LLQ tenderness without costovertebral angle tenderness or peritoneal signs. Given the ongoing hematuria, warfarin was temporarily withheld, while her other medications were continued. Laboratory investigations including a complete blood count, coagulation profile, comprehensive metabolic panel, liver function tests, urinalysis, and urine culture were performed. Pertinent findings included an INR of 4.09, which had decreased from 4.6 but remained above the therapeutic range, numerous red blood cells on urinalysis, and the presence of leukocyte esterase. The urine culture was negative for bacterial growth, and hemoglobin remained stable at 122 g/L (reference range: 120–160 g/L). A multiphasic computed tomography (CT) urogram was performed on the day of admission (approximately three days after symptom onset) revealing non-enhancing hyperdense suburothelial collections (Hounsfield unit: 25) extending from the left renal collecting system to the mid-ureter, with associated left pelviectasis (Fig. 1A,B). The radiologist interpreted these findings as consistent with suburothelial hemorrhage, raising suspicion for an AGL attributed to supratherapeutic anticoagulation. Notably, a contrast-enhanced CT scan of the abdomen and pelvis from one month prior had shown no abnormalities in the kidneys or ureters.

The patient remained hemodynamically stable throughout her hospitalization. Withholding one additional dose of warfarin resulted in the resolution of hematuria and LLQ pain. She was advised to continue holding warfarin for two additional days following discharge before resuming at a reduced dose of 5 mg daily (previously 7 mg on Fridays and 5 mg on other days). Invasive diagnostic procedures such as ureteroscopy or retrograde pyelography were not pursued given the patient's hemodynamic stability, absence of hydronephrosis or suspicious enhancing masses, and the radiologist's diagnostic impression. Outpatient follow-up was arranged in urology for cystoscopic evaluation of the lower urinary tract along with urine cytology as well as a follow-up CT scan to ensure resolution of findings. Additionally, she was scheduled for hematology follow-up for warfarin titration.

At her most recent follow-up, her symptoms had completely resolved, she remained asymptomatic, and her INR had stabilized at 2.57. The cystoscopic evaluation did not reveal any lower urinary tract pathologies, and the urine cytology was normal. The follow-up CT scan also showed complete resolution of the suburothelial hemorrhage (Fig. 1C,D).

DISCUSSION

Subepithelial hemorrhage of the urothelium, also known as AGL¹, is a rare condition with less than 60 cases reported in the literature. It classically affects the renal pelvis^{1,2} and presents as flank pain and hematuria mostly in the elderly.¹⁻⁶ Previously, AGL was diagnosed histologically post partial or radical nephrectomies due to its radiographic resemblance to renal neoplasms (e.g., evidence of filling defects on CT imaging).⁷ Today, improvements in imaging modalities and increased awareness of AGL have largely eradicated unnecessary nephrectomies, and AGL is mainly diagnosed radiologically.^{3,10} Cardin et al., demonstrated for the first time that AGL can be managed conservatively with follow-ups in patients with coagulopathies or anticoagulant use and unremarkable ureterorenoscopic examinations.⁵ Similarly, Gayer and colleagues conservatively managed AGL in seven patients by correcting their coagulopathies or discontinuing anticoagulants leading to full symptom resolution.¹⁰ Despite the modern adoption of such conservative approaches, complications such as refractory hematuria requiring selective arterial embolization¹⁰ and emphysematous pyelonephritis requiring urgent nephrectomy⁴ should not be missed. Moreover, although the etiology of AGL remains poorly understood, it has been associated with trauma, hypertension, diabetes, amyloidosis, and coagulopathies.^{3,5,6} Of note, anticoagulant use is a common feature of several reported cases of AGL.^{6,7,12-15}

In our report, we present the case of a 52-year-old female patient who presented to the ED complaining of left flank pain and hematuria, who was on warfarin therapy due to a prior mitral valve replacement. Laboratory investigation indicated an elevated INR (4.09), which is higher than the acceptable upper limit of 3.5 for mechanical mitral valves.⁹ A CT urogram showed suburothelial hemorrhage extending from the left renal collecting system to the proximal ureter. The extension of the hemorrhage to the level of the proximal ureter is an interesting and uncommon finding in our case as AGL typically affects the renal pelvis.¹ Nevertheless, our findings are consistent with previous reports where AGL was attributed to anticoagulant use and/or elevated INR levels.^{7,12-15} Importantly, the resolution of hematuria with cessation of warfarin further reinforces the utility of conservative management, especially in patients on anticoagulant therapy. While there are no clear follow-up guidelines or recommendations, cystoscopy of the lower urinary tract, outpatient assessment of underlying factors such as coagulopathies or hypertension, or CT imaging can be considered. However, with the resolution of symptoms and the absence of worrisome malignant features, follow-ups may be of little value.

CONCLUSIONS

With less than 60 cases reported to date, suburothelial hemorrhage is an uncommon presentation associated with certain risk factors. Here, we present the case of a suburothelial hemorrhage

secondary to supratherapeutic warfarin dosing and show that its discontinuation leads to full symptom resolution. Our report further demonstrates that AGL can be managed conservatively by correcting underlying contributing factors, such as an unoptimized anticoagulant dosage. Further follow-up with cystoscopy, repeat CT imaging, and outpatient assessment of risk factors can be considered depending on the patient profile and the presence of malignant features.

DRAFT

REFERENCES

1. Antopol W, Goldman L. Subepithelial hemorrhage of renal pelvis simulating neoplasm. *Urol Cutaneous Rev* 1948;52:189-95.
2. Zelaya R, Byerly D, Zarka A. Spontaneous suburothelial hemorrhage associated with forniceal rupture and retroperitoneal hemorrhage. *Cureus* 2022;14:e23728. <https://doi.org/10.7759/cureus.23728>
3. Şeker KG, Arıcan G, Başar MM, et al. A rare case mimicking collecting system tumor: Antopol-Goldman lesion. *Bull Urooncol* 2018;17:150-2. <https://doi.org/10.4274/uob.1052>
4. Komagamine J, Kano Y. Renal pelvic haemorrhage complicating emphysematous pyelonephritis. *Eur J Case Rep Intern Med* 2023. https://doi.org/10.12890/2023_004241
5. Cardin AL, Daneman A, Navarro OM, et al. Antopol-Goldman lesion of the kidney diagnosed by radiology: A case report of observation. *Acta Radiol* 2008;49:715-7. <https://doi.org/10.1080/02841850802056009>
6. Eccher A, Gobbo S, Grassi A, et al. Subepithelial pelvic hematoma (Antopol-Goldman lesion) simulating renal neoplasm: Report of a case and review of the literature. *Int J Surg Pathol* 2009;17:264-7. <https://doi.org/10.1177/1066896908330482>
7. Sahin TK, Uysal F, Ozturk A, et al. Spontaneous subepithelial hemorrhage of renal pelvis and ureter (Antopol-Goldman lesion) in hemophilia A patient with inhibitor. *Medicine* 2020;99:e18995. <https://doi.org/10.1097/MD.00000000000020851>
8. Altay B, Barisik CC, Erkurt B, et al. Subepithelial pelvic hematoma of the kidney (Antopol-Goldman lesion). *Turk J Urol* 2015;41:48-50. <https://doi.org/10.5152/tud.2014.48208>
9. Whitlock RP, Sun JC, Femes SE, et al. Antithrombotic and thrombolytic therapy for valvular disease. *Chest* 2012;141:e576S-600S. <https://doi.org/10.1378/chest.11-2305>
10. Gayer G, Zissin R, Apter S, et al. Spontaneous suburothelial hemorrhage in coagulopathic patients: CT diagnosis. *AJR Am J Roentgenol* 2011;197:W524-8. <https://doi.org/10.2214/AJR.11.6474>
11. Branchereau J, Raynaud JP, Justrabo E, et al. Syndrome d'Antopol-Goldman bilatéral associé à un déficit congénital sévère en facteur V. *Prog Urol* 2010;20:604-7. <https://doi.org/10.1016/j.purol.2009.12.002>
12. Fishman M, Pollack H, Arger P, et al. Radiographic manifestations of spontaneous renal sinus hemorrhage. *AJR Am J Roentgenol* 1984;142:1161-4. <https://doi.org/10.2214/ajr.142.6.1161>
13. Viamonte M, Viamonte M, Rywlin AM, et al. Subepithelial hemorrhage of renal pelvis simulating neoplasm (Antopol-Goldman lesion). *Urology* 1980;16:647-9. [https://doi.org/10.1016/0090-4295\(80\)90581-6](https://doi.org/10.1016/0090-4295(80)90581-6)
14. Miller V, Witten DM, Shin MS. Computed tomographic findings in suburothelial hemorrhage. *Urol Radiol* 1982;4:11-4. <https://doi.org/10.1007/BF02924017>
15. Kossol JM, Patel SK. Suburothelial hemorrhage. *J Comput Assist Tomogr* 1986;10:157-<https://doi.org/10.1097/00004728-198601000-00038>

FIGURES AND TABLES

Figure 1. Multiphase computed tomography urogram showing a left-sided Antopol Goldman lesion and its interval resolution. The axial non-contrast and coronal delayed excretory-phased images demonstrate hyperdense suburothelial collections along the left renal pelvis and proximal ureter as shown by the arrows (panels A&B), which showed complete resolution on followup imaging (panels C&D).



