

A rapid evidence assessment for extracorporeal magnetic stimulation to treat urinary incontinence in menRyan Wong¹, Blayne Welk²¹ Schulich Medical School, Western University, London, ON, Canada; ² Department of Surgery and Epidemiology and Biostatistics, Western University, London, ON, Canada**Cite as:** Wong R, Blayne W. A rapid evidence assessment for extracorporeal magnetic stimulation to treat urinary incontinence in men. *Can Urol Assoc J* 2025 October 27; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.9320>

Published online October 27, 2025

Corresponding author: Dr. Blayne Welk, Department of Surgery and Epidemiology and Biostatistics, Western University, London, ON, Canada; bkwelk@gmail.com

ABSTRACT**Introduction:** Extracorporeal magnetic stimulation (EMS) is a non-invasive treatment for urinary incontinence (UI) in women, but its effectiveness in men is not well known. This review identifies and evaluates the evidence supporting EMS for treating UI in adult men.**Methods:** We systematically searched the MEDLINE, CINAHL, and PEDro databases up to November 2024. Studies included EMS alone or compared to other treatments in men with UI. A qualitative assessment of the evidence was carried out. Study quality

was assessed using the Downs and Black checklist for randomized and non-randomized studies and the Cochrane Risk of Bias 2 tool for randomized controlled trials (RCTs).

Results: Of 285 studies screened, nine met the inclusion criteria, encompassing 181 men treated with EMS, mostly post-prostatectomy. Four RCTs, with qualities ranging from fair to good, and small sample sizes (n=16–36), found EMS led to earlier continence compared to pelvic floor muscle therapy (PFMT) and was superior to sham treatment. One study reported significant improvements in urodynamic measures after EMS, and another showed a 48% reduction in 24-hour pad usage, sustained at 12.5 months. Several studies indicated that EMS accelerated symptom improvement compared to PFMT, but had similar long-term outcomes. All studies using validated quality-of-life measures reported significant improvements after EMS.**KEY MESSAGES**

- Across nine studies (181 men treated with EMS), EMS led to earlier continence recovery and was superior to sham treatment; one study showed a 48% reduction in 24-hour pad usage sustained for over a year.
- Evidence quality is limited by small sample sizes, methodologic concerns, and lack of standardized protocols; most studies were rated as fair, with some risk of bias
- Larger, high-quality studies are needed to confirm the benefits of EMS and to better define its role in managing male urinary incontinence.

Conclusions: Evidence for EMS in treating male UI is limited but generally positive. EMS may promote faster continence recovery than PFMT, with similar long-term outcomes. Larger, high-quality studies are needed to confirm these findings and guide clinical practice and recommendations for different subgroups.

INTRODUCTION

Incontinence is a common and impactful complication after prostate cancer treatment. Approximately 8 to 25% of patients will experience persistent post-prostatectomy incontinence at 12 months postoperatively,¹ and this is a major contributor to decreased general health-related quality of life.² Current AUA/SUFU guidelines for post-prostatectomy incontinence suggest the use of conservative options such as pelvic floor muscle therapy (PFMT) and surgical interventions such as implantation of a male sling or an artificial urinary sphincter (AUS).^{3,4} These options have known limitations. For instance, traditional PFMT often has poor patient compliance,⁵ and there are complications associated with the AUS such as infection, mechanical failure, erosion, and persistent leakage.⁴

Electromagnetic stimulation (EMS) was first described by Galloway et al., in 1998 as a non-invasive alternative to functional electrical stimulation (FES).⁶ During EMS treatment, patients sit in a special chair equipped with an electromagnetic coil located in the seat. The activated coil stimulates contractions in pelvic floor muscles, and in some dual-coil chairs, depolarizations in motor nerves in the sacral root. These magnetic currents can penetrate through the body tissue without significant alteration, allowing a patient to be treated in a non-invasive manner.^{6,7} This was a major improvement over FES which required the insertion of a transrectal probe to stimulate deeper structures in the pelvis.⁸ EMS has been shown to be safe and effective for incontinence in women and treats both overactive bladder symptoms and stress incontinence.⁹ In this population it is a non-invasive, well tolerated treatment option with very rare potential adverse effects and limited contraindications.^{9,10}

In a 2015 systematic review of conservative management for postprostatectomy incontinence, Anderson et al., concluded that while EMS be beneficial in recovery, the evidence was still limited.¹¹ In this rapid literature review, we identified evidence for the use of EMS in men with urinary incontinence and identified the therapeutic benefits, gaps in current knowledge, and areas for future research.

METHODS

This rapid literature review was performed according to the structure of rapid reviews outlined by the World Health Organization,¹² and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines¹³.

Identification of studies

We searched MEDLINE, CINAHL, and PEDro from inception to November 2024. Literature search strategies were developed with the expertise of a reference librarian; search terms and the PICO statement are included in the Appendix. The reference lists of the identified studies were also checked to find additional articles relevant to our research question. All references were

imported into Covidence and duplicates were removed. Studies were then screened in duplicate by two independent reviewers (R.W. and B.W.).

We included all original research articles evaluating the outcomes of men being treated with EMS for urinary incontinence. We also included studies that compared EMS with other methods of pelvic floor rehabilitation such as PFMT, or FES. We excluded abstracts, editorials, commentaries, and other non-experimental studies. We excluded studies that had male and female participants without a subgroup analysis for only male participants. Studies screened at the title and abstract stage were excluded if they did not meet one or more of these inclusion criteria. Rationales for exclusions were recorded at the full-text screening stage.

Data extraction and quality assessment

Data was extracted by two authors independently (R.W., B.W.) and included primary author, year of article publication, study design, inclusion and exclusion criteria for study sample, number of patients analyzed, type of pelvic floor therapies, frequency of treatments and outcome measures. All studies were assessed for methodological quality using the Downs and Black scale consisting of 27 questions that assessed: reporting, external validity, internal validity/bias, and internal validity/confounders.¹⁴ Based on scoring on the Downs and Black scale, studies were classified as excellent, good, fair, or poor¹⁵ (Table 1). Randomized controlled trials included in this review were evaluated using the Cochrane Risk of Bias tool (RoB2) that assesses the quality of studies on the basis of methods of randomization, deviations from intended intervention, missing outcome data, measurement of outcomes, and reporting of results.^{16,17}

Analysis

Due to the limited studies, and the heterogeneity in treatment protocols, EMS device, and patient populations, a qualitative analysis of the results was performed, and no meta-analysis was carried out.

RESULTS

We identified 285 studies in the initial search of three databases. After removing duplicates, and applying our inclusion criteria, 31 were studies eligible for full-text screening. After detailed review, and snowball searches of other systematic reviews, we had 9 studies included in the present review (Figure 1).

Study details

Out of nine studies included in this review, four were RCTs,^{18–21} four were cohort studies,^{22–25} and one was a quasi-experimental study²⁶. All studies reported data at baseline and after interventions. Table 2 summarizes the studies, their design, and their objective. Seven of the nine studies were specifically in men after radical prostatectomy. Three studies originated from Japan, two from Italy, and one each from Germany, Korea, China, and Taiwan.

The most common comparison groups were FES, or PFMT. Out of the studies included, 3 studies examined the effect of EMS alone,^{22,23,25} two compared the effects of EMS with PFMT,^{20,21} and one compared a group of patients that received no treatment to those who received EMS and PFMT.²⁶ One study compared EMS to FES,²⁴ and another compared FES, EMS and no treatment.¹⁹ Lastly, one study employed a crossover design where patients were either assigned a sham group first followed by EMS treatment, or vice versa.¹⁸

Participants

There was a total of 390 male participants studied, of which 181 were treated with EMS (Table 3). The number of participants in each study ranged from 10 to 178. Of the RCTs included, total sample sizes ranged from 16 to 36. In the one study by Suzuki et al. that did not specifically recruit men with postprostatectomy incontinence, all participants were diagnosed with idiopathic urgency incontinence based on urodynamic studies, and 16 of 39 participants were male.¹⁸ Three studies included participants with stress (SUI), urgency (UUI) and mixed incontinence (MUI).^{22,25,26} Yokoyama et al., included only those with SUI,²⁴ and Terzoni et al., included those with SUI and UUI.²³ Three studies did not specify the type of incontinence being treated.^{19–21} In the seven studies that reported baseline incontinence measures, there was an average of 3.85 episodes of incontinence per day,^{18,23,25} 24 hour pad weights of 477.7g,^{19,20,24} and use of 5.4 pads per day.²² No studies excluded patients based incontinence severity.

Intervention

Participants who received EMS had treatment spanning between 4 and 10 weeks, with 2 to 3 sessions per week that each lasted 10 to 20 minutes. To administer EMS, three studies used the NeoControl® therapy system,^{19,22,23} three used the BioCon 2000W™,^{20,21} one used a magnetic stimulator from Nihon Kohden,¹⁸ and two did not specify the device used. Each study had its own treatment protocol, which generally consisted of multiple short treatments over the course of 4-10 weeks. An estimate of the total number of treatment contractions was calculated for each study, with most protocols delivering approximately 500,000 contractions.

Outcomes

The studies used various diagnostic tests, such as pad weights, pad usage, bladder diaries, and urodynamic studies (UDS). All studies except for two also evaluated quality of life (QoL) with validated and reliable questionnaires such as the Urinary Incontinence Quality of Life Scale (I-QoL), the Urogenital Distress Inventory (UDI-6), the International Consultation on Incontinence Questionnaire-Short Form (ICIQ-SF), the International Prostate Symptom Score (IPSS), Incontinence Impact Questionnaire (IIQ-7).^{18–21,23,25,26}

There were two small cohort studies that included urodynamic studies; one found that EMS significantly increased mean maximum cystometric capacity and Valsalva leak point pressure²³, and in the other, functional bladder capacity.²⁵ Two small studies (a cohort study and a cross-over study) found that EMS decreased the number of episodes of incontinence by 35-36% by the conclusion of treatment.^{18,25} In one of the larger cohort studies, EMS reduced 24-hour pad usage by 48% with persistent reductions at 12.5 months without maintenance therapy in men who had undergone radical prostatectomy.²² However, other authors reported less durable success, with Yokoyama et al., finding that 3 of 6 patients that had significant improvements in their I-QoL scores had returned to baseline within 12 months.²³ Two studies showed that EMS improved lower urinary tract symptoms (LUTS) and urinary leakage quicker than pelvic floor muscle therapy, however at the end of the studies both groups reached similar levels of improvement.^{19,26} Terzoni et al. also found that EMS resulted in quicker reductions in urine leakage compared to FES²⁴. All seven studies that used validated QoL questionnaires found that the use of EMS significantly improved QoL measures compared to baseline.^{18–21,23,25,26} However, when compared to the PFMT in one study, there was no difference in the improvement of QoL measures²⁶.

Critical appraisal

All studies were assessed using the Downs and Black checklist, which allowed the quality assessment of comparative and non-comparative studies (Table 4). The mean score of all studies in this review was 17.6/27, with non-RCTs averaging 16.6/27, and RCTs averaging a score of 18.8/27. Seven studies were assessed as being methodologically fair,^{19–21,23–26} and one was methodologically poor.²² The one study assessed as methodologically good was a small cross-over RCT of EMS and sham; it had a critical score of 23/27 due to the use of blinding, randomization and a sham comparator.¹⁸ The study that was assessed as methodologically poor was a cohort study and scored 14/27.²² Notably, none of the studies reported power calculations to indicate that they had a sufficient sample size to detect a clinically significant result. Moreover, none of the studies stated the proportion of patients asked who agreed to participate in each study (and there was no flow diagram of participants through the various phases of a trial, as recommended by the CONSORT statement for RCTs).²⁷ RCTs were also assessed using the Cochrane Risk of Bias 2 (RoB2). Two of four RCTs included in this review showed an overall “high” risk of bias, and 2 showed “some concern”. None of the RCTs were enrolled into a clinical trial registry, and therefore it was impossible to ascertain if studies were analyzed according to a pre-specified plan. As such, all studies were scored as “some concern” in the domain of “Selection of Reported Results.” Visual representations of risk of bias assessments are shown in Figure 2.

DISCUSSION

There is currently limited literature exploring the impact of EMS in the treatment of incontinence in men. To our knowledge, this review represents the first review evaluating the therapeutic benefits of EMS for treating incontinence in men. Much of the foundational research in EMS has focused on women. The more recent literature on EMS therapy in women has reported a consistent benefit; for example a recent systematic review concluded that EMS was effective at treating incontinence and improving patients’ QoL;⁹ the authors identified 11 studies published over the last 15 years (four with a strong quality score), and the majority of these studies demonstrated a benefit with EMS for women with incontinence. These results are similar to another recent systematic review of EMS for women.¹⁰ This aligns with our key finding that in men, EMS appears to improve QoL and may be effective at treating incontinence. However, our conclusions should be tempered by the quality of the literature identified: we only identified four RCTs, representing 108 patients, and all had at least some concern of bias. Additionally, the studies used a variety of treatment protocols and devices; most of the devices were not “high-frequency” devices that use a higher-intensity magnetic field and produce supramaximal contractions, as seen with most contemporary devices.

The studies in this review showed that in general, men treated with EMS have improvements in incontinence episodes, lower urinary tract symptoms, and QoL measures, and in some urodynamic parameters. However, it is still uncertain as to how long these improvements last due to conflicting study results: one small study showed that some patients returned to baseline I-QoL scores within 12 months,²³ while another study showed lasting reductions in pad usage at 12.5 months of follow-up.²² This speaks to the lack of long-term follow-up for the majority of these studies (with most limited to less than 6 months). It is also uncertain if EMS is more effective than PFMT. Authors have both shown an advantage with EMS versus PFMT in QOL,²¹ and similar outcomes at 6 months (with a quicker initial recovery

with EMS).^{20,26} An advantage over PFMT may be attributed to an adherence advantage for EMS in the real world due to the passive nature of the treatment. The identified studies used a variety of both qualitative evaluations, such as the I-QoL questionnaire, in addition to quantitative evaluations such as 24-hour pad weights or urodynamic studies. Therefore, it was challenging to compare the efficacy of EMS between studies that used different outcome measures of incontinence.

RCTs unsurprisingly scored higher on the Downs and Black checklist than non-RCTs. However, this was largely due to a single RCT that scored 23/27. The quality of the remaining RCTs was moderate, with studies rated between 16/27 to 19/27. These scores were impacted by a lack of blinding and limitations as to how data was reported across the studies. For instance, half of the RCTs in this review did not report on the number and characteristics of participants who were lost to follow-up. Reporting of these characteristics is important in understanding bias of the included results of a study. Among non-RCTs, the gaps in quality were mainly due to the lack of adverse event reporting. Moreover, no study reported power calculations, and several studies were missing key information such as how patients were recruited. As a result of these gaps in study design, the quality of most studies in this review limit the extrapolation of results to widespread clinical practise.

While this review did not use post-prostatectomy incontinence as part of the study inclusion criteria, seven of nine articles in this review studied only patients with post-prostatectomy incontinence. Therefore, the strongest support from the literature is specific this to patient population. This is appropriate, as RP is a common cause of incontinence in men. In the final stages of completion of this project, one newly published study was identified. The publication date precluded formal inclusion in our systematic review, however it is relevant to include a summary of the study. In a randomized, quadruple-blind, sham-controlled clinical trial, Unal *et al.*, found that EMS is superior to sham treatment at improving QoL and rates of incontinence episodes in those with post-prostatectomy incontinence.²⁸ They also found that EMS improved clinical parameters such as anxiety and sexual function. From a methodological perspective, this study included well described inclusion and exclusion of patients, description of participants lost to follow-up, adverse events, and both qualitative and quantitative assessments of incontinence. This study adds to the evidence supporting EMS for male incontinence, however it is still a small study (n=20 men/arm), and used a unique treatment protocol and device. Our study conclusions are not altered

This review had a few limitations. We identified a small number of studies, most of a moderate quality, that were conducted over a decade ago. The patient populations, intervention parameters, and outcomes were variable, and it was not possible to formally meta-analyse these study results. None of the studies had a sample size large enough to look at potential subgroups which may benefit, such as those with less severe incontinence, or those who were unable/unwilling to do PFMT. Importantly, to our knowledge, none of the studies used an EMS system that is commercially available in North America; this limits the generalisability of these previous studies.

CONCLUSIONS

Preliminary studies suggest that extracorporeal magnetic stimulation may be effective in treating men with urinary incontinence. Men may derive a benefit similar to pelvic floor muscle therapy, with improvements in the amount of urinary incontinence, and in incontinence related quality of

life. However, each of these studies have methodology limitations, and a risk of bias. Given the need for more low-risk treatment options for post-prostatectomy incontinence, larger, methodologically sound, appropriately controlled randomized trials are needed to evaluate the efficacy of EMS.

DRAFT

REFERENCES

1. Holm HV, Fosså SD, Hedlund H, et al. How should continence and incontinence after radical prostatectomy be evaluated? A prospective study of patient ratings and changes with time. *J Urol* 2014;192:1155-61. <https://doi.org/10.1016/j.juro.2014.03.113>
2. Penson DF, Feng Z, Kuniyuki A, et al. General Quality of life 2 years following treatment for prostate cancer: What influences outcomes? results from the Prostate Cancer Outcomes Study. *JCO* 2003;21:1147-4. <https://doi.org/10.1200/JCO.2003.07.139>
3. Sandhu JS, Breyer B, Comiter C, et al. Incontinence after prostate treatment: AUA/SUFU guideline. *J Urol* 2019;202:369–78. <https://doi.org/10.1097/JU.0000000000000314>
4. Breyer BN, Kim SK, Kirkby E, et al. Updates to incontinence after prostate treatment: AUA/GURS/SUFU guideline (2024). *J Uro* 2024;212:531-8. <https://doi.org/10.1097/JU.0000000000004088>
5. Wang L, Li Y, Qi Z, et al. Barriers and facilitators of the implementation of the application of pelvic floor muscle training in patients with prostate cancer: A scoping review. *Front Public Health* 2023;11:1191508. <https://doi.org/10.3389/fpubh.2023.1191508>
6. Galloway NTM, El-Galley RES, Sand PK, et al. Extracorporeal magnetic innervation therapy for stress urinary incontinence. *Urology* 1999;53:1108–11. [https://doi.org/10.1016/S0090-4295\(99\)00037-0](https://doi.org/10.1016/S0090-4295(99)00037-0)
7. Galloway NTM, El-Galley RES, Sand PK, et al. Update on extracorporeal magnetic innervation (ExMI) therapy for stress urinary incontinence. *Urology* 2000;56:82-6. [https://doi.org/10.1016/S0090-4295\(00\)00686-5](https://doi.org/10.1016/S0090-4295(00)00686-5)
8. Moore K. Treatment of urinary incontinence in men with electrical stimulation: Is practice evidence-based? *Journal of WOCN* 2000;27:20–31. [https://doi.org/10.1016/S1071-5754\(00\)90037-8](https://doi.org/10.1016/S1071-5754(00)90037-8)
9. Strojek K, Straczyńska A, Radzimińska A, et al. The effects of extracorporeal magnetic innervation in the treatment of women with urinary incontinence: A systematic review. *JCM* 2023;12:5455. <https://doi.org/10.3390/jcm12175455>
10. Leonardo K, Rahardjo HE, Afriansyah A. Non-invasive high-intensity focused electromagnetic therapy in women with urinary incontinence: A systematic review and meta-analysis. *NeuroUrol Urology* 2025;425:424-33. <https://doi.org/10.1002/nau.25658>
11. Anderson CA, Omar MI, Campbell SE, et al. Conservative management for postprostatectomy urinary incontinence. *Cochrane Database Syst Rev* 2015;1:CD001843. <https://doi.org/10.1002/14651858.CD001843.pub5>
12. Tricco AC, Langlois Etienne V, Straus SE, et al. Rapid reviews to strengthen health policy and systems: a practical guide. Geneva: World Health Organization. Available at: <https://iris.who.int/handle/10665/258698> (accessed May 12, 2025)
13. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. <https://doi.org/10.1136/bmj.n71>
14. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomized and non-randomized studies of healthcare interventions. *J Epidemiol Community Health* 1998;52:377-84. <https://doi.org/10.1136/jech.52.6.377>

15. Hooper P, Jutai JW, Strong G, et al. Age-related macular degeneration and low-vision rehabilitation: A systematic review. *Can J Ophthalmol* 2008;43:180-7. <https://doi.org/10.3129/i08-001>
16. Higgins JPT, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomized trials. *BMJ* 2011;343:d5928. <https://doi.org/10.1136/bmj.d5928>
17. Sterne JAC, Savović J, Page MJ, et al. RoB 2: A revised tool for assessing risk of bias in randomized trials. *BMJ* 2019;366:l4898. <https://doi.org/10.1136/bmj.l4898>
18. Suzuki T, Yasuda K, Yamanishi T, et al. Randomized, double-blind, sham-controlled evaluation of the effect of functional continuous magnetic stimulation in patients with urgency incontinence. *Neurourol Urodyn* 2007;26:767-72. <https://doi.org/10.1002/nau.20423>
19. Yokoyama T, Nishiguchi J, Watanabe T, et al. Comparative study of effects of extracorporeal magnetic innervation vs. electrical stimulation for urinary incontinence after radical prostatectomy. *Urology* 2003;63:264-7. <https://doi.org/10.1016/j.urology.2003.09.024>
20. Koo D, So SM, Lim JS. Effect of Extracorporeal magnetic innervation (ExMI) pelvic floor therapy on urinary incontinence after radical prostatectomy. *Korean J Urol* 2009;50:23. <https://doi.org/10.4111/kju.2009.50.1.23>
21. Liu F, Yao L, Mai H, et al. Extracorporeal magnetic innervation in the treatment of urinary incontinence after radical prostatectomy. *J Clin Rehab Tissue Eng Res*;12:3289-92. **[AU1: Can't find this reference. What year was it published?]**
22. Wöllner J, Neisius A, Hampel C, et al. "Extracorporeal magnetic innervation." *Urologe* 2012;51:1432-7. <https://doi.org/10.1007/s00120-012-2969-4>
23. Yokoyama T, Inoue M, Fujita O, et al. Preliminary Results of the effect of extracorporeal magnetic stimulation on urinary incontinence after radical prostatectomy: A pilot study. *Urol Int* 2005;74:224-8. <https://doi.org/10.1159/000083553>
24. Terzoni S, Montanari E, Mora C, et al. Electrical stimulation versus extracorporeal magnetic innervation for urinary incontinence after radical prostatectomy. *Int J of Uro Nursing* 2013;7:121-5. <https://doi.org/10.1111/ijun.12008>
25. Chang P-C, Wu C-T, Huang S-T, et al. Extracorporeal magnetic innervation increases functional bladder capacity and quality of life in patients with urinary incontinence after robotic-assisted radical prostatectomy. *Urol Sci* 2015;26:250-3. <https://doi.org/10.1016/j.urols.2015.06.286>
26. Terzoni S, Montanari E, Mora C, et al. Reducing urine leakage after radical retropubic prostatectomy: Pelvic Floor exercises, magnetic innervation or no treatment? A quasi-experimental study. *Rehab Nurs* 2013;38:153-60. <https://doi.org/10.1002/rnj.72>
27. Hopewell S, Chan A-W, Collins GS, et al. CONSORT 2025 statement: Updated guideline for reporting randomized trials. *BMJ* 2025;389:e081123. <https://doi.org/10.1136/bmj-2024-081123>
28. Unal B, Sarsan A, Yıldız N, et al. Efficacy of magnetic stimulation in men with urinary incontinence after radical prostatectomy: A randomized, quadruple-blind, sham-controlled clinical trial. *Neurourol Urodyn* 2025;44:1140-8. <https://doi.org/10.1002/nau.70055>

FIGURES AND TABLES

Figure 1. Process of study selection according to PRISMA flow diagram.




Figure 2. Visual representation of Cochrane risk of bias (RoB)2 assessment.

	D1	D2	D3	D4	D5	Overall
Suzuki et al., 2007	+	+	+	+	!	!
Koo et al., 2009	-	!	+	!	!	-
Yokoyama et al., 2004	!	-	-	-	!	-
Liu et al., 2009	!	+	+	!	!	!
Domains:					Judgement:	
D1: Bias arising from the randomization process					High risk	-
D2: Bias arising from deviations from intended interventions					Some Concern	!
D3: Bias due to missing outcome data					Low risk	+
D4: Bias in measurement of the outcome						
D5: Bias in selection of the reported result						

Quality index	Quality score (n=27)
Excellent	26–27
Good	20–25
Fair	15–19
Poor	≤14

Study	Country	Design	Type of incontinence	Study objective
Yokoyama et al, 2004 ¹⁹	Japan	RCT	Not specifically reported	To compare the efficacy of EMS vs FES on incontinence after retropubic RP
Yokoyama et al, 2005 ²³	Japan	Cohort	SUI, UII	To understand the clinical effects of EMS on incontinence after retropubic RP
Suzuki et al, 2007 ¹⁸	Japan	RCT	UII	To understand if there are sex differences in therapeutic outcomes when using EMS
Liu et al, 2008 ²¹	China	RCT	Not specifically reported	To compare EMS vs PFMT in post-RP incontinence
Koo et al, 2009 ²⁰	Korea	RCT	Not specifically reported	To compare the efficacy of EMS vs. PFMT on urinary incontinence after RP
Wöllner et al, 2012 ²²	Germany	Cohort	SUI, UII, MUI	To investigate the effectiveness and adverse events of EMS in the treatment of urinary incontinence
Terzoni et al, 2013 ²⁶	Italy	Quasi-experimental	SUI, UII, MUI	To compare the efficacy of PFMT and EMS to treat post-RP incontinence
Terzoni et al, 2013 ²⁴	Italy	Cohort	SUI	To compare the effects of EMS and FES on involuntary leakages of urine in patients who underwent RP
Chang et al, 2015 ²⁵	Taiwan	Cohort	SUI, UII, MUI	To investigate the effect of EMS health-related QoL in patients with post-RP incontinence

EMS: extracorporeal magnetic stimulation; FES: functional electrical stimulation; min: minutes; MUI: mixed urinary incontinence; PFMT: pelvic floor muscle therapy; RCT: randomized controlled trial, RP: radical prostatectomy; SUI: stress urinary incontinence; UII: urgency urinary incontinence.

Table 3. Summary of the studies identified in this systematic review

Study	Assigned groups	Types of intervention	Total EMS dose	Outcome measures	Results
Yokoyama et al, 2004 ¹⁹	EMS: 12 FES: 12 Control: 12	FES: anal electrode pulses of 20 Hz square waves at a 300 µs pulse duration and a maximal output current of 24 mA were used for 15 minutes twice a day for 1 month EMS: 10 Hz, intermittently for 10 minutes, followed by a rest period of 2 minutes, and a second treatment at 50 Hz intermittently for 10 minutes; 2x per week for 2 months Control: PFMT supine, anal muscle contracture with verbal instruction to selectively contract anal sphincter	576 000 contractions in 8 weeks	Primary: 24h pad weight Secondary: IQoL adverse events	No significant differences were noted between either FES or EMS group, and the control group after 3 months of treatment, and no statistical differences in 24h pad weight between FES and EMS.
Yokoyama et al, 2005 ²³	EMS 10	EMS: 10 Hz for 10 min + 50 Hz for 10 min; 2 sessions/week for 2 months	576 000 contractions in 8 weeks	Primary: Daily urinary frequency, 1-h pad weight, UDS Secondary: IQoL	EMS improves urodynamic parameters and I-QoL scores from baseline at the end of treatment. However, 30% of participants returned to baseline IQoL scores by 12 months post-treatment.
Suzuki et al, 2007 ¹⁸	EMS to Sham: 9 Sham to EMS: 7	EMS: 10 Hz continuously with a stimulating pulse width of 300µs for 20 min; 1 session/week for 10 weeks	120 000 contractions in 10 weeks	Primary: Leaks/week Secondary: ICIQ-SF, UDS	There were no significant sex differences in reductions of number of

		Sham: 1 Hz in 5-sec “on”–5-sec “off” pulsing manner with 300µs-width and 20%, or less, of the maximum output for 20 min; 1 session/week for 10 weeks			leaks/weeks, or total ICIQ-SF score. EMS decreased leak frequency by 35.2%.
Liu et al, 2008 ²¹	EMS: 12 PFMT: 12	EMS: 10 Hz for 10 minutes, followed by a 3-minute rest and a second treatment at 50 Hz for 20 minutes; 30 minutes 2x/week for 6 weeks PFMT: Contraction for 3 seconds followed by a 3-second rest, 20 minutes once, 3x/day for 6 weeks	792 000 contractions in 6 weeks	Primary: IQoL, ICIQ-SF	After 3 and 6 months of treatment, participants treated with EMS had significantly lower QoL and ICIQ-SF scores than those receiving PFMT.
Koo et al, 2009 ²⁰	EMS: 16 PFMT: 16	EMS: 10hz for 10 min, 50hz for 10 min; 20 minutes 2x/week for 8 weeks - Total of 16 Sessions PFMT: pelvic floor muscle contractions held for 10 seconds and relaxed for 10 seconds. 10 reps per session, 10 sessions daily	576 000 contractions in 8 weeks	Primary: 24-hour pad weight and pad usage Secondary: IQoL	EMS significantly decreased 24h pad weight and pad usage compared to PFMT at 2 and 3 months after the start of treatment. There were no significant differences between treatment groups at 6 months.
Wöllner et al, 2012 ²²	EMS: 57	EMS: 25 Hz for 10 min, 50hz for 10 min at; 2–3 times per week for a total of 12 sessions	540 000 contractions in 4-6 weeks	Primary: Pads used/24h Secondary: Patient satisfaction, duration of therapy, and adverse events	EMS decreases pads used/24h with persistent or further reductions at 12.5 months.
Terzoni et al, 2013 ²⁶	EMS: 69 PFMT: 87 Control: 22	EMS: SUI & MUI: 50Hz, 5s on, 5s off 10 minutes for First treatment, 20 minutes subsequent. 2x/week for 6 weeks	Stress & mixed: 345 000 contractions	Primary: IPSS at 1 month Secondary: 24h pad weight	EMS reduces urine leakage quicker than PFMT within the first 6 weeks of treatment, however,

Extracorporeal magnetic stimulation to treat UI in men

		<p>UUI: 5Hz, 5s on, 5s off. 10 minutes for First treatment, 20 minutes subsequent. 2x/week for 6 weeks</p> <p>PFMT: 7 exercise schemes of progressively increasing difficulty; 2x daily after bladder emptying for 6 months</p> <p>Control: No treatment</p>	<p>ns in 6 weeks</p> <p>UUI: 34 500 contractions in 6 weeks</p>		<p>there was no difference to recovery at 6 months of treatment.</p>
Terzoni et al, 2013 ²⁴	EMS: 22 FES: 18	<p>EMS: 50 Hz, 5s on, 5s off. 10 min for the first session and 20 min in subsequent sessions, 2x per week for 6 weeks</p> <p>FES: biphasic currents at 50 Hz; 6 s of stimulation and 12 s of recovery time 10 min in first session or 15 min for the second, 2x per week for 6 weeks</p>	345 000 contractions in 6 weeks	Primary: Leakage (grams/day)	At 6 weeks of treatment, EMS resulted in quicker reductions in urine leakage compared to FES.
Chang et al, 2015 ²⁵	EMS: 13	<p>EMS: Pulse-field frequency was 10 Hz applied intermittently for 10 minutes, followed by a 1-minute rest period, then a second treatment at 50 Hz applied intermittently for 10 minutes; 20 minutes, 2x/week, 2 months</p>	576 000 contractions in 8 weeks	<p>Primary: Incontinence episodes, 3-day dairy outcomes</p> <p>Secondary: UDI6, IIQ-7, IPSS-QoL</p>	EMS resulted in significantly decreased incontinence episodes, improved urodynamic parameters and significantly lower UDI 6 scores.

EMS: extracorporeal magnetic stimulation; FES: functional electrical stimulation; ICIQ-UI SF: International Consultation on Incontinence Questionnaire-Urinary Incontinence Short Form; IPSS-QoL: International Prostate Symptom Score-QoL Index; I-QOL: Urinary Incontinence Quality of Life Scale for female patients; min: minutes; MUI: mixed urinary incontinence; PFMT: pelvic floor muscle therapy; SUI: stress urinary incontinence; UDI-6: Urogenital Distress Inventory; UDS: urodynamic studies; UUI: urgency urinary incontinence.

Table 4. Quality assessment scores based on the modified Downs and Black checklist

Study	Score	Quality index
Yokoyama et al, 2004 ¹⁹	16/27	Fair
Yokoyama et al, 2005 ²³	16/27	Fair
Suzuki et al, 2007 ¹⁸	23/27	Good
Liu et al, 2008 ²¹	19/27	Fair
Koo et al, 2009 ²⁰	17/27	Fair
Wöllner et al, 2012 ²²	14/27	Poor
Terzoni et al, 2013 ²⁶	18/27	Fair
Terzoni et al, 2013 ²⁴	16/27	Fair
Chang et al, 2015 ²⁵	19/27	Fair

DRAFT