

Case - Long-term management of oligo-progressive metastatic castrate sensitive prostate cancer with recurrence-directed radiotherapyShan Grewal^{1,3}, Hsin-Pei Hu^{1,2}, Andre Gouveia^{1,2}, Theodoros Tsakiridis^{1,2,3}¹Department of Radiation Oncology, Juravinski Cancer Center, Hamilton Health Sciences, Hamilton, ON, Canada;²Department of Oncology, Juravinski Cancer Center, Hamilton Health Sciences, Hamilton, ON, Canada; ³Center for Discovery in Cancer Research, McMaster University, Hamilton, ON, Canada**Cite as:** Grewal S, Hu H-P, Gouveia A, et al. Case - Long-term management of oligo-progressive metastatic castrate sensitive prostate cancer with recurrence-directed radiotherapy. *Can Urol Assoc J* 2025 November 25; Epub ahead of print. <http://dx.doi.org/10.5489/cuaj.9363>

Published online November 25, 2025

Corresponding author: Dr. Theos Tsakiridis, Radiation Oncology, Juravinski Cancer Center, Hamilton, ON, Canada; tsakirt@mcmaster.ca

INTRODUCTION

Although localized prostate cancer (PC) has a favourable prognosis with a 5-year progression-free survival (PFS) rate of 65-85%, many patients develop distant metastatic disease, which carries a reported 5-year survival rate of 41% in Canada in 2023 [1]. The survival rate for patients with metastatic castrate-sensitive PC (CSPC) has further improved through the use of androgen deprivation therapy (ADT), androgen pathway inhibitors (ARPIs), docetaxel chemotherapy, and metastasis- or recurrence-directed therapy (RDT). ADT remains a cornerstone in the management of recurrent and metastatic PC, but studies have reported its impact on patient-reported quality of life [2], increased cardiovascular morbidity and mortality [3], and osteoporotic skeletal fracture risks [4]. This side effect profile demonstrates a need to explore pathways to limit and/or postpone its use.

KEY MESSAGES

- While ADT is a cornerstone in the management of recurrent and metastatic prostate cancer, it can significantly impact patient-reported QoL and increases cardiovascular and osteoporotic morbidity and mortality.
- Metastasis- or recurrence-directed therapy (RDT) with the use of stereotactic body radiotherapy (SBRT) has shown promising efficacy in managing oligometastatic disease in various cancer types.
- Future studies may further define the role of RDT in the long-term management of metastatic prostate cancer to delay systemic therapy in certain clinical scenarios.

RDT includes stereotactic body radiation therapy (SBRT) to precisely deliver high-dose radiation to metastatic sites with or without concomitant systemic therapy, elective nodal irradiation, and surgery. SBRT has shown promising efficacy in managing oligometastatic disease in various cancer types.

This case report details a patient with CSPC who presented with episodes of oligo-progression, resisted treatment with standard of care ADT over a period of 9 years, and was managed with nine courses of RDT with limited ADT. It highlights the potential of RDT in addressing some therapeutic challenges posed by progressing metastatic PC.

CASE REPORT

A 61-year-old male presented with an elevated PSA of 5.53 ng/mL in 2012. Medical history was significant for hypertension, sleep apnea, knee arthroplasty, and burn trauma surgery. He underwent laparoscopic radical prostatectomy revealing pT3a, N0, ISUP Grade Group 3 [Gleason Score of 7 (4 + 3)] prostatic adenocarcinoma. Post-surgery PSA was 0.066 ng/mL, and it increased to 0.22 ng/mL in April 2014 with a doubling time of 6 months. He then participated in the RTOG-0534 trial and was randomized to prostatic bed radiotherapy alone, 6480 cGy in 36 daily fractions, for salvage treatment (Fig. 1A).

His PSA reached a nadir of 0.09 ng/mL in October 2014, but it progressed to 2.4 ng/mL over the next four years, with a 10-month doubling time (Fig. 2A). The patient denied physical symptoms, but a bone scan and CT scans in June 2018 revealed metastatic foci in the chest wall (L-3rd rib) and spine (T7) (Fig. 1B, 1D). He refused standard of care ADT and requested local therapy alone. Both metastatic sites were treated with low dose radiotherapy of 2000 cGy in five consecutive daily fractions using 3D-conformal techniques without complications (Fig. 1C, 1E).

His PSA decreased significantly to 0.48 ng/mL by November 2018 but rose again to 1.0 ng/mL after 10 months (Fig. 2A). Prostate-specific membrane antigen - positron emission tomography PSMA-PET/CT (¹⁸F-DCFPyL) scan showed no activity in previously detected sites but identified new foci in the right scapula, right ilium, sacrum, and two iliac nodes (Fig. 1F). The patient once again declined ADT, opting for further courses of RDT (Fig. 1G-L). All newly identified sites were treated with SBRT in March 2020 using moderate dose (3000cGy in five fractions delivered every other day). He had no observable toxicity and had rapid biochemical response (Fig. 2A).

DISCUSSION

We report a patient with oligo-progressive CSPC who underwent well-tolerated, lower biological effective dose RDT, guided by high-sensitivity modern PSMA-PET imaging, to achieve biochemical disease control over a period of 5 years while deferring ADT for a significant amount of time. This treatment approach is unconventional and diverges from clinical guidelines, driven by the patient's repeated refusal of systemic therapy. Although unconventional, this presented case and surmounting research underscore the promise of RDT for efficacy in managing metastatic lesions, prompting further consideration of the timing and necessity of ADT

in certain clinical scenarios. Once systemic therapy was initiated in this patient, the combined use of ADT and SBRT yielded durable biochemical remission consistent with the synergy seen in studies [5,6].

The intensification of systemic therapy with the addition of ARPIs to ADT +/- chemotherapy has been elevated to the standard of care for de novo metastatic CSPC after recent studies demonstrated failure-free and overall survival benefit with their use [7-9]. Within this patient population, recent studies have identified an emerging role for consolidative ablative local radiotherapy to active disease sites in metastatic PC. The STAMPEDE and PEACE1 trials showed improved PFS and overall survival benefit with the addition of prostate radiotherapy to standard of care ADT in patients with low volume metastatic CSPC [10,11]. The ORIOLE and STOMP phase II randomized trials demonstrated that compared to surveillance, SBRT for oligometastases in metastatic PC with ≤ 3 lesions improved PFS (61% vs. 19% at 6 months) and ADT-free survival (34% vs. 8% at 5 years), with few serious toxicity events [5, 12]. These outcomes reflect the broader trend in oligometastatic cancers, where the SABR-COMET Phase II trial found patients with ≤ 5 metastases who received ablative therapy in addition to the standard-of-care saw enhanced 5-year progression-free survival (17% vs. 3%) and median overall survival (50 months vs. 28 months) [13]. His PSA steadily rose to 3.3 ng/mL in the subsequent 2 years. A PSMA-PET/CT scan in July 2022 identified disease recurrence at the previously treated T7 vertebra and two novel sites in a perirectal and left external iliac lymph node (Fig. 1M). At this point, the patient agreed to ADT alongside SBRT (3000cGy in five fractions delivered every other day) to the new metastatic sites (Fig. 1N–S). The patient had no observable gastrointestinal or neurologic toxicity from the treatments. Undetectable PSA levels were obtained in September 2022 and in January 2023. At that point ADT was discontinued on patient's request. The patient has remained in biochemical remission (undetectable PSA) as testosterone began to rise to eugonadal testosterone levels (T=5.2 nmol/L) in October 2023. At the time of the preparation of this report, the patient enjoyed 18 months of eugonadal complete biochemical remission (Fig. 2A).

DISCUSSION

We report a patient with oligo-progressive CSPC who underwent well-tolerated, lower biological effective dose RDT, guided by high-sensitivity modern PSMA-PET imaging, to achieve biochemical disease control over a period of 5 years while deferring ADT for a significant amount of time. This treatment approach is unconventional and diverges from clinical guidelines, driven by the patient's repeated refusal of systemic therapy. Although unconventional, this presented case and surmounting research underscore the promise of RDT for efficacy in managing metastatic lesions, prompting further consideration of the timing and necessity of ADT in certain clinical scenarios. Once systemic therapy was initiated in this patient, the combined use of ADT and SBRT yielded durable biochemical remission consistent with the synergy seen in studies [5,6]. The intensification of systemic therapy with the addition of ARPIs to ADT +/- chemotherapy has been elevated to the standard of care for de novo metastatic CSPC after recent

studies demonstrated failure-free and overall survival benefit with their use [7-9]. Within this patient population, recent studies have identified an emerging role for consolidative ablative local radiotherapy to active disease sites in metastatic PC. The STAMPEDE and PEACE1 trials showed improved PFS and overall survival benefit with the addition of prostate radiotherapy to standard of care ADT in patients with low volume metastatic CSPC [10,11]. The ORIOLE and STOMP phase II randomized trials demonstrated that compared to surveillance, SBRT for oligometastases in metastatic PC with ≤ 3 lesions improved PFS (61% vs. 19% at 6 months) and ADT-free survival (34% vs. 8% at 5 years), with few serious toxicity events [5, 12]. These outcomes reflect the broader trend in oligometastatic cancers, where the SABR-COMET Phase II trial found patients with ≤ 5 metastases who received ablative therapy in addition to the standard-of-care saw enhanced 5-year progression-free survival (17% vs. 3%) and median overall survival (50 months vs. 28 months) [13].

In the absence of systemic therapy, metastasis-directed therapies alone, including SBRT and surgery, are capable of prolonging ADT-free survival in oligometastatic PC [14]. The findings align with the broader objective of metastasis- and recurrence-directed therapy in advanced PC: to prolong ADT-free survival, delay systemic treatment initiation and/or reduce systemic treatment duration, improving quality of life while controlling disease progression. Using a reversal of the conventional paradigm, studies have begun reporting RDT in oligometastatic CSPC with short-course ADT added as an adjunct treatment. Although there were no detected differences in the time to development of CRPC or eugonadal PFS rates, the randomized phase II trial RADIOSA showed significantly improved clinical and biochemical PFS with the addition of 6-month ADT in patients with metachronous oligometastatic CSPC treated with SBRT [15]. Such findings support a growing emphasis on personalized approaches to RDT and ADT.

CONCLUSIONS

Overall, recurrence-directed SBRT has evidently emerged as a promising and transformative tool to help manage of early metastatic prostate cancer. The case we present here suggests that SBRT, in combination with PSMA-PET imaging, may be a useful tool for long-term management of oligo-progressive CSPC that could help delay systemic therapy in certain clinical scenarios and/or complement short term ADT as an approach to preserving quality of life in patient with limited metastatic burden.

REFERENCES

1. Canadian cancer statistics 2023. Canadian cancer statistics advisory committee in collaboration with the Canadian Cancer Society, Statistics Canada and the Public Health Agency of Canada. Available at: https://cdn.cancer.ca/-/media/files/research/cancer-statistics/2023-statistics/2023_PDF_EN.pdf (accessed July 19, 2025)
2. Gay HA, Sanda MG, Liu J, et al. External beam radiation therapy or brachytherapy with or without short-course neoadjuvant androgen deprivation therapy: Results of a multicenter, prospective study of quality of life. *Int J Radiat Oncol Biol Phys* 2017; 98:304-17. <https://doi.org/10.1016/j.ijrobp.2017.02.019>
3. Saigal CS, Gore JL, Krupski TL, et al. Androgen deprivation therapy increases cardiovascular morbidity in men with prostate cancer. *Cancer* 2007;110:1493-500. <https://doi.org/10.1002/cncr.22933>
4. Shahinian VB, Kuo Y-F, Freeman JL, et al. Risk of fracture after androgen deprivation for prostate cancer. *N Engl J Med* 2005;352:154-64. <https://doi.org/10.1056/NEJMoa041943>
5. Ost P, Reynders D, Decaestecker K, et al. Surveillance or metastasis-directed therapy for oligometastatic prostate cancer recurrence: A prospective, randomized, multicenter, phase 2 trial. *J Clin Oncol* 2018;36:446-53. <https://doi.org/10.1200/JCO.2017.75.4853>
6. Bolla M, Van Tienhoven G, Warde P, et al. External irradiation with or without long-term androgen suppression for prostate cancer with high metastatic risk: 10-year results of an EORTC randomized study. *Lancet Oncol* 2010;11:1066-73. [https://doi.org/10.1016/S1470-2045\(10\)70223-0](https://doi.org/10.1016/S1470-2045(10)70223-0)
7. James ND, de Bono JS, Spears MR, et al. STAMPEDE Investigators. Abiraterone for prostate cancer not previously treated with hormone therapy. *N Engl J Med* 2017;377:338-51. <https://doi.org/10.1056/NEJMoa1702900>
8. Chi KN, Agarwal N, Bjartell A, et al. TITAN Investigators. Apalutamide for metastatic, castration-sensitive prostate cancer. *N Engl J Med* 2019;381:13-24. <https://doi.org/10.1056/NEJMoa1903307>
9. Smith MR, Hussain M, Saad F, et al. ARASENS Trial investigators. Darolutamide and survival in metastatic, hormone-sensitive prostate cancer. *N Engl J Med* 2022;386:1132-42. <https://doi.org/10.1056/NEJMoa2119115>
10. Parker CC, James ND, Brawley CD, et al. STAMPEDE Trial collaborative group. Radiotherapy to the prostate for men with metastatic prostate cancer in the UK and Switzerland: Long-term results from the STAMPEDE randomised controlled trial. *PLoS Med* 2022;19:e1003998. <https://doi.org/10.1371/journal.pmed.1003998>
11. Bossi A, Foulon S, Maldonado X, et al. PEACE-1 investigators. Efficacy and safety of prostate radiotherapy in de novo metastatic castration-sensitive prostate cancer (PEACE-1): a multicentre, open-label, randomized, phase 3 study with a 2 × 2 factorial design. *Lancet* 2024;404:2065-76. [https://doi.org/10.1016/S0140-6736\(24\)01865-8](https://doi.org/10.1016/S0140-6736(24)01865-8)
12. Phillips R, Shi WY, Deek M, et al. Outcomes of observation vs. stereotactic ablative radiation for oligometastatic prostate cancer: The ORIOLE phase 2 randomized clinical trial. *JAMA Oncol* 2020;6:650-9. <https://doi.org/10.1001/jamaoncol.2020.0147>
13. Palma DA, Olson R, Harrow S, et al. Stereotactic ablative radiotherapy for the comprehensive treatment of oligometastatic cancers: Long-term results of the SABR-COMET phase 2 randomized trial. *J Clin Oncol* 2020;38:2830-38. <https://doi.org/10.1200/JCO.20.00818>

14. Glicksman RM, Metser U, Vines D, et al. Curative-intent metastasis-directed therapies for molecularly-defined oligorecurrent prostate cancer: A prospective phase 2 trial testing the oligometastasis hypothesis [Formula presented]. *Eur Urol* 2021;80:374-82. <https://doi.org/10.1016/j.eururo.2021.02.031>
15. Marvaso G, Corrao G, Zaffaroni M, et al. ADT with SBRT vs. SBRT alone for hormone-sensitive oligorecurrent prostate cancer (RADIOSA): A randomized, open-label, phase 2 clinical trial. *Lancet Oncol* 2025;26:300-11. [https://doi.org/10.1016/S1470-2045\(24\)00730-7](https://doi.org/10.1016/S1470-2045(24)00730-7)

DRAFT

FIGURES AND TABLES

Figure 1. Imaging and radiotherapy (RT) plans for primary and metastatic prostate cancer. (A) Radiation plan for prostate bed treated in 2014. (B, D) Bone scan in June 2018 revealed metastatic foci in the third rib and T7 spine (green arrows). (C, E) Three-dimensional conformal radiation plans (2000 cGy in 5 fractions) for metastases in the T7 spine and third rib. (F) Prostate-specific membrane antigen positron emission tomography (PSMA-PET) scan in January 2020 showed complete response in previous metastases but identified several new sites (green arrows from top to bottom): Right scapula (G), sacrum (I), two right external iliac lymph nodes (one shown) and right ilium (K). Stereotactic body radiation therapy (SBRT) plans (3000 cGy in 5 fractions) for the metastatic sites at right scapula (H), sacrum (J), and one of the right external iliac nodes and right ilium (L). (M) PSMA-PET scan in June 2022 showed disease recurrence at the T7 vertebra (red arrow; N), plus novel sites in the right external iliac lymph node (top green arrow; P) and perirectal lymph node (bottom green arrow; R). SBRT plans (3000 cGy in 5 fractions) for the re-treatment of T7 spine (O) and the treatment of the right external iliac lymph node (P) and perirectal lymph node (S).





Figure 2. Summary of disease course, prostate-specific antigen (PSA) and testosterone measurements, and radiotherapy treatments. (A) Patient PSA and testosterone trends since diagnosis. (B) Summary of disease sites and corresponding treatment regimen.



DRAFT