

Case - Penile mycobacterium avium complex infection

Angie K. Puerto Nino^{1,2}, Adrienne K. Chan^{3,4}, Scott Walsh⁵,
Ron Kodama^{1,6}, Sarah Neu^{1,6}

¹Division of Urology, Department of Surgery, University of Toronto, Toronto, ON, Canada; ²Faculty of Medicine, University of Helsinki, Helsinki, Finland; ³Division of Infectious Diseases, Sunnybrook Health Sciences Centre, Toronto, ON, Canada; ⁴Division of Clinical Public Health, Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada; ⁵Division of Dermatology, Department of Medicine, University of Toronto, Toronto, ON, Canada; ⁶Division of Urology, Sunnybrook Health Sciences Center, Toronto, ON, Canada

Cite as: Puerto Nino AK, Chan AK, Walsh S, et al. Case - Penile mycobacterium avium complex infection. *Can Urol Assoc J* 2026;20(9):E381-4. <http://dx.doi.org/10.5489/cuaj.9604>

Published online May 4, 2026

INTRODUCTION

Nontuberculous mycobacteria (NTM), including those in the *Mycobacterium avium* complex (MAC), are a common environmental microorganism that can be found in soil, water sources, household plumbing, and various moist environments, including the oral cavity.¹ MAC are increasingly recognized as opportunistic pathogens.² Although MAC are associated with pulmonary disease and disseminated infection in immunocompromised patients, the incidence of isolated extrapulmonary manifestations is rising.³ We present a rare case of penile infection caused by MAC to contribute to the understanding of its clinical presentation and management. Because penile involvement is exceptionally uncommon, we performed a literature review to contextualize this case and to illustrate the broader spectrum of genitourinary NTM infections. In doing so, we highlight the diagnostic challenges that such unusual presentations pose and offer insights to help clinicians recognize and treat similar rare infections.

CASE REPORT

A 59-year-old male with a history of suspected iatrogenic urethral injury during catheter placement at the age of 10 and Peyronie's disease presented with worsening voiding urinary tract symptoms. The patient was diagnosed with significant stricture disease from the mid-penile to the distal bulbar urethra measuring 4 cm in length, and 12-French in diameter. Subsequently, the patient underwent one-stage dorsal onlay anterior urethral reconstruction with bilateral buccal mucosa grafting.

Eight months after the urethral reconstruction, the patient developed distal penile pain and an erythematous plaque on the dorsum of his penis. Biopsy of the lesion showed features of lichen planus and he received treatment with betamethasone dipropionate. The patient did not report improvement of his symptoms after steroid cream. Six months later, meatal stenosis and hardness within his corporal bodies and the glans were found on the patient's physical exam. MRI of the penis showed lesions in the glans extending to the urethra (Figure 1A). Excisional biopsy and 4 core needle biopsies of the glans lesion showed a suppurative granulomatous process with excessive epithelial reactions. The urology team referred the patient to the infectious diseases and dermatology services. Stains and culture for bacteria, fungi, and acid-fast bacilli (AFB) were negative. Initially, the patient received treatment for pyogenic superinfection with ciprofloxacin for two weeks, and then, amoxicillin-clavulanate for two weeks to cover for the remote possibility of actinomycosis.

No improvement was seen, and further studies were performed. Blastomyces and Histoplasma serology, urine Histoplasma Ag, and urine Blastomyces Ag were negative. C-reactive protein, anti-neutrophil cytoplasmic antibody, HIV, and syphilis serology were negative. The patient's urine culture for AFB showed 1/3 positive for MAC. MRI repeated three months after initial MRI showed a 3.2 × 2.2 cm enlarging heterogeneous T2 signal-enhancing lesion almost replacing the penile glans (Figure 1B). The lesion involved the most distal portion of both cavernosa. Additionally, along the ventral aspect of the spongiosum, there was a new 0.8 cm enhancing soft tissue nodule. The lesion progressed, leading to splitting of the glans (Figure 2A). Therefore, the patient underwent a repeat biopsy which showed necrotizing granuloma-




Figure 1. Results of magnetic resonance imaging of the penis.

KEY MESSAGES

- *Mycobacterium avium* complex (MAC) is a nontuberculous mycobacterium typically associated with pulmonary disease or disseminated infection in the immunocompromised. Isolated genitourinary involvement is rare.
- We report a unique case of penile MAC infection in a 59-year-old man with a history of urethral reconstruction. The patient developed progressive penile lesions, ultimately confirmed as MAC. He was treated with prolonged multidrug therapy achieving clinical improvement.
- Given the rarity of penile involvement, we review the literature to highlight diagnostic challenges, therapeutic considerations, and the need for awareness of atypical penile MAC infection.

tous inflammation. Grocott methenamine silver (GMS) and AFB stains were negative as was polymerase chain reaction (PCR) testing for both *M. tuberculosis* complex and MAC. Urine AFB culture showed 4/4 positive results for MAC and the isolate was sent for antimicrobial susceptibility testing to the National Microbiology Laboratory of Canada by broth microdilution minimum inhibitory concentration (MIC) panel (ug/mL): amikacin 256, ciprofloxacin 4, clarithromycin 4, clofazamine 0.25, doxycycline >8, linezolid 32, minocycline >8, rifabutin 0.5, rifampin >4, streptomycin >32, trimethoprim-sulfamethoxazole 4/76. With negative fungal cultures and without an alternative diagnosis, in the context of a recent reconstruction procedure, the patient was therefore treated empirically for penile MAC and started on rifabutin (due to drug interactions with rifampin), azithromycin, and ethambutol.

After the patient was started on the therapy, urine cultures for AFB repeated at three and five months continued to be positive for MAC. He also had bronchoalveolar lavage (BAL) for an asymptomatic 8 mm pulmonary nodule found on chest CT that was culture positive for MAC. He continued to have distal penile pain and urethral discharge. During this time he had a supra-pubic tube (SPT) for urinary drainage. Moxifloxacin was added to his therapy. Repeated MRIs




Figure 2. Clinical findings of the patient's physical exam.

showed a progressive decrease in the size of the lesion without fistulous tracts. Physical exam continued to show slow improvement at six and 16 months as demonstrated in Figures 2B and 2C. The patient received a total of 20 months of antibiotic therapy. Afterwards, the patient was found to have meatal stenosis and underwent a distal first-stage anterior urethroplasty up to the fossa navicularis, and his SPT was removed.

Table 1. Publications of penile MAC infection

Publication	Clinical presentation	Diagnosis	Treatment	Clinical progress
Niedrach et al, 1989 ¹⁰	Painful abscess	MAC, <i>Streptococcus haemolyticus</i> , and <i>Clostridium perfringens</i>	Surgical drainage and cefoxitin for 4 days followed by penicillin V potassium	No recurrence 4 months post-intervention
Ralph et al, 1998 ¹¹	Penile ulcer infiltrating the corpus cavernosum	Cutaneous MAC	8 months of: - Rifabutin 300 mg/d - Clarithromycin 500 mg BID - Ethambutol 400 mg TID	Resolution of the ulcer at the end of the 8 months
Fonda-Pascual et al, 2018 ¹²	- Several yellowish firm papules on the penile shaft and foreskin - Indurated ulcer with mamelonated borders and a fibrinous base on the foreskin	MAC cutaneous involvement with fusocellular pseudotumoral pattern	Not reported	Not reported

MAC: mycobacterium avium complex.

After this procedure and to date, the patient remains asymptomatic and is voiding without difficulty. His MAC cultures continue to be negative now, 25 months out.

DISCUSSION

Penile lesions may result from a wide variety of etiologies, including inflammation, infection, trauma, or malignancy, making them a significant diagnostic challenge. Given this broad differential, careful clinical assessment supported by histopathological and microbiological testing is essential to establish an accurate diagnosis and to guide treatment. This is particularly important in infections caused by rare pathogens, such as nontuberculous mycobacteria, which can mimic more prevalent conditions. Infectious causes of penile lesions often include sexually transmitted infections, such as syphilis, herpes simplex virus, and human papillomavirus. Other common culprits are bacterial infections such as cellulitis or folliculitis, and fungal infections like candidiasis. Failure to correctly identify the etiology can lead to misdiagnosis, which can result in ineffective treatment, increased morbidity, or disease progression and dissemination.⁴

Penile infections caused by MAC are rare and may present with nonspecific symptoms, including chronic inflammation, nodules, ulcerations, or abscess formation. These clinical manifestations can easily be mistaken for other infectious or neoplastic conditions. Because of this diagnostic overlap, a high degree of clinical suspicion and advanced diagnostic methods are often necessary to distinguish MAC from more common causes.⁵ The exact incidence of MAC disease remains uncertain but is rarely described, and its pathway of dissemination is not yet fully understood. In this case, it is possible that the patient was colonized with MAC, as is commonly

found in the human oral cavity,¹ and that the infection may have been introduced to the genital region during placement of the buccal graft at the time of urethroplasty. Table 1 describes the different reports published to date on the different presentations of penile MAC infection. Other reported genitourinary presentations include asymptomatic sterile pyuria,⁶ hematuria,⁷ granulomatous prostatitis,^{7,8} or symptoms of urinary infection and sepsis.⁹

The diagnosis can involve a combination of culture, molecular testing, and histopathological examination to confirm the microorganism and exclude other granulomatous diseases or malignancies. Confirmation of MAC infection in genitourinary specimens usually requires AFB staining, urine culture on specialized mycobacterial media, and molecular methods such as PCR or gene sequencing to achieve definitive identification.^{6,13}

An important caveat is that the clinical significance of MAC isolation from urine must be interpreted with caution, as it is most often due to contamination from environmental sources rather than true urinary tract infection, especially in non-sterile specimens, such as midstream urine samples. Molecular characterization has shown that environmental strains of MAC can be found in hospital water systems and may contaminate urine samples, leading to false positive results for infection.¹⁴

Distinguishing between colonization, contamination, and true infection requires correlation with clinical findings and, ideally, molecular typing to differentiate environmental from pathogenic strains. Urine cultures for atypical organisms, including nontuberculous mycobacteria, should be performed only when there is a strong

clinical suspicion, such as in this case where there were pathological findings of granulomatous disease.

Managing penile MAC infections is often challenging. To our knowledge, no specific protocol for its management has been established. A wide variety of treatments was used in the reported cases (Table 1). Medical treatment requires prolonged multidrug therapy tailored to the antimicrobial susceptibility results, based on our experience.

In some cases, particularly with localized disease or abscess formation, surgical intervention may be necessary. Standard regimens generally include at least three drugs, including ethambutol and a macrolide;¹⁵ however, treatment outcomes can vary, highlighting the importance of individualized management based on the microorganism's susceptibility to medical treatment.¹⁶ The emergence of drug resistance and patient intolerance to standard regimens further complicate care, underscoring the need for novel treatment options and personalized approaches.¹⁷

CONCLUSIONS

This review highlights the diagnostic difficulties and treatment challenges posed by penile MAC infections, with particular attention to its nonspecific clinical presentation. Ongoing research is needed to improve diagnostic approaches and develop more effective treatment strategies for these rare cases.

COMPETING INTERESTS: The authors have no relevant conflicts of interest to report.

This paper has been peer reviewed.

REFERENCES

- Kompa KG, Trotter CA, Hyman CL, et al. Disseminated mycobacterium avium complex myositis in a patient with graft-versus-host disease. *Open Forum Infect Dis* 2022 Jul 29;9:ofac385. <https://doi.org/10.1093/ofid/ofac385>
- Kim, BG, Jhun, B.W, Kim H, et al. Treatment outcomes of mycobacterium avium complex pulmonary disease according to disease severity. *Sci Rep* 2022. <https://doi.org/10.1038/s41598-022-06022-z>
- Marshall JE, Mercado RA, Lipner EM, et al. Nontuberculous mycobacteria testing and culture positivity in the United States. *BMC Infect Dis* 2024;24:288. <https://doi.org/10.1186/s12879-024-09216-0>
- Moule MG, Cirillo JD. Mycobacterium tuberculosis dissemination plays a critical role in pathogenesis. *Front Cell Infect Microbiol* 2020;10:65. <https://doi.org/10.3389/fcimb.2020.00065>
- Sousa C, Marchiori E, Youssef A, et al. Chest imaging in systemic endemic mycoses. *J Fungi (Basel)* 2022;8:1132. <https://doi.org/10.3390/jof8111132>
- Yang K, Samplaski M, Mazzulli T, et al. *Can Urol Assoc J* 2016;10:E186-8. <http://dx.doi.org/10.5489/cuoj.3363> Published online May 12, 2016
- Clark R, Cardona L, Valainis G, et al. Genitourinary infections caused by mycobacteria other than *Mycobacterium tuberculosis*. *Tubercle* 1989;70:297-300. [https://doi.org/10.1016/0041-3879\(89\)90025-1](https://doi.org/10.1016/0041-3879(89)90025-1)
- Mikolich DJ, Mates SM. Granulomatous prostatitis due to *Mycobacterium avium* complex. *Clin Infect Dis* 1992;14:589-91. <https://doi.org/10.1093/cid/14.2.589>
- Huang CT, Chen CY, Chen HY, et al. Genitourinary infections caused by nontuberculous mycobacteria at a university hospital in Taiwan, 1996-2008. *Clin Microbiol Infect* 2010;16:1585-90. <https://doi.org/10.1111/j.1469-0691.2010.03180.x>
- WL Niedrach, RM Lerner, CA Linke. Penile abscess involving the corpus cavernosum: A case report. *J Urol* 1989;141:374-5. [https://doi.org/10.1016/S0022-5347\(17\)40773-7](https://doi.org/10.1016/S0022-5347(17)40773-7)
- Ralph SG, Wilcox MH, Wilson JD. Mycobacterium avium-intracellulare complex causing penile ulceration in an immunocompetent man. *Int J STD AIDS* 1998;9:56-7. <https://doi.org/10.1258/0956462981920900>
- Fondo-Pascual P, Arana-Raja A, Moreno-Arrones OM, et al. Penile *Mycobacterium avium* complex spindle cell pseudotumor. *Indian J Dermatol Venereol Leprol* 2018;84:199-202. https://doi.org/10.4103/ijdv.IJDVL_403_17
- Hara R, Kitada S, Iwai A, et al. Diagnostic validity of gastric aspirate culture in nontuberculous mycobacterial pulmonary disease. *Ann Am Thorac Soc* 2020;17:1536-41. <https://doi.org/10.1513/AnnalsATS.201911-852OC>
- Alvarez J, García I. G, Aranaz A, et al. Genetic diversity of mycobacterium avium isolates recovered from clinical samples and from the environment: Molecular characterization for diagnostic purposes. *J Clin Microbiol* 2008;46:1246-51. <https://doi.org/10.1128/JCM.01621-07>
- Daley C. L, Iaccarino J. M, Lange C, et al. Treatment of nontuberculous mycobacterial pulmonary disease: An official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Clin Infect Dis* 2020;71:e1-e36. <https://doi.org/10.1093/cid/cia241>
- Zimenkov D. Variability of mycobacterium avium complex isolates drug susceptibility testing by broth microdilution. *Antibiotics (Basel)* 2022;11:1756. <https://doi.org/10.3390/antibiotics11121756>
- Calcagno A, Coppola N, Sarmati L, et al. Drugs for treating infections caused by nontuberculous mycobacteria: A narrative review from the study group on mycobacteria of the Italian Society of Infectious Diseases and Tropical Medicine. *Infection* 2024;52:737-65. <https://doi.org/10.1007/s15010-024-02183-3>

CORRESPONDENCE: Dr. Sarah Neu, Division of Urology, Department of Surgery, University of Toronto, Toronto, ON, Canada; sarah.neu@sunnybrook.ca