

Case - Extraskelatal Ewing sarcoma of the prostate masquerading as small cell neuroendocrine carcinoma

Areeb Hassan¹, Eric Winquist², Glenn Bauman², Brant A. Inman^{2,3}, Terence Tang⁴

¹Schulich School of Medicine and Dentistry, Western University, London, ON, Canada;

²Department of Oncology, Schulich School of Medicine and Dentistry, Western University and Verspeeten Family Cancer Centre, London Health Sciences Centre, London, ON, Canada;

³Division of Urology, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada; ⁴Department of Radiation Oncology, Odette Cancer Centre, Sunnybrook Health Sciences Centre, Toronto, ON, Canada

Cite as: Hassan A, Winquist E, Bauman G, et al. Case - Extraskelatal Ewing sarcoma of the prostate masquerading as small cell neuroendocrine carcinoma. *Can Urol Assoc J* 2026;20(11):E481-4. <http://dx.doi.org/10.5489/cuaj.9626>

Published online July 20, 2026

INTRODUCTION

Ewing sarcoma (ES) is a rare, small, round, blue cell tumor that is characterized by specific chromosomal translocations. Most ES arise in bone, but 20–30% are extraskelatal (EES).¹ EES occur in soft tissue and viscera and is generally diagnosed in adults over the age of 35.¹ Most ES arise from a gene translocation/fusion between a gene of the FET family (e.g., FUS, EWSR1, TAF15) and a gene of the ETS transcription factor family (e.g., ERG, FLI1, FEV).²⁻⁴

Histologically, ES appears similar to other small, round, blue cell tumors, such as small cell neuroendocrine carcinoma (SCNC), neuroblastoma, and lymphomas, making the histologic diagnosis of ES challenging. In this case report, we present the diagnostic and treatment challenges associated with EES of the prostate, which was initially diagnosed as SCNC.

CASE REPORT

A 33-year-old male mechanic presented with progressive obstructive lower urinary tract symptoms, rectal pain, and decreased stool calibre. On digital rectal exam, he had a firm, irregular mass involving the right prostate lobe. His prostate-specific antigen (PSA) was 0.73 with a free-to-total ratio of 0.18. Computed tomography (CT) scan revealed an enlarged, diffusely heterogeneous prostate with a focal mass in the right posterolateral gland with extracapsular extension. There was no evidence of metastatic disease, confirmed by a bone scan.

Pelvic magnetic resonance imaging (MRI) showed diffuse signal intensity throughout the central and peripheral zones of the prostate, with suspected extraprostatic extension from mid-gland to apex abutting the right anterior rectal wall. Measured prostate volume was 92 mL (Figure 1).

A transrectal ultrasound (TRUS)-guided biopsy was done, demonstrating small, round, blue cells that stained positive for CKPAN, synaptophysin, and Ki-67 on immunohistochemistry, with weak focal staining of chromogranin. They stained negative for NKX3.1, GATA 3, CK7, S100, Melan-A, CD45, TTF-1, Rb, P40, CD56, and desmin (Figure 2). Given the positive staining for neuroendocrine markers, a preliminary diagnosis of high-grade neuroendocrine carcinoma of the prostate was made. An outside second opinion was sought. Meanwhile, the case was reviewed at a local tumor board, and as the patient was very symptomatic, curative-intent chemotherapy and radiation were recommended based on a provisional diagnosis of limited-stage prostatic SCNC.

The patient was prescribed cisplatin and etoposide, and radiation to the prostate—60 Gy in 20 fractions—was delivered concurrent with the fourth cycle of chemotherapy. A subsequent pelvic MRI demonstrated a dramatic reduction in prostate volume to 34 mL. After six cycles of chemotherapy and all planned radiation, the external pathology report was issued and indicated diffuse CD99 expression with variable positivity in synaptophysin and chromogranin, but negative staining for NKX3.1, PSA, PSAP, INSM1, WT1, and MYF4. The working diagnosis was revised more broadly to a malignant small, round, blue cell tumor, recognizing the possibility of a fusion-associated sarcoma. To clarify this, further genetic testing via the Trusight RNA Pan-Cancer sequencing panel was requested.

The RNA panel identified the presence of an oncogenic EWSR1:FEV fusion, resulting in a final diagnosis of EES of the prostate, and the strong diffuse cytokeratin staining was interpreted as aberrant. The patient's case was discussed at a sarcoma multidisciplinary conference with experts from Canada and the U.S., and an ES regimen was recommended. As the patient had already completed six cycles of cisplatin and etoposide, a further eight cycles of alternating vincristine, doxorubicin, and cyclophosphamide (VAC)




Figure 1. Pre-treatment pelvic magnetic resonance imaging with T2-weighted images of the prostate in the (A) sagittal, (B) axial, and (C) coronal planes showing an enlarged prostate gland with extracapsular extension in the right posterolateral peripheral zone. (D) T1-weighted image of the prostate in the axial plane with evidence of extracapsular extension abutting the rectum.




Figure 2. Histologic images from initial prostate biopsy with (A) hematoxylin and eosin staining, as well as immunohistochemistry with (B) negative staining for NKX3.1, (C) positive staining for synaptophysin, and (D) weak focal staining for chromogranin.

with ifosfamide and etoposide (IE) chemotherapy was advised.

One member of the team (EW) met with the patient and his partner. The change in cancer diagnosis and treatment plan was discussed with complete transparency and an apology. The patient acknowledged a good understanding of the rarity of his diagnosis, the extra time required to confirm this, as well as the rationale for additional chemotherapy. Greatly improved lower urinary tract symptoms and evidence of tumor response on imaging helped with the delivery of this difficult news.

The patient completed the additional chemotherapy despite experiencing much fatigue over the last few months of treatment. Repeat TRUS-guided biopsy was negative for residual malignancy. The patient currently remains well, with restaging scans up to 16 months post-treatment confirming no local or distant recurrence (Figure 3).

DISCUSSION

In this case report, EES of the prostate was initially diagnosed and treated as SCNC, given the limited information available. Diagnosing ES in adults can be difficult due to its rarity, which has been reflected in other case reports, including EES of the kidney initially diagnosed as renal cell carcinoma and intracranial Ewing sarcoma initially diagnosed as a glioblastoma.^{5,6}

EES involving the prostate is exceedingly rare, with only nine other cases described in the literature.⁷ Our patient presented with non-specific symptoms, and his low PSA could not differentiate between EES or SCNC, as neither malignancy secretes PSA. Furthermore, ES and SCNC both appear as small, round, blue cell tumors with scant cytoplasm, and both can stain positive for cytokeratin, synaptophysin, and other neuroendocrine markers.^{5,8-10}

What established the diagnosis of EES in our patient was the identification of a pathognomonic EWSR1:FEV fusion via the TruSight RNA Pan-Cancer panel. This panel uses next-generation RNA sequencing to identify 1385 genes linked to oncogenesis and is superior to detecting fusion genes compared to DNA sequencing.¹¹ Given its high sensitivity, specificity, and short assay time, RNA sequencing is a valuable diagnostic tool that should be used earlier and more widely in diagnostic pathways; however, availability and cost may be barriers to implementation.

In North America, ES treatment consists of induction chemotherapy with nine alternating cycles of VAC and IE chemotherapy, followed by local treatment with

surgery or radiation.¹ After local treatment, consolidative chemotherapy with five alternating cycles of VAC and IE is given. Our patient initially received six cycles of dual-agent chemotherapy as opposed to a longer course of multi-agent induction chemotherapy.

Additionally, the patient received cisplatin, which is not commonly used in ES, and radiation was initiated concurrent with chemotherapy. Furthermore, the biologically equivalent dose of radiation that was delivered to the patient exceeded the standard doses used for ES. There is evidence to suggest that dose escalation improves local control in ES, and this may have contributed to the excellent disease response.¹² Radiation was delivered in a hypofractionated manner at 3 Gy per fraction vs. the standard 1.8–2 Gy per fraction regimens, although the latter were designed to minimize the late effects of radiation in the pediatric population and may be less relevant in adults.

Regarding local treatment options, previous evidence has suggested that surgery offers superior local control compared to radiation, but most of these studies were retrospective in nature and subject to selection bias.^{13,14} In our patient, radiation was given because of the initial concern for SCNC, which is a radiosensitive disease where surgery has a limited role; however, a delayed cystoprostatectomy—possibly with en bloc proctocolectomy—was considered but ultimately not undertaken given the negative post-treatment biopsies and the long-term morbidity of the procedure.

While the patient's treatment did deviate from the typical ES paradigm, there is no established standard of care for prostatic EES, given its rarity. For example, in one case report, a 33-year-old patient with prostatic EES underwent a prostatovesiculectomy, regional lymph node dissection, and pelvic floor resection followed by postoperative chemotherapy.⁷ In contrast, in another case report, a 29-year-old patient with prostatic EES was treated with neoadjuvant chemotherapy followed by a total pelvic exenteration.¹⁵ The variations in practice in managing prostatic EES are likely influenced by both local physician expertise and the rarity of the disease.

Despite deviations from the classic ES paradigm, the treatment our patient received has been effective thus far, with a marked reduction in prostate volume and a complete pathologic response on post-treatment biopsy. Further followup imaging and prostate biopsies are planned.




Figure 3. Post-treatment pelvic magnetic resonance imaging with T2-weighted images of the prostate in the (A) sagittal, (B) axial, and (C) coronal planes show response to treatment with a reduction in prostate size and no further evidence of extracapsular disease. (D) T1-weighted image of the prostate in the axial plane illustrates no evidence of extracapsular extension.

CONCLUSIONS

This case report of EES of the prostate highlights the challenges associated with diagnosing and treating rare prostatic malignancies, especially when establishing the definitive diagnosis is delayed and impacts treatment.

This paper has been peer reviewed.

COMPETING INTERESTS: The authors have no relevant conflicts of interest to report.

REFERENCES

1. Abboud A, Masrouha K, Saliba M, et al. Extraskelatal Ewing sarcoma: Diagnosis, management and prognosis (review). *Oncol Lett* 2021;21. <https://doi.org/10.3892/ol.2021.12615>
2. Mehmood K, Shehzad A. Rare site and age of presentation: Prostatic sarcoma of the Ewing family in an adolescent. *Indian J Musculoskelet Radiol* 2022;4:115-8. https://doi.org/10.25259/IJMSR_40_2021
3. Jo VY. EWSR1 fusions: Ewing sarcoma and beyond. *Cancer Cytopathol* 2020;128:229-31. <https://doi.org/10.1002/cncy.22239>
4. Karlina I, Schroeder BA, Kirgizov K, et al. Latest developments in the pathobiology of Ewing sarcoma. *J Bone Oncol* 2022;35:100440. <https://doi.org/10.1016/j.jbo.2022.100440>
5. Rangankar V, Dhande A, Khaladkar SM, et al. Primary Ewing's sarcoma of the kidney mimicking renal cell carcinoma with widespread metastasis: A case report and a brief review of the literature. *Cureus* 2024;16:e64896. <https://doi.org/10.7759/cureus.64896>
6. Nishant. The diagnostic dilemma of intracranial Ewing's sarcoma: A rare case study. *Biomed J Sci Tech Res* 2024;58. <https://doi.org/10.26717/BJSTR.2024.58.009163>

7. Esch L, Barski D, Bug R, et al. Prostatic sarcoma of the Ewing family in a 33-year-old male - a case report and review of the literature. *Asian J Urol* 2016;3:103-6. <https://doi.org/10.1016/j.ajur.2015.11.007>
8. Abdelghany K, Macaulay R, Tang Z, et al. Pulmonary Ewing sarcoma mimicking small cell lung cancer. *Clin Lung Cancer* 2022;23:e330-4. <https://doi.org/10.1016/j.clcl.2022.04.006>
9. Yoshida A. Ewing and Ewing-like sarcomas: A morphological guide through genetically-defined entities. *Pathol Int* 2023;73:12-26. <https://doi.org/10.1111/pin.13293>
10. Nadal R, Schweizer M, Kryvenko ON, et al. Small cell carcinoma of the prostate. *Nat Rev Urol* 2014;11:213-9. <https://doi.org/10.1038/nrurol.2014.21>
11. Illumina. TruSight RNA Pan-Cancer Panel. Available at: <https://www.illumina.com/products/by-type/clinical-research-products/trusight-rna-pan-cancer.html> (accessed September 21, 2025)
12. Laskar S, Sinha S, Chatterjee A, et al. Radiation therapy dose escalation in unresectable Ewing sarcoma: Final results of a phase 3 randomized controlled trial. *Int J Radiat Oncol Biol Phys* 2022;113:996-1002. <https://doi.org/10.1016/j.ijrobp.2022.04.024>
13. Bacci G, Ferrari S, Longhi A, et al. Local and systemic control in Ewing's sarcoma of the femur treated with chemotherapy, and locally by radiotherapy and/or surgery. *J Bone Joint Surg* 2003;1 85:107-14. <https://doi.org/10.1302/0301-620x.85b1.12781>
14. DuBois SG, Krailo MD, Gebhardt MC, et al. Comparative evaluation of local control strategies in localized Ewing sarcoma of bone: A report from the Children's Oncology Group. *Cancer* 2015;121:467-75. <https://doi.org/10.1002/cncr.29065>
15. Borges da Ponte C, Leitão TP, Miranda M, et al. Prostate Ewing sarcoma/PNET: A case of long survival in a highly aggressive malignancy. *Urology* 2021;154:e11-2. <https://doi.org/10.1016/j.urology.2021.05.006>

CORRESPONDENCE: Dr. Areeb Hassan, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada; mhassan2028@meds.uwo.ca