Dean et al. Routine use of loop diuretics during morcellation does not improve holmium laser enucleation of the prostate outcomes: A prospective, double-blind, randomized controlled trial

APPENDIX

Supplementary Table 1. Subgroup analysis >100 cc prostates			
	Furosemide (n=35)	**Control (n=38)**	**p**
Baseline characteristics			
Age (years) ± SD	72.3±7.0	69.4±6.5	0.091
BMI (kg/m^2) ± SD	28.6±4.0	29.0±4.6	0.683
American Society of Anesthesiologists Score (ASA) ± SD	2.5±0.6	2.6±0.5	0.938
Preoperative prostate volume (cm^3) ± SD	158.5±64.5	158.1±49.4	0.978
Indwelling Foley or CIC (%)	7 (43.8)	6 (28.6)	0.352
Anticoagulation (%)	5 (14.3)	2 (5.3)	0.196
Perioperative characteristics			
Mean total procedure time (min) ± SD	73.8±37.3	72.3±26.8	0.941
Mean enucleation time (min) ± SD	37.2±15.9	41.5±18.7	0.298
Mean morcellation time (min) ± SD	11.5±6.9	16.3±14.0	0.067
Mean specimen weight (g) ± SD	107.9±56.6	120.8±59.8	0.351
Postoperative outcomes			
Same-day TOV (SDTOV) (%)	25 (71.4)	35 (92.1)	**0.021***
Failed SDTOV (%)	3 (12.0)	6 (17.4)	0.590
Same-day discharge (SDD) (%)	21 (63.4)	28 (77.8)	0.201
Mean length of stay (hours) ± (SD)	15.2±16.3	13.6±21.8	0.732
ER-30 day (%)	1 (2.9)	2 (5.4)	0.595
Re-admission 30-day (%)	1 (2.9)	1 (2.6)	0.954
30-day complications: Total (%)	2 (5.9)	4 (10.5)	0.484
TURP syndrome/fluid overload	0 (0)	0 (0)	1.000